Report

Cytogenetic and Genome Research

Cytogenet Genome Res 2015;145:78–179 DOI: 10.1159/000430927 Published online: July 14, 2015

Third Report on Chicken Genes and Chromosomes 2015

Prepared by

Michael Schmid¹ Jacqueline Smith³ David W. Burt³ Bronwen L. Aken⁵ Parker B. Antin²⁶ Alan L. Archibald³ Chris Ashwell²⁹ Perry J. Blackshear³⁰ Clarissa Boschiero⁴⁸ C. Titus Brown^{31, 32} Shane C. Burgess²⁷ Hans H. Cheng³³ William Chow⁶ Derrick J. Coble⁴⁰ Amanda Cooksey²⁸ Richard P.M.A. Crooijmans¹⁷ Joana Damas⁸ Richard V.N. Davis³⁴ Dirk-Jan de Koning¹⁸ Mary E. Delany³⁶ Thomas Derrien²⁰ Takele T. Desta¹⁰ Ian C. Dunn³ Matthew Dunn⁶ Hans Ellegren¹⁹ Lél Eöry³ Ionas Erb²⁴ Marta Farré⁸ Mario Fasold² Damarius Fleming⁴⁰ Paul Flicek⁵ Katie E. Fowler¹¹ Laure Frésard²² David P. Froman³⁸ Valerie Garceau³ Paul P. Gardner^{51, 52} Almas A. Gheyas³ Darren K. Griffin¹² Martien A.M. Groenen¹⁷ Thomas Haaf¹ Olivier Hanotte¹⁰ Alan Hart^{3, 13} Julien Häsler⁷ S. Blair Hedges³⁹ Jana Hertel² Kerstin Howe⁶ Allen Hubbard³⁵ David A. Hume³ Pete Kaiser⁴ Darek Kedra²⁴ Stephen J. Kemp¹⁴ Christophe Klopp²³ Kalmia E. Kniel³⁵ Richard Kuo³ Sandrine Lagarrigue²¹ Susan J. Lamont⁴⁰ Denis M. Larkin⁸ Raman A. Lawal¹⁰ Sarah M. Markland³⁵ Fiona McCarthy²⁸ Heather A. McCormack³ Marla C. McPherson³⁶ Akira Motegi⁵³ Stefan A. Muljo⁴¹ Andrea Münsterberg¹⁵ Rishi Nag⁵ Indrajit Nanda¹ Michael Neuberger⁷ Anne Nitsche² Cedric Notredame²⁴ Harry Noyes¹⁴ Rebecca O'Connor¹² Elizabeth A. O'Hare⁴³ Andrew J. Oler⁴² Sheila C. Ommeh⁴⁹ Helio Pais¹⁶ Michael Persia⁴⁴ Frédérique Pitel²² Likit Preeyanon³¹ Pablo Prieto Barja²⁴ Elizabeth M. Pritchett³⁵ Douglas D. Rhoads⁴⁵ Charmaine M. Robinson³⁷ Michael N. Romanov¹² Max Rothschild⁴⁰ Pierre-François Roux²¹ Carl J. Schmidt³⁵ Alisa-Sophia Schneider¹ Matt Schwartz⁴⁶ Steve M. Searle⁶ Michael A. Skinner⁹ Craig A. Smith⁵⁰ Peter F. Stadler² Tammy E. Steeves⁵¹ Claus Steinlein¹ Liang Sun³⁵ Minoru Takata⁵⁴ Igor Ulitsky²⁵ Qing Wang³⁵ Ying Wang³⁶ Wesley C. Warren²⁷ Jonathan M.D. Wood⁶ David Wragg²² Huaijun Zhou³⁶

¹Department of Human Genetics, University of Würzburg, Würzburg, and ²Bioinformatics Group, Department of Computer Science and Interdisciplinary Centre for Bioinformatics, University of Leipzig, Leipzig, Germany; ³Division of Genetics and Genomics, The Roslin Institute and R(D)SVS, and ⁴Division of Infection and Immunity, The Roslin Institute, University of Edinburgh, Edinburgh, ⁵European Bioinformatics Institute, EMBL, and ⁶Wellcome Trust Sanger Institute, Hinxton, ⁷Medical Research Council Laboratory of Molecular Biology, Cambridge, ⁸Department of Comparative Biomedical Sciences, Royal Veterinary College, University of London, and ⁹Section of Virology, Department of Medicine, Imperial College London, London, ¹⁰School of Life Sciences, University of Nottingham, Nottingham,

Supported by Biotechnology and Biological Sciences Research Council (BBSRC); USDA Agriculture and Food Research Initiative Competitive Grant; NIH Intramural Research Program of the National Institute of Allergy and Infectious Diseases; European Union FP-7 project QUANTOMICS; Deutsche Forschungsgemeinschaft (DFG); Swiss National Science Foundation; Lady Tata Memorial Trust; French 'Agence Nationale de la Recherche' EpiBird; Intramural Research Program of the NIEHS, NIH; Department for International Development (DFID). Correspondence to: Michael Schmid Department of Human Genetics, University of Würzburg Biozentrum, Am Hubland DE–97074 Würzburg (Germany) E-Mail m.schmid@biozentrum.uni-wuerzburg.de

Jacqueline Smith Division of Genetics and Genomics The Roslin Institute, University of Edinburgh Edinburgh EH25 9RG (UK) E-Mail jacqueline.smith@roslin.ed.ac.uk

David W. Burt Division of Genetics and Genomics The Roslin Institute, University of Edinburgh Edinburgh EH25 9RG (UK) E-Mail dave.burt@roslin.ed.ac.uk

KARGER 125

E-Mail karger@karger.com www.karger.com/cgr

© 2015 S. Karger AG, Basel 1424–8581/15/1452–0078\$39.50/0

¹¹ School of Human and Life Sciences, Canterbury Christ Church University, and ¹² School of Biosciences, University of Kent, Canterbury, ¹³Biological Services, Royal Hallamshire Hospital, University of Sheffield, Sheffield, ¹⁴Institute of Integrative Biology, University of Liverpool, Liverpool, ¹⁵ School of Biological Sciences, University of East Anglia, Norwich, and ¹⁶ Weatherall Institute of Molecular Medicine, University of Oxford, John Radcliffe Hospital, Oxford, UK; ¹⁷Animal Breeding and Genomics Centre, Wageningen University, Wageningen, The Netherlands; ¹⁸Department of Animal Breeding and Genetics, Swedish University of Agricultural Sciences, and ¹⁹Department of Evolutionary Biology, Evolutionary Biology Centre, Uppsala University, Uppsala, Sweden; ²⁰Institut de Génétique et Développement de Rennes, Université de Rennes1, and ²¹Equipe Génétique et Génomique, PEGASE, INRA, Rennes, ²²GenPhySE, INRA, and ²³Plateforme Bioinformatique Toulouse Midi-Pyrénées, INRA, Biométrie et Intelligence Artificielle, Castanet-Tolosan, France; ²⁴Comparative Bioinformatics Group, Bioinformatics and Genomics Programme, Centre for Genomic Regulation, Barcelona, Spain; ²⁵Department of Biological Regulation, Weizmann Institute of Science, Rehovot, Israel; ²⁶Department of Cellular and Molecular Medicine, ²⁷College of Agriculture and Life Sciences, and ²⁸ Department of Veterinary Science and Microbiology, University of Arizona, Tucson, Ariz., ²⁹ Department of Poultry Science, Raleigh, N.C., ³⁰Department of Biochemistry, Duke University School of Medicine, NIEHS, Durham, N.C., ³¹Microbiology and Molecular Genetics, Michigan State University, ³²Computer Science and Engineering, Michigan State University, and ³³Avian Disease and Oncology Laboratory, USDA, ARS, East Lansing, Mich., ³⁴Department of Biology, University of Delaware, and ³⁵Department of Animal and Food Sciences, University of Delaware, Newark, Del., ³⁶Department of Animal Science, University of California, Davis, Calif., ³⁷Division of Science and Environmental Policy, California State University Monterey Bay, Seaside, Calif., ³⁸Department of Animal and Rangeland Sciences, Oregon State University, Corvallis, Oreg., ³⁹Center for Biodiversity, Temple University, Philadelphia, Pa., ⁴⁰Department of Animal Science, Iowa State University, Ames, Iowa, ⁴¹National Institute of Allergy and Infectious Diseases, NIH, and ⁴²Medical Science and Computing Inc., High-Throughput Sequencing Bioinformatics Specialist, Computational Biology Section, Bethesda, Md., ⁴³Division of Endocrinology, Diabetes and Nutrition, Department of Medicine, University of Maryland School of Medicine, Baltimore, Md., 44 Virginia Tech University, Department of Animal and Poultry Sciences, Blacksburg, Va., ⁴⁵Department of Biological Sciences, Science and Engineering, University of Arkansas, Fayetteville, Ark., ⁴⁶Department of Genetics, Harvard Medical School, Boston, Mass., and ⁴⁷The Genome Institute, Washington University School of Medicine, St. Louis, Mo., USA: ⁴⁸Departamento de Zootecnia, University of São Paulo, Piracicaba, Brazil; ⁴⁹ Animal Biotechnology Group, Institute of Biotechnology Research, Jomo-Kenyatta University of Agriculture and Technology, Nairobi, Kenya; ⁵⁰Department of Anatomy and Developmental Biology, Monash University, Clayton, Vic., Australia; ⁵¹School of Biological Sciences, University of Canterbury, and ⁵²Biomolecular Interaction Centre, University of Canterbury, Christchurch, New Zealand; ⁵³Department of Radiation Genetics, Graduate School of Medicine, Kyoto University Yoshida Konoe, and ⁵⁴Department of Late Effects Studies, Radiation Biology Center, Kyoto University Yoshida Konoe, Kyoto, Japan

Contributions received from November 4, 2014 to March 19, 2015; accepted for publication by M. Schmid.

The Chicken Leads the Way in Avian Genomics

(Prepared by J. Smith)

Following on from the First Report on Chicken Genes and Chromosomes [Schmid et al., 2000] and the Second Report in 2005 [Schmid et al., 2005], we are pleased to publish this long-awaited Third Report on the latest developments in chicken genomics. The First Report highlighted the availability of genetic and physical maps, while the Second Report was published as the chicken genome sequence was released. This report comes at a time of huge technological advances (particularly in sequencing methodologies) which have allowed us to examine the chicken genome in detail not possible until now. This has also heralded an explosion in avian genomics, with the current availability of more than 48 bird genomes [Zhang G et al., 2014b; Eöry et al., 2015], with many more planned.

Third Report on Chicken Genes and Chromosomes 2015 The chicken has long been a model organism for genetic and developmental studies and now takes its place as a model genome, opening up the fields of phylogenetics and comparative genomics like never before. This report summarizes the current efforts to complete the gaps in the genome and describes the progress of genomic annotation, particularly with respect to noncoding RNAs and genetic variants. Reviews of comparative genomics, avian evolution and sex determination are also given. Transcriptomic case studies are described as are developments in epigenetic studies. We also report on the development of the National Avian Research Facility (http://www.narf. ac.uk/) which is home to various chicken resources including antibodies, chicken lines and transgenic protocols and which hosts information currently available on the sequenced avian genomes.

The Chicken Genome: Current Status of Genome Assembly and Annotations

(Prepared by D.W. Burt, L. Eöry, A.L. Archibald, B.L. Aken, P. Flicek, K. Howe, W. Chow, M. Dunn, J.M.D. Wood, R. Nag, and W.C. Warren)

In 2004, the assembly and annotation of the draft chicken genome assembly was reported [International Chicken Genome Sequencing Consortium, 2004], which represented the first avian (and livestock) genome to be sequenced. The chicken genome assembly has proved an invaluable research tool for the avian research community, and efforts to improve the genome and its annotation have continued.

Current Status

The initial draft of the chicken genome was based on a single female red junglefowl bird (UDC 001) and was assembled using a whole-genome sequencing strategy based on Sanger sequencing (7× coverage), including BAC, fosmid and plasmid paired-end reads. Additional data sources have been incrementally added, including high-resolution genetic linkage maps [Groenen et al., 2009], BAC maps [Ren et al., 2003], radiation hybrid maps [Morisson et al., 2007], targeted genome sequences (gaps and a BAC-based sequence of the Z sex chromosome [Bellott et al., 2010]) and using next-generation sequencing data (14× coverage of 454 and 74× coverage of Illumina reads) [Ye et al., 2011]. In addition, improvements in genome assembly software have increased the quality of these assemblies, for example, removing artefacts due to allelic variants and multigene families [Ye et al., 2011].

The chicken karyotype comprises 39 chromosome pairs as follows: 10 pairs of large autosomes (chromosomes 1-10), 28 pairs of microchromosomes (chromosomes 11-38), and a pair of sex chromosomes (chromosomes W and Z). Chromosome Z is a large chromosome and present as a pair in males. Chromosome W is a microchromosome and present only in the heterogametic (ZW) females [Masabanda et al., 2004]. The current genome assembly, Gallus_gallus-4.0 (Galgal4, GCA_ 000002315.2) covers 1.03 Gb or 96% of the predicted genome size. This includes assembled sequence for all 10 large autosomes, 19 of the 28 microchromosomes, and both sex chromosomes (tables 1, 2). In addition, it includes sequence that is yet to be assigned to an assembled chromosome (linkage groups LGE22C19W28_E50C23 and LGE64) and 14,093 unplaced scaffolds. The contig N50 is 280 kb (n = 27,041), and the scaffold N50 is 12.9

Table 1. Genome statistics for the chicken reference genome assembly

Statistics	
Genome assembly	Galgal4, Nov 2011 (GCA_000002315.2)
Database version	Ensembl 78
Base pairs	1,072,544,763
Golden path length	1,046,932,099
Gene counts	
Coding genes	15,508
Small noncoding genes	1,558
miRNA	1,049
rRNA	29
snoRNA	227
snRNA	79
misc_RNA	150
Mt_rRNA	2
Mt_tRNA	22
Pseudogenes	42
Gene transcripts	17,954
Genescan gene predictions	40,572

Data available from the Ensembl genome database (www.ensembl.org).

Mb (n = 16,847). The contig N50 is defined as the contig length at which 50% of the total number of bases in the assembly are in contigs of that length or greater.

Overlaps with available cDNA clones suggest that less than 5% of coding genes are missing from the final assembly; however, gene duplications and GC-rich sequences are still difficult regions to assemble and pose a particular problem for chicken. Chromosome 16 (which contains the MHC region) and the W sex chromosome are a rich source of duplicated genes and repeats, and are poorly represented. In addition, sequenced contigs have not been assigned to the smallest microchromosomes (GGA29–31 and GGA33–38). Further work is needed to complete the chicken genome by sequencing gaps (so far specific gaps of 18 Mb and 27 Mb have been defined on the ordered and unordered chromosomes, respectively) and missing chromosomes.

The full utility of the chicken genome requires a comprehensive annotation; enumerating and defining all the transcribed and regulatory regions. Until recently, the main tools of gene prediction and annotation for species with few resources (such as full-length cDNA sequences in human and mouse) were based on gene homology with sequences from other species, for example the Ensembl

Assembly unit	Sequence name	INSDC accession	Total length (including gaps)	Present in Galgal4
Chromosome	1	CM000093.3	195,276,750	yes
Chromosome	2	CM000094.3	148,809,762	yes
Chromosome	3	CM000095.3	110,447,801	yes
Chromosome	4	CM000096.3	90,216,835	ves
Chromosome	5	CM000097.3	59,580,361	yes
Chromosome	6	CM000098.3	34,951,654	yes
Chromosome	7	CM000099.3	36,245,040	yes
Chromosome	8	CM000100.3	28,767,244	yes
Chromosome	9	CM000101.3	23,441,680	ves
Chromosome	10	CM000102.3	19,911,089	yes
Chromosome	11	CM000103.3	19,401,079	yes
Chromosome	12	CM000104.3	19,897,011	yes
Chromosome	13	CM000105.3	17,760,035	yes
Chromosome	14	CM000106.3	15,161,805	yes
Chromosome	15	CM000107.3	12,656,803	yes
Chromosome	16 (NOR)	CM000108.3	535,270	yes
Chromosome	17	CM000109.3	10,454,150	yes
Chromosome	18	CM000110.3	11,219,875	yes
Chromosome	19	CM000111.3	9,983,394	yes
Chromosome	20	CM000112.3	14,302,601	yes
Chromosome	21	CM000113.3	6,802,778	yes
Chromosome	22	CM000114.3	4,081,097	yes
Chromosome	23	CM000115.3	5,723,239	yes
Chromosome	24	CM000116.3	6,323,281	yes
Chromosome	25	CM000124.3	2,191,139	yes
Chromosome	26	CM000117.3	5,329,985	yes
Chromosome	27	CM000118.3	5,209,285	yes
Chromosome	28	CM000119.3	4,742,627	yes
Chromosome	29	N/A	N/A	no
Chromosome	30	N/A	N/A	no
Chromosome	31	N/A	N/A	no
Chromosome	32	CM000120.2	1,028	yes
Chromosome	33	N/A	N/A	no
Chromosome	34	N/A	N/A	no
Chromosome	35	N/A	N/A	no
Chromosome	36	N/A	N/A	no
Chromosome	3/	IN/A	IN/A	no
Chromosome	38	N/A	N/A	no
Sex chromosome	VV	CM000121.3	1,248,174	yes
Sex chromosome	L	CM000122.3	82,363,669	yes
Linkage group Linkage group Unplaced scaffolds	LGE22C19W28_E50C23 LGE64	CM000123.3 CM000367.2	965,146 799,899	yes yes yes (14,093 sequences)

Table 3. Summary of chicken RNA-seq expression profiles

Tissue	Transcript models	%
Embryo	13,326	74.2
Heart	10,710	59.7
Testes	14,600	81.3
Bone-derived macrophage	11,339	63.2
Embryo somites	11,350	63.2
DF1 cell line	9,266	51.6
Kidney	10,688	59.5
Breast muscle	9,161	51.0
Brain cerebellum	10,591	59.0
Brain	10,890	60.7
Miscellaneous	11,324	63.1
Total expressed	16,915	94.2
Predicted transcript models	17,954	

gene annotation system [Cunningham et al., 2015]. This approach has been successful at defining protein-coding regions, but not so successful when translated across the large evolutionary distance from mammals to birds, in particular for defining rapidly evolving genes such as innate immune genes or noncoding genes, such as micro-RNAs (miRNAs) and long noncoding RNAs (lncRNAs). Developments in RNA-seq have changed this landscape, capable of both defining and quantifying small and large RNAs, either coding or noncoding [Robertson et al., 2010]. The Ensembl gene annotation pipeline has been extended to incorporate these types of data to improve gene predictions for both coding and noncoding genes [Flicek et al., 2011, 2014].

In 2012, an International Avian RNAseq Consortium [Smith et al., this report] was formed and agreed to share data from a wide range of tissues to be used for gene discovery. Collaboration between this Consortium and Ensembl completed a new annotation of the chicken protein-coding genes (table 1) and the RNA-seq data provided direct experimental evidence for 94% of the predicted transcripts (table 3). All the gene models and RNA-seq data are available to end-users from the Ensembl Genome Browser (www.ensembl.org). This combined evidence-based approach available in Ensembl release 78 predicted 15,508 protein-coding genes in the chicken, which compares with 20,364 (13,990 chicken orthologs, 15,197 human orthologs) and 22,606 (13,741 chicken orthologs, 15,072 mouse orthologs) predictions in human and mouse, respectively. These comparisons suggest that mammals may have ~6,000 protein-coding

genes not found in the chicken. This may be an artefact or a real biological finding, which remains to be tested. Interestingly, it appears that ~274 genes are truly missing in the avian lineage [Lovell et al., 2014], yet are conserved in human, thus potentially serving as naturally evolved models to study the physiological consequence. More limited predictions are available for noncoding RNAs (ncRNAs) in the chicken, which include 1,534 small ncRNA genes in 31 distinct families: 1,049 miRNAs, 29 rRNAs, 227 small nucleolar RNAs (snoRNAs), 79 small nuclear RNAs (snRNAs), and a further 150 (27 families) miscellaneous RNAs. This compares with 9,673 small ncRNAs and 14,817 lncRNAs in human, and 5,860 small ncRNAs and 5,762 lncRNAs in mouse. These comparisons suggest that many more ncRNAs in the chicken are still to be discovered, in particular the lncRNA class. Only 42 pseudogenes were detected in the chicken compared to 14,415 and 8,015 predictions in human and mouse, respectively. The reason for the very low number in the chicken when compared to mammals may be the sequence specificity of reverse transcription by avian LINES [International Chicken Genome Sequencing Consortium, 2004]. Mammalian LINES are more promiscuous and able to retrotranspose most mRNAs.

Ongoing Improvements

To ensure the chicken's continued utility as a model organism, we are undertaking improvements to the chicken genome assembly. Several different approaches are currently being used to further improve the assembly, including single molecule sequencing from PacBio [W. Warren, unpubl. results] to extend reads and fill in gaps, and chromosome sorting [Burt and Fergusson-Smith, unpubl. results] to target sequencing of the tiny chromosomes (GGA29–31 and GGA33–38), which are currently missing from the genome assembly. Microchromosomes are gene dense, and so even these tiny chromosomes are likely to be of significance [Burt, 2002; International Chicken Genome Sequencing Consortium, 2004].

Within the assembled chromosomes, it is important to ensure that the ordering and orientation of the contigs is correct. Optical mapping (http://en.wikipedia.org/wiki/ Optical_mapping) to create high-resolution, chromosome-wide restriction maps based on rare-cutters is used to complement the creation of sequenced contigs [Burt and Dunn, unpubl. results]. Combining these methods will provide accurate data on: (1) confirmation of chromosome assemblies, (2) defining gaps and duplicated regions, (3) assembly of sequence contigs for microchromosomes, and (4) defining a route for completion of the chicken genome assembly. To facilitate the latter and to represent the genome diversity of the chicken, the Chicken Genome Consortium plans to join the Genome Reference Consortium (www.ncbi.nlm.nih.gov/projects/genome/assembly/grc) to share best practice and coordinate activities. Currently, all data associated with the chicken genome project (sequence, BAC-ends, fosmids, BAC clones, etc.) are being evaluated using the gEVAL Genome Browser (http://geval.sanger.ac.uk) to define gaps, errors in assembly, etc. The long-term aim will be to produce a chicken genome assembly of a 'gold' standard to be used as the reference avian genome.

Future Directions

As the chicken genome assembly improves, we will ensure that genome annotation improves and expands. For genome assembly, long read single-molecule sequencing has already shown great future application, with de novo assembled N50 contig length reaching >1 Mb and 90% of the genome assembled in >250-kb length contigs [W. Warren, unpubl. results]. This long-range contiguity promises to achieve the goal of mostly complete copies of each chicken chromosome. In addition, the physical assignment of sequenced microchromosomes will be necessary to form a definition. For gene annotation, PacBio full-length sequences are in progress and will be used to identify missing genes, join gene fragments, define transcription start/end sites and annotate novel transcript isoforms. These annotations will include both coding and noncoding RNA genes. Significant progress has been made in establishing the Bird Gene Nomenclature Committee but more needs to be done, with a greater role for the avian research community [Burt et al., 2009]. Knowledge of gene regulation is a critical component of genomics research. We will pursue methods for identifying transcription factor binding sites, ChIP-seq histone marks, DNA methylation sites and define functional elements, such as enhancers, silencers and promoters (http://www. animalgenome.org/community/FAANG/). As more individual chicken genomes are sequenced, structural and single nucleotide variations will be identified and submitted to the public archives. With a well annotated chicken genome, these genome variants will be associated with functional elements, if any, such as genes or regulatory elements. Quantitative trait loci (QTLs) and genomewide associations are important tools for identifying regions of the genome associated with specific phenotypes. Over 4,300 QTLs have been identified for chicken (http:// www.animalgenome.org/cgi-bin/QTLdb/GG/index). It is anticipated that, with growing variation and expression

Third Report on Chicken Genes and Chromosomes 2015 data, expression QTLs will be identified for chicken in the future. The final goal will be to associate functional variants with specific phenotypes.

The Avian RNAseq Consortium: A Community Effort to Annotate the Chicken Genome

(Prepared by J. Smith, D.W. Burt, and the Avian RNA-seq Consortium)

Publication of the chicken genome sequence in 2004 [International Chicken Genome Sequencing Consortium, 2004] highlighted the beginning of a revolution in avian genomics. Progression of DNA sequencing technologies and data handling capabilities has also meant that genome sequencing and assembly is now a relatively simple, fast and inexpensive procedure. The success seen with the chicken genome was soon followed by the completion of the zebra finch genome [Warren et al., 2010], an important model for neurobiology [Clayton et al., 2009], again based on Sanger sequencing. In recent years, the rapid advances in next-generation sequencing (NGS) technologies, hardware and software have meant that many more genomes can now be sequenced faster and cheaper than ever before [Metzker, 2010]. The first avian genome to be sequenced by NGS methods was the turkey [Dalloul et al., 2010], which was also integrated with genetic and physical maps, thus providing an assembly of high quality, even at the chromosome level. Recently, NGS has been used to sequence the genomes of a further 42 avian species, as part of the G10K initiative [Genome 10K Community of Scientists, 2009]. In addition, there have also been 15 other genome assemblies recently published, each with a focus on a unique aspect of avian biology, including the Japanese quail (domestication) [Kawahara-Miki et al., 2013], Puerto Rican parrot (speciation) [Oleksyk et al., 2012], scarlet macaw (speech, intelligence and longevity) [Seabury et al., 2013], medium and large ground finches (speciation) [Parker et al., 2012; Rands et al., 2013], collared and pied flycatchers (speciation) [Ellegren et al., 2012], peregrine and saker falcons (predatory lifestyle) [Zhan et al., 2013], rock pigeon (domestication) [Shapiro et al., 2013], the ground tit (adaptation to high altitude) [Cai et al., 2013], and the northern bobwhite (population history) [Halley et al., 2014]. Through November 2014, there are currently 56 avian genome sequences completed, either published or in press (table 4). A new project, B10K (web.bioinfodata.org/B10K), proposes sequencing of all avian genomes; this would include all 40 orders, 231 families, 2,268 genera and 10,476

Table 4. Avian	species	with	sequenced	genomes
----------------	---------	------	-----------	---------

Abbreviation	Latin name	Common name	Abbreviation	Latin name	Common name
ACACH	Acanthisitta chloris	rifleman	GALGA	Gallus gallus	chicken
AMAVI	Amazona vittata	Puerto Rican parrot	GAVST	Gavia stellata	red-throated loon
ANAPL	Anas platyrhynchos domestica	Pekin duck	GEOFO	Geospiza fortis	medium ground finch
APAVI	Apaloderma vittatum	bar-tailed trogon	GEOMA	Geospiza magnirostris	large ground finch
APTFO	Aptenodytes forsteri	emperor penguin	HALAL	Haliaeetus albicilla	white-tailed eagle
ARAMA	Ara macao	scarlet macaw	LEPDI	Leptosomus discolor	cuckoo roller
BALRE	Balearica regulorum gibbericeps	grey crowned crane	MANVI	Manacus vitellinus	golden-collared manakin
BUCRH	Buceros rhinoceros silvestris	rhinoceros hornbill	MELGA	Meleagris gallopavo	wild turkey
CALAN	Calypte anna	Anna's hummingbird	MELUN	Melopsittacus undulatus	budgerigar
CAPCA	Caprimulgus carolinensis	Chuck-will's widow	MERNU	Merops nubicus	northern carmine bee-eater
CARCR	Cariama cristata	red-legged seriema	MESUN	Mesitornis unicolor	brown mesite
CATAU	Cathartes aura	turkey vulture	NESNO	Nestor notabilis	kea
CHAPE	Chaetura pelagica	chimney swift	NIPNI	Nipponia nippon	crested ibis
CHAVO	Charadrius vociferus	killdeer	OPHHO	Opisthocomus hoazin	hoatzin
CHLMA	Chlamydotis macqueenii	MacQueen's bustard	PELCR	Pelecanus crispus	Dalmatian pelican
COLLI	Columba livia	rock pigeon	PHACA	Phalacrocorax carbo	great cormorant
COLST	Colius striatus	speckled mousebird	PHALE	Phaethon lepturus	white-tailed tropicbird
COLVI	Colinus virginianus	northern bobwhite	PHORU	Phoenicopterus ruber	American flamingo
CORBR	Corvus brachyrhynchos	American crow	PICPU	Picoides pubescens	downy woodpecker
COTJA	Coturnix japonica	Japanese quail	PODCR	Podiceps cristatus	great crested grebe
CUCCA	Cuculus canorus	common cuckoo	PSEHU	Pseudopodoces humilis	ground tit
EGRGA	Egretta garzetta	little egret	PTEGU	Pterocles gutturalis	yellow-throated sandgrouse
EURHE	Eurypyga helias	sunbittern	PYGAD	Pygoscelis adeliae	Adélie penguin
FALCH	Falco cherrug	saker falcon	STRCA	Struthio camelus	ostrich
FALPE	Falco peregrinus	peregrine falcon	TAEGU	Taeniopygia guttata	zebra finch
FICAL	Ficedula albicollis	collared flycatcher	TAUER	Tauraco erythrolophus	red-crested turaco
FICHY	Ficedula hypoleuca	pied flycatcher	TINGU	Tinamus guttatus	white-throated tinamou
FULGL	Fulmarus glacialis	northern fulmar	TYTAL	Tyto alba	barn owl

species of birds. The chicken genome remains the bestdescribed genome and is used as a reference upon which the annotations of other assemblies are based. Assembly and annotation of the genome continues to improve. However, gaps and unaligned regions remain (particularly for some of the smallest microchromosomes), which can cause practical problems in the analysis and annotation of important loci, especially for those representing gene families. Other approaches, such as long reads generated by Pacific Biosciences (PacBio) sequencing, chromosome sorting and optical maps are being used to resolve these assembly issues [Warren and Burt, pers. commun.]. Specific genome features also require further study; for example, noncoding RNAs (ncRNAs), annotation of rare transcripts, confirmation of alternatively spliced transcripts, mapping of transcription start sites and identification of conserved regions. One method by which some of these goals can be achieved is through analysis of transcriptomic sequence data, or 'RNA-seq' data.

With a view to addressing some of these issues, we decided to collect as much RNA-seq data from the chicken research community as possible. This was the beginning of what we have termed 'The Avian RNAseq Consortium'. Since the start of the Consortium at the end of 2011, it now includes 50 people from 27 different institutions (fig. 1) who have contributed to the effort to create a detailed annotation of the chicken genome by either providing RNA-seq data or by helping to analyze the combined data.

We currently have 21 different data sets (representing more than 1.5 Tb of data) with more data being added (fig. 2; table 5). These data represent transcriptome sequences from many different chicken tissues and from many different experimental conditions, including several infection/disease cases. These data were submitted to public archives, collected at The Roslin Institute and then passed on to the Ensembl team who used the information to help annotate the latest chicken genome assembly, Galgal4 as part of Ensembl release 71 (April 2013) (table 6).



Fig. 1. Worldwide locations of current RNAseq Consortium members.



Fig. 2. Comparison of the different relative amounts of RNA-seq data from each tissue. Tissues from different data providers are shown separately as they have all been subject to different treatments/stimuli. Numbered data sets are as referred to in table 5.

Table 5. Details of RNA-seq data sets

Da	ta set	Description of data	Reads, bp	Sequencing
1	Antin	whole embryo	35	Illumina SE
2	Blackshear	LPS-stimulated macrophages vs. control CEFs	51	Illumina PE
3	Burgess/McCarthy	miRNA from various red junglefowl tissues (adrenal gland, adipose, cerebellum, cere- brum, testis, ovary, heart, hypothalamus, kidney, liver, lung, breast muscle, sciatic nerve, proventriculus, spleen)	50	Illumina SE
4 5 6	Burt/Smith	spleen: infectious bursal disease virus infected vs. control lung and ileum: avian influenza infected vs. control (high path H5N1 and low) lung short-read data	36 36 25	Illumina SE Illumina SE Illumina SE
7	de Koning/Dunn/McCormack	bone from 70-week-old Leghorns	100	Illumina PE
8 9	Frésard/Pitel	brain from epileptic vs. non-epileptic birds pooled whole embryos (stage HH26)	380-400 100	Roche 454 Illumina PE
10	Froman/Rhoads	testes: roosters with high-mobility sperm vs. low-mobility sperm	35	Illumina SE
11	Garceau/Hume	embryo, DF1 cell line and bone marrow-derived macrophages	100	Illumina PE
12	Hanotte/Kemp/Noyes/Ommeh	Newcastle disease virus infection vs. control (trachea and lung epithelial cells)	50	SOLiD SE
13	Häsler/Oler/Muljo/Neuberger	DT40 cells	60	Illumina PE
14	Kaiser	bone marrow-derived dendritic cells from 6-week-old birds (control, DCs + LPS); BMDMs from 6-week-old birds (control, BMDMs + LPS); heterophils isolated from blood of day-old chicks (control, het + LPS)	100	Illumina PE
15	Lagarrigue/Roux	abdominal adipose tissue and liver tissue from 14-week-old broilers	100	Illumina PE
16	Lamont	livers of 8 individuals, 28-day-old broiler males: 4 controls, 4 heat-stressed	100	Illumina SE
17	Munsterberg/Pais	somites injected with anti-mir206 vs. non-injected	50	Illumina PE
18	Schmidt	tissues from heat-stressed and control birds (liver, brain, spleen, thymus, bursa, kidney, ileum, jejunum, duodenum, ovary, heart, breast, monocyte)	42-50	Illumina SE
19	Schwartz/Ulitsky	whole embryo stages – HH4/5; HH11; HH14/15; HH21/22; HH25/26; HH32; HH36 – stranded	80/100	Illumina PE
20	Skinner	CEFs	100	Illumina PE
21	Wang/Zhou	lung from Fayoumi and Leghorn birds: control and H5N3 infected	75	Illumina SE
	PMDMa - Pana mannayy daniyyad	maaranhaaraa CEE - shialaan amhuwa fikushlasta DC - dan duitia salla hat - hatayan hil	L. LDC 1:	om olyroo ook o <i>u</i> i doo

BMDMs = Bone marrow-derived macrophages; CEF = chicken embryo fibroblasts; DC = dendritic cell; het = heterophils; LPS = lipopolysaccharide; PE = paired end; SE = single end.

Table 6. Ensembl 71 annotation statistics of 17,108 genes

Genes	Description	Biotype
15,495	Ensembl	protein-coding
42	Ensembl	pseudogene
2	mt_genbank_import	Mt_rRNA
22	mt_genbank_import	Mt_tRNA
13	mt_genbank_import	protein-coding
1,049	ncRNA	miRNA
150	ncRNA	misc_RNA
29	ncRNA	rRNA
227	ncRNA	snoRNA
79	ncRNA	snRNA

This new annotation includes 15,495 protein-coding genes, 1,049 miRNAs, 456 ncRNAs and 42 pseudogenes. This gene build is primarily concerned with coding genes, but there are many more noncoding genes which remain un-annotated. Consortium members have analyzed the RNA-seq data for long ncRNAs (lncRNAs) [manuscript in preparation], snoRNAs [Gardner et al., 2015] and other features of interest. Around 14,000 potential lncRNA genes have thus far been identified from the RNA-seq data. Ensembl release 71 marked a significant update in the annotation of the chicken genome with gene models based on experimental data. Table 7 shows how this gene

	Ensembl 70	Ensembl 71	Ensembl 77
Assembly	WashUC2, May 2006	Galgal4, Nov 2011	Galgal4, Nov 2011
Base pairs	1,050,947,331	1,072,544,086	1,072,544,763
Coding genes	16,736	15,508	15,508
Short noncoding genes	1,102	1,558	1,558
Pseudogenes	96	42	42
Gene transcripts	23,392	17,954	17,954

Table 7. Comparison	of Ensembl	gene	builds
---------------------	------------	------	--------

build was the first to use the Galgal4 assembly and, through the use of RNA-seq data, was able to help remove assembly errors and reduce the number of predicted gene transcripts by identifying incorrectly predicted genes from previous builds and improving identification of short ncRNAs. The significance of this community effort is indicated by the fact that the current Ensembl 77 gene set has not changed since Ensembl release 71, with only difference being reflected in the total number of base pairs. This is due to the correction of one particular scaffold on the Z chromosome (which was reflected in Ensembl release 74).

The availability of these data will allow for the further development of a chicken expression atlas by providing the ability to analyze transcript levels across tissues (http://geneatlas.arl.arizona.edu/). It will also enable development of exon capture technology for the chicken and has already proved to be of great use in helping annotate the other avian genomes which have now been sequenced. On-going collection of RNA-seq data will remain a valuable resource as genomic analysis of avian species continues to expand.

Methods

Ensembl Gene Build

The chicken gene build from Ensembl release 71 was done using standard Ensembl annotation procedures and pipelines, mostly focussed on protein-coding sequences. Briefly, vertebrate UniProtKB proteins were downloaded and aligned to the Galgal4 (GCA_000002315.2) assembly with Genewise (http://www.ebi.ac.uk/Tools/psa/genewise/) in order to annotate protein-coding models. Uni-Prot assigns protein existence (PE) levels to each of their protein sequences. The PE level indicates the type of evidence that supports the existence of a protein sequence, and can range from PE 1 ('Experimental evidence at protein level') to PE 5 ('Protein uncertain'). Only PE 1 and PE 2 proteins from UniProtKB were used for the Genewise step. RNA-seq models were annotated using the Ensembl RNA-seq pipeline and models from both the Genewise and the RNA-seq pipelines were used as input for the final protein-coding gene set. Chicken cDNAs and RNA-seq models were also used to add UTRs in the 5' and 3' regions. Some missing gene models were recovered by aligning chicken, zebra finch and turkey translations from Ensembl release 65 (December 2011) to the new chicken genome assembly.

RNA-seq Gene Models

Raw reads were aligned to the genome using BWA [Li and Durbin, 2009] to identify regions of the genome that are actively transcribed. The results from all tissues were used to create one set of alignment blocks roughly corresponding to exons. Read pairing information was used to group exons into approximate transcript structures called proto-transcripts. Next, partially mapped reads from both the merged (combined data from all tissue samples) and individual tissues were re-aligned to the proto-transcripts using Exonerate [Slater and Birney, 2005], to create a merged and tissue-specific sets of spliced alignments. For each gene, merged and tissue-specific transcript isoforms were computed from all observed exon-intron combinations, and only the best supported isoform was reported.

Annotation of ncRNAs

The following ncRNA gene types were annotated: ribosomal RNA (rRNA), small nuclear RNA (snRNA), small nucleolar RNA (snoRNA), microRNA precursors (miRNA), miscellaneous other RNA (misc_RNA). Most ncRNA genes in Ensembl are annotated by first aligning genomic sequence against RFAM [Burge et al., 2013], using BLASTN (parameters W = 12, B = 10000, V = 10000, -hspmax 0 -gspmax 0 -kap -cpus = 1), to identify likely ncRNA loci. The BLAST [Altschul et al., 1990] hits are clustered, filtered for hits above 70% coverage, and used

to seed an Infernal [Nawrocki and Eddy, 2013] search with the corresponding RFAM covariance model, to measure the probability that these targets can fold into the structures required. Infernal's cmsearch is used to build ncRNA models. miRNAs are predicted by BLASTN (default parameters) of genomic sequence slices against miRBase [Kozomara and Griffiths-Jones, 2014] sequences. The BLAST hits are clustered, filtered to select the alignment with the lowest p value when more than one sequence aligns at the same genomic position, and the aligned genomic sequence is checked for possible secondary structure using RNAFold [Hofacker et al., 1994]. If evidence is found that the genomic sequence could form a stable hairpin structure, the locus is used to create a miRNA gene model. Transfer RNAs (tRNAs) were annotated as part of the raw compute process using tRNAscan-SE with default parameters [Schattner et al., 2005]. All results for tRNAscan-SE are available through Ensembl; the results are not included in the Ensembl gene set because they are not annotated using the standard evidencebased approach (i.e. by aligning biological sequences to the genome) that is used to annotate other Ensembl gene models.

Summary

The availability of this collection of chicken RNA-seq data within the consortium has allowed:

- annotation of 17,108 chicken genes, 15,495 of which are protein-coding (Ensembl 71)
- identification of ~14,000 putative lncRNA genes (with >23,000 transcripts suggested)
- annotation of miRNAs, snoRNAs, and other ncRNAs
- future generation of an expression atlas which will allow comparisons of expression over many tissues
- an improved avian reference for comparative analyses with 48 other avian genomes [Zhang G et al., 2014b]

Future Directions

The next stage in progressing annotation of the avian genomes will concentrate on the analysis of data generated by PacBio sequencing, in conjunction with stranded RNA-seq data from a wide variety of tissues. PacBio technology allows for very long read lengths, producing reads with average lengths of 4,200–8,500 bp, with the longest reads over 30,000 bp. This enables sequencing of fulllength transcripts. Extremely high accuracy means that de novo assembly of genomes and detection of variants with >99.999% accuracy is possible. Individual molecules can also be sequenced at 99% reliability. The high sensitivity of the method also means that minor variants can be detected even when they have a frequency of <0.1%[http://www.pacificbiosciences.com/products/smrttechnology/smrt-sequencing-advantage/]. We currently have brain transcriptomic PacBio data generated from a female Brown Leghorn J-line chicken [Blyth and Sang, 1960]. This will be analyzed alongside stranded RNA-seq data that has been generated from 21 different tissues. The advantage of using strand-specific sequence information is that it provides an insight into antisense transcripts and their potential role in regulation and strand information of ncRNAs as well as aiding in accurately quantifying overlapping transcripts. It is particularly useful for finding unannotated genes and ncRNAs. This strategy should allow us to obtain full-length transcript sequences, identify novel transcripts and low-level transcripts, map transcription start and stop sites and confirm further ncRNAs.

Get Involved

If you are interested in helping further the annotation of the avian genomes, and you can provide avian RNAseq data or can help with the analysis of such data, then please contact Jacqueline Smith (jacqueline.smith@roslin.ed.ac.uk) or Dave Burt (dave.burt@roslin.ed.ac.uk).

Avian RNAseq Consortium Members

J. Smith, I. Dunn, V. Garceau, D. Hume, P. Kaiser, R. Kuo, H. McCormack, D. Burt (Roslin Institute); A. Cooksey, F. McCarthy, P.B. Antin, S. Burgess (University of Arizona); A. Münsterberg, H. Pais (University of East Anglia); A. Oler (NIH National Institute of Allergy and Infectious Diseases); S. Searle (Wellcome Trust Sanger Institute); P. Flicek, B.L. Aken, R. Nag (European Molecular Biology Laboratory, European Bioinformatics Institute and Wellcome Trust Sanger Institute); C. Schmidt (University of Delaware); C. Klopp (INRA Toulouse); P. Prieta Barja, I. Erb, D. Kedra, C. Notredame (CRG, Barcelona); D. Froman (Oregon State University); D.-J. de Koning (Swedish University of Agricultural Sciences, Uppsala); D. Rhoads (University of Arkansas); I. Ulitsky (Weizmann Institute of Science, Rehovot); J. Häsler, M. Neuberger (in memoriam) (MRC, Cambridge); L. Frésard, F. Pitel (INRA, Auzville); M. Fasold, P. Stadler (University of Leipzig); M. Schwartz (Harvard Medical School); M. Skinner (Imperial College London); O. Hanotte (University of Nottingham); P. Blackshear (NIEHS, North Carolina); S. Lagarrigue, P.-F. Roux (INRA Agrocampus Ouest); T. Derrien (University of Rennes); S. Ommeh (Jomo-Kenyatta University of Agriculture and Technology, Kenya); S. Muljo (NIH NIAID, Bethesda); S.

Kemp, H. Noyes (University of Liverpool); S. Lamont, D. Fleming, D.J. Coble (Iowa State University); Y. Wang, H. Zhou (UC Davis).

Availability of RNA-seq Data

Data have been submitted to the public databases under the following accession numbers:

Antin/Burgess/McCarthy/Schmidt data: BioProject ID: PRJNA204941 (Sequence Read Archive); Blackshear data: PRJEB1406 (European Nucleotide Archive); Burt/ Smith data: E-MTAB-2908, E-MTAB-2909, E-MTAB-2910 (Array Express); de Koning/Dunn/McCormack data: E-MTAB-2737 (Array Express); Frésard/Pitel data: SRP033603 (Sequence Read Archive); Froman/ Rhoads data: BioProject ID: PRJNA247673 (Sequence Read Archive); Garceau/Hume data: E-MTAB-3048 (Array Express); Hanotte/Kemp/Noyes/Ommeh data: E-MTAB-3068 (Array Express); Häsler/Oler/Muljo/Neuberger data: GSE58766 (NCBI GEO); Kaiser data: E-MTAB-2996 (Array Express); Lagarrigue/Roux data: SRP042257 (Sequence Read Archive); Lamont data: GSE51035 (NCBI GEO); Münsterberg/Pais data: GSE58766 (NCBI GEO); Schwartz/Ulitsky data: SRP041863 (Sequence Read Archive); Skinner data: PRJEB7620 (European Nucleotide Archive); Wang/Zhou data: GSM1385570, GSM1385571, GSM1385572, GSM1385573 (NCBI GEO).

Noncoding RNAs in the Chicken Genome

(Prepared by J. Hertel, M. Fasold, A. Nitsche, I. Erb, P. Prieto, D. Kedra, C. Notredame, T.E. Steeves, P.P. Gardner, and P.F. Stadler)

Noncoding RNAs (ncRNAs) are some of the evolutionarily most conserved biomolecules and are essential for many cellular processes [Jeffares et al., 1998]. These include the ribosomal RNAs (rRNAs) and transfer RNAs (tRNAs) involved in translation, the small nucleolar RNAs (snoRNAs) and RNase P/MRP RNAs involved in maturing ncRNAs, and the microRNAs (miRNAs) involved in gene regulation [Cech and Steitz, 2014]. The recently rediscovered and expanded class of 'long noncoding RNAs' (lncRNAs) are involved in diverse functions, including dosage compensation (e.g. Xist, roX1, roX2) and forming scaffolds for recruiting proteins to form functional complexes (e.g. HOTAIR) [Erdmann et al., 2000; Cech and Steitz, 2014].

The First Report on Chicken Genes and Chromosomes 2000 [Schmid et al., 2000] made high-resolution

genetic and physical maps readily available and cemented the chicken as a model genome. However, coverage of ncRNAs was restricted to a brief mention of the Z-linked *MHM* (male hypermethylated) locus, adjacent to the candidate sex-determining gene, *DMRT1* (doublesex and mab-3-related transcription factor 1), subsequently described by Teranishi et al. [2001] and reviewed in The Second Report on Chicken Genes and Chromosomes 2005 [Schmid et al., 2005]. Briefly, the *MHM* region is hypermethylated and transcriptionally silent in males (ZZ), but in females (ZW), *MHM* is hypomethylated and transcribed into an lncRNA suspected to play a role in the downregulation of *DMRT1*. For an up-to-date review of the role of *MHM* in sex determination, see the contribution by Smith et al. in this report.

Compiled prior to the publication of the draft chicken genome [International Chicken Genome Sequencing Consortium, 2004], coverage of ncRNAs in the Second Report on Chicken Genes and Chromosomes 2005 [Schmid et al., 2005] was limited to 3 ncRNA classes, but each was reviewed in detail. Namely, there was extensive coverage of *MHM* (described above), the rRNAs (28S, 18S, 5.8S and 5S) and telomerase RNA. A number of recent studies have since expanded the number of chicken ncRNAs. These include miRNAs [Glazov et al., 2008; Shao et al., 2012], lncRNAs [Chodroff et al., 2010; Necsulea et al., 2014], and the entire complement of ncRNAs [International Chicken Genome Sequencing Consortium, 2004; Zhang et al., 2009; Gardner et al., 2015].

The ncRNAs reviewed in this contribution fall into 3 major classes: (1) housekeeping RNAs such as rRNAs, tRNAs, spliceosomal RNAs, and snoRNAs as well as a few minor classes, (2) miRNAs and related small RNAs, and (3) lncRNAs.

Housekeeping RNAs

Ribosomal RNAs. The rRNA operon comprising the 18S, 5.8S and 28S rRNAs is located in more than 100 copies on a microchromosome [Muscarella et al., 1985]. The pol-III transcribed 5S rRNA is organized as 2 tandem repeats on chromosome 9 [Daniels and Delany, 2003]. In addition, the mitogenome contains its own copy of the LSU and SSU rRNA [International Chicken Genome Sequencing Consortium, 2004].

Transfer RNAs. The predicted number of tRNAs encoded in the chicken genome is relatively modest according to tRNAscan-SE [Lowe and Eddy, 1997]. Just 299 tRNA genes are predicted, which is just 4 more than in *Saccharomyces cerevisiae* [Chan and Lowe, 2009]. All 20 canonical tRNAs are represented with copy numbers

Table 8. Number of ncRNAs identified in the chicken genome to date and the proportion confirmed by RNA-seq experiments

RNA family	Homology-based predictions, n	Confirmed by RNA-seq, n
tRNAs	300 ^a	280 (93.3%)
rRNAs	22	10 (45.5%)
microRNA	427	280 (65.6%)
snoRNA box C/D	106	90 (84.9%)
snoRNA box H/ACA	68	48 (70.6%)
snoRNA – Cajal body	12	12 (100%)
spliceosomal snRNAs	77	35 (45.4%)

ranging from 6 (Trp) to 36 (Ala), there is 1 selenocysteine tRNA and 4 pseudogenized tRNAs. There are also several significant tRNA clusters. Namely, 10 Cys tRNAs are clustered in a 5-kb region on chromosome 27, 10 Arg and Tyr tRNAs are clustered in a 40-kb region on chromosome 2, and 8 Glu, Leu, Lys, and Val tRNAs can be found in a 4-kb region on chromosome 16.

Spliceosomal RNAs. The chicken genome encodes both a major and a minor spliceosome. The components of the major spliceosome (U1, U2, U4, U5, and U6) are present in multiple copies (many of which are presumably pseudogenes) as in most vertebrates, while most sn-RNAs of the minor spliceosome (U4atac, U6atac, U11, and U12) are single-copy genes [Marz et al., 2008].

Other Nuclear RNAs. There are single copies of the RNase P and RNase MRP RNAs on chromosomes 8 and 4, respectively; 4 copies of the RNA component of the signal recognition particle in a 1-Mb region on chromosome 5, and a cluster comprising 3 distinct Y RNAs in a 17-kb region on chromosome 2. Of the 2 vault RNA loci (chromosomes 5 and 12), the one on chromosome 5 appears to be a pseudogene. The telomerase RNA and the U7 snRNA are also represented as single copies.

Small Nucleolar RNAs. Covariance models of known snoRNAs were used to identify snoRNAs in chicken using INFERNAL. In addition, an independent homology-based approach using snoStrip was applied. Both results were merged leaving 106 box C/D and 68 box H/ACA snoRNAs, of which 90 (84.9%) and 48 (70.6%) were confirmed with RNA-seq, respectively.

miRNAs and Related Small ncRNAs

An INFERNAL search for miRNAs using those mi-RNA families from miRBase that have at least 1 representative in vertebrates returned 427 miRNAs. Of these, 280 genes are confirmed with RNA-seq. Only 51 miRNAs annotated in chicken seem to be species-specific, since they do not show any significant homology in other vertebrates. Only 18 of these miRNAs are organized in 8 families, while the remaining 33 miRNAs come as single sequences with no paralogs at all. Although miRNAs are lost infrequently in animal genomes, it appears that 3 miRNA families (i.e. mir-139, mir-425 and mir-1287) have been lost completely in chicken. Before the divergence of chicken, a few other miRNAs have also been lost (e.g. 8 at the branch leading to the split of Sauropsida and 4 at the ancestor of Dinosauria/Aves).

The recent availability of comprehensive strand-specific high-throughput RNA sequencing data for several developmental stages and tissues allows for further experimental confirmation of the current catalog of chicken ncRNAs. Only few traces of small ncRNA are found in total RNA libraries due to size selection for larger fragments. In small RNA-seq protocols, the size fraction of 17–25 nt is selected for library preparation. The resulting sequencing data comprises not only processed miRNA sequences, for which those protocols were originally developed, but contains also fragments from snoRNAs, tRNAs and other housekeeping RNAs. Table 8 shows the fraction of major ncRNA families that could be experimentally validated [Gardner et al., 2015].

lncRNAs

While many classes of short RNAs have been characterized in recent years, lncRNAs are less well understood. They are operationally defined as noncoding transcripts that are longer than 200 bp. Here, we only describe lncRNAs falling in intergenic regions, thus excluding intronic transcripts or those overlapping a coding transcript. There is still much debate about whether lncRNAs constitute mainly transcriptional noise or in their majority have functions such as reviewed in Rinn and Chang [2012]. Our approach revealed a high degree of sequence conservation in other bird species as well as reptiles, thus hinting at a large percentage of functional transcripts.

RNA-seq data from 20 different tissues comprising 238 read libraries was mapped to the Ensembl 71 chicken genome. Ab-initio prediction of transcripts was then performed using the software Cufflinks [Trapnell et al., 2010]. After merging predictions from individual libraries, only intergenic transcripts (at least 1 kb away from coding genes) were kept. An evaluation of coding potential as well as repeat content was used to extensively filter

Table 9. Number of lncRNAs identified in the chicken genome to date along with those remaining after application of different approaches to increase the likelihood that these are functional

Category	Number
Ab-initio predicted transcripts	10,364
Conserved beyond chicken	9,386
Conserved in at least 10 species	5,058
Conserved in at least 40 species	1,251
Expressed by at least 0.1 RPKM in 1 of 123 libraries	8,345
Expressed by at least 0.1 RPKM on average in at least 1 of 9	
tissues	6,360
Of these, expressed in only 1 tissue	3,475
Of these, expressed in all 9 tissues	318

the transcripts thus obtained. To further cut down on false positives, only multi-exonic transcripts were retained. Systematic phylogenomic profiling against 52 other bird and reptile species using the procedure described in Derrien et al. [2012] was performed to reveal the degree of conservation of the spliced sequences. Transcript expression was studied on a reduced set of 9 tissues (comprising 123 libraries) with a sufficient number of replicates (between 4 and 36) each.

We report 10,364 putative lncRNA transcripts forming 7,597 gene loci. Less than 10% of these transcripts appear to be chicken-specific, almost half of them are conserved in at least 10 species, and more than 10% are conserved in at least 40 species (table 9). Expression of ~80% of transcripts could be detected in at least one of the selected tissues. Generally we found a strong bias toward tissue-specific transcripts: more than half of the expressed transcripts are expressed in a single tissue, and only ~5% are found ubiquitously expressed in all tissues.

Synopsis

The ncRNAs in the chicken genome form a valuable collection for a number of comparative analyses as the number of ncRNA-derived pseudogenes in the avian genomes is remarkably low [International Chicken Genome Sequencing Consortium 2004; Gardner et al., 2015]. This, therefore, forms a useful way to discriminate pseudogenized ncRNAs from functional copies in related genomes where synteny is likely to have been preserved. A further use for the chicken ncRNA annotations is the identification of 'missing' ncRNAs. The chicken genome has the most complete collection of microchromosome sequences currently available, and has proven essential to explain the consistent 'loss' of essential ncRNAs in the avian lineage such that these have been traced to the poor

ability of sequencing methods for capturing the microchromosomes [Gardner et al., 2015]. As more transcriptomes from more bird species, from more developmental stages and from more tissue types become available, new ncRNAs will undoubtedly be located. However, identifying the function of these novel ncRNAs will remain a major research challenge.

Genome Sequencing in Birds and Evolutionary Inferences from Avian Genome Sequences (Prepared by H. Ellegren)

Developments in avian genome analysis well illustrate the overall progress in genome sequencing of non-model species offered by next-generation sequencing (NGS) technology [Ellegren, 2014]. It took 6 years between the reports of the first 2 avian genome sequences in 2004 (chicken, Gallus gallus [International Chicken Genome Sequencing Consortium, 2004]) and 2010 (zebra finch, Taeniopygia guttata [Warren et al., 2010]), respectively, both sequenced with Sanger technology. In 2010, the genome sequence of turkey (Meleagris gallopavo) was also reported, the first based on NGS [Dalloul et al., 2010]. Subsequently, there has been an exponential increase in the number of avian genomes reported, recently topping by the parallel sequencing and analysis of 45 new genomes by the Avian Phylogenomics Consortium [Jarvis et al., 2014]. By now (January 2015), a total of some 60 bird species have been subject to genome sequencing (table 10). The precise number is somewhat dependent on what criteria for coverage, sequence continuity and other assembly statistics one sets for defining a genome to have been sequenced. Moreover, as new genomes are continuously reported, the total number changes quickly. All 34 currently recognized orders of living birds in the avian tree of life are represented in the list of sequenced species. This offers an unprecedented resource to comparative and evolutionary studies, some brief examples of which will be provided here and limited to work that specifically has made use of wholegenome sequence data. For a general review of avian evolutionary genomics, see Ellegren [2013].

Molecular Evolution

From alignments of 2 or more avian genomes, it is possible to analyze a number of aspects related to the accumulation of nucleotide substitutions and the forces driving this process. For neutral sites, the rate of sequence divergence should reflect the underlying rate of mutation. Under the assumption that 4-fold degenerate sites evolve

Latin name	Species name	Reference	Latin name	Species name	Reference
Acanthisitta chloris	rifleman	Jarvis et al., 2014	Gavia stellata	red-throated loon	Jarvis et al., 2014
Amazona vittata	Puero Rican amazon	Oleksyk et al., 2012	Geospiza fortis	medium ground-finch	Zhang G et al., 2012
Anas platyrhynchos	Pekin duck	Huang et al., 2013	Geospiza magnirostris	large ground finch	Rands et al., 2013
Antrostomus carolinensis	Chuck-will's-widow	Jarvis et al., 2014	Haliaeetus albicilla	white-tailed eagle	Jarvis et al., 2014
Apaloderma vittatum	bar-tailed trogon	Jarvis et al., 2014	Haliaeetus leucocephalus	bald eagle	Jarvis et al., 2014
Aptenodytes forsteri	emperor penguin	Li C et al., 2014	Leptosomus discolor	cuckoo-roller	Jarvis et al., 2014
Aquila chrysaetos	golden eagle	Doyle et al., 2014	Lyrurus tetrix	black grouse	Wang et al., 2014
Ara macao	scarlet macaw	Seabury et al., 2013	Manacus vitellinus	golden-collared manakin	Jarvis et al., 2014
Balearica regulorum	grey crowned crane	Jarvis et al., 2014	Meleagris gallopavo	turkey	Dalloul et al., 2010
Buceros rhinoceros	rhinoceros hornbill	Jarvis et al., 2014	Melopsittacus undulatus	budgerigar	Jarvis et al., 2014
Calypte anna	Anna's hummingbird	Jarvis et al., 2014	Merops nubicus	carmine bee-eater	Jarvis et al., 2014
Cariama cristata	red-legged seriema	Jarvis et al., 2014	Mesitornis unicolor	brown mesite	Jarvis et al., 2014
Cathartes aura	turkey vulture	Jarvis et al., 2014	Nestor notabilis	kea	Jarvis et al., 2014
Chaetura pelagica	chimney swift	Jarvis et al., 2014	Nipponia nippon	crested ibis	Jarvis et al., 2014
Charadrius vociferus	killdeer	Jarvis et al., 2014	Opisthocomus hoazin	hoatzin	Jarvis et al., 2014
Chlamydotis macqueenii	MacQueen's bustard	Jarvis et al., 2014	Pelecanus crispus	Dalmatian pelican	Jarvis et al., 2014
Colinus virginianus	northern bobwhite	Halley et al., 2014	Phaethon lepturus	white-tailed tropicbird	Jarvis et al., 2014
Colius striatus	speckled mousebird	Jarvis et al., 2014	Phalacrocorax carbo	great cormorant	Jarvis et al., 2014
Columba livia	pigeon	Shapiro et al., 2013	Phoenicopterus ruber	American flamingo	Jarvis et al., 2014
Corvus brachyrhynchos	American crow	Jarvis et al., 2014	Picoides pubescens	downy woodpecker	Jarvis et al., 2014
Corvus cornix	hooded crow	Poelstra et al., 2014	Podiceps cristatus	great crested grebe	Jarvis et al., 2014
Coturnix japonica	Japanese quail	Kawahara-Miki et al., 2013	Pseudopodoces humilis	ground tit	Cai et al., 2013; Qu et al., 2013
Cuculus canorus	common cuckoo	Jarvis et al., 2014	Pterocles gutturalis	yellow-throated sandgrouse	Jarvis et al., 2014
Egretta garzetta	little egret	Jarvis et al., 2014	Pygoscelis adeliae	Adélie penguin	Li C et al., 2014
Eurypyga helias	sunbittern	Jarvis et al., 2014	Serinus canaria	canary	Frankl-Vilches et al., 2015
Falco cherrug	saker falcon	Zhan et al., 2013	Struthio camelus	ostrich	Jarvis et al., 2014
Falco peregrinus	peregrine falcon	Zhan et al., 2013	Taeniopygia guttata	zebra finch	Warren et al., 2010
Ficedula albicollis	collared flycatcher	Ellegren et al., 2012; Smeds et al., 2015	Tauraco erythrolophus	red-crested turaco	Jarvis et al., 2014
Fulmarus glacialis	northern fulmar	Jarvis et al., 2014	Tinamus guttatus	white-throated tinamou	Jarvis et al., 2014
Gallus gallus	chicken	ICGSC, 2004	Tyto alba	barn owl	Jarvis et al., 2014

neutrally, Nam et al. [2010] estimated the neutral divergence rate at $1.2-2.2 \times 10^{-9}$ per site in 3 avian lineages (the ancestral bird lineage and the lineages leading to chicken and zebra finch, respectively), with evidence for rate variation among lineages. Zhang G et al. [2014b] found a mean rate of 1.9×10^{-9} at 4-fold degenerate sites across the species sequenced by the Avian Phylogenomics Consortium, again with significant variation among lineages. Intriguingly, they also found a positive correlation between rate of divergence and number of species per order, suggesting an association (though not necessarily causative) between divergence and diversification.

Selection in coding sequences has often been studied by estimating the ratio of the non-synonymous (d_N) and the synonymous (d_S) substitution rates (i.e. d_N/d_S). For example, the approach can be used to identify fast and adaptively evolving genes with codons or whole sequences showing $d_N/d_S > 1$. Functional annotation of genes based on gene ontology has indicated that certain categories of genes may be overrepresented among adaptively evolving genes in different bird lineages [Nam et al., 2010; Huang et al., 2013; Zhan et al., 2013; Li C et al., 2014; Zhang G et al., 2014b]. d_N/d_S can also be used to study

DOI: 10.1159/000430927

how the intensity of selection varies among lineages in relation to life history [Weber et al., 2014b] as well as how it varies within genomes. For example, it has been suggested that the high rate of recombination in microchromosomes reduces the effect of Hill-Robertson interference, leading to lowered d_N/d_S estimates under the assumption that slightly deleterious mutations contribute to d_N [Nam et al., 2010; Zhang G et al., 2014b]. The rationale here is that selection is more efficient in the removal of slightly deleterious mutations when recombination uncouples loci and reduces the effects (interference) of linked selection. However, recombination is also related to the process of GC-biased gene conversion in avian genomes [Weber et al., 2014a], and this can easily lead to erroneous conclusions on selection in coding sequences [Ratnakumar et al., 2010]. There is clear evidence that the recombination hot-spot in the pseudoautosomal region of avian sex chromosomes has a strong effect on sequence evolution [Smeds et al., 2014].

Genome Evolution

It is well known that birds have the smallest genomes among amniotes. Genome assemblies are in the range of 1.0–1.3 Gb, although cytogenetic analyses indicate that there might be species with somewhat larger genomes. The relative compactness of avian genomes is evident from shorter introns and less intergenic DNA than in mammals and reptiles. It seems largely to be due to a lower transposable element activity in the avian lineage but also to a high rate of deletions [International Chicken Genome Sequencing Consortium, 2004; Zhang G et al., 2014b], which might be driven by high recombination rates in avian chromosomes [Nam and Ellegren, 2012].

That the avian karyotype is unusually stable is also well established; the majority of bird species have a diploid chromosome number of 2n = 76-80. Genome sequence data now allow for more quantitative assessments of this stability and, in particular, analyses of intrachromosomal rearrangements not detected in previous cytogenetic work or in work based on low-resolution linkage maps. Kawakami et al. [2014] estimated the rate of inversion at 1.5 and 2.0 events per million years in 2 song bird lineages, respectively, with a mean inversion size of ~7 Mb. Across more distantly related groups of birds, a mean rate of 1.25 rearrangement events (mostly inversions) per million years was observed, with increased rate seen in some lineages [Romanov et al., 2014; Zhang G et al., 2014b]. This is considerably lower than that seen in mammals, particularly in rodents [Zhao and Bourque, 2009]. Karyotypic stability coupled with a low repeat activity is also manifested in highly similar amounts of assembled DNA per syntenic chromosome of divergent avian lineages (table 11).

The access to whole-genome sequences provides new possibilities to elucidate the character and extent of evolutionary conservation in avian genomes. Sequences that are conserved across distantly related species are likely to be subject to functional constraints and to evolve under purifying selection. The proportion of constrained sequence has been estimated to 7.5% [Zhang G et al., 2014b], which is at least twice as high as the proportion of exonic sequence. This demonstrates that protein-coding genes are not the only, maybe not even the dominating, functionally important category of sequences in avian genomes. Sequences involved in regulation of gene expression are likely to constitute a large part of the conserved elements, some of which are conserved across both birds and mammals.

Demography

DNA sequence data is often used in attempts to infer the demography of species and to test different demographic scenarios and estimate divergence times. A major limitation of traditional methods for such inference has

Table 11. Amount of assembled sequence in syntenic chromosomes 1–28 of 3 bird species in which scaffolds have been assigned to and ordered along chromosomes through genetic linkage analysis

Chromosome	Assembled sequence, Mb				
	collared flycatcher	zebra finch	chicken		
1 ^a	119.8	119.6	201.0		
1A ^a	74.8	73.7	-		
2	157.4	156.4	154.9		
3	115.7	112.6	113.7		
4A ^b	21.2	69.8	94.2		
4 ^b	70.3	20.7	_		
5	64.6	62.4	62.2		
6	37.2	36.3	37.4		
7	39.3	39.8	38.4		
8	32.0	28.0	30.7		
9	26.8	27.2	25.6		
10	21.3	20.8	22.6		
11	21.7	21.4	21.9		
12	21.9	21.6	20.5		
13	18.6	17.0	18.9		
14	17.4	16.4	15.8		
15	14.9	14.4	13.0		
16 ^c	-	< 0.01	0.43		
17	12.4	11.6	11.2		
18	13.1	11.2	10.9		
19	11.9	11.6	9.9		
20	15.6	15.7	14.0		
21	8.1	6.0	7.0		
22	5.7	3.4	3.9		
23	7.9	6.2	6.0		
24	8.0	8.0	6.4		
25	2.7	1.3	2.0		
26	7.6	4.9	5.1		
27	5.5	4.6	4.8		
28	6.1	5.0	4.5		
Zď	59.7	74.6	72.9		

^a Chicken chromosome 1 corresponds to chromosomes 1 and 1A in flycatcher and zebra finch, the result of a fission in the galliform lineage.

^b Chicken chromosome 4 corresponds to chromosomes 4 and 4A in zebra finch and flycatcher, the result of a fission in the passeriform (flycatcher and zebra finch) lineage.

^c Chromosome 16 is for some reason very difficult to sequence in birds.

^d The smaller size of the Z chromosome in flycatcher owes mainly to a relatively high proportion of scaffolds that could not be ordered with confidence on the Z chromosome by linkage analysis. been the inability to model changes in effective population size through time. However, this has come to change by the introduction of the pairwise sequentially Markovian coalescent (PSMC) model that infers local time to the most recent common ancestor for each region of the genome and uses coalescent theory to deduce the effective population size (N_e, basically the number of reproducing individuals, at equilibrium) at different points in time [Li and Durbin, 2011]. The method, in its original form, requires the access of a diploid genome sequence where sequencing has been done at sufficient depth (approx. >15×) that the vast majority of heterozygous sites have been correctly called.

PSMC-based estimates of ancestral population sizes have recently been presented for several bird species and have given us the first detailed insights into how bird populations have varied in size through time [Nadachowska-Brzyska et al., 2015]. The crested ibis (*Nipponia nippon*), now one of the most rare and endangered species on earth, was once common and had an N_e of ~70,000 one million years ago [Li S et al., 2014]. It subsequently went through 2 population bottlenecks, reducing N_e to 10,000 about 10,000 years ago. During modern times, the decline has continued.

The Adélie (*Pygoscelis adeliae*) and emperor (*Apteno-dytes forsteri*) penguins showed steady increases in N_e during the period between 1 million and 100,000 years ago, from <100,000 to 200,000 and 600,000, respectively [Li C et al., 2014]. Except for a decrease to 400,000 in the Adélie penguin about 60,000 years ago, the population sizes of the 2 penguins seem to have been stable during the last 100,000 years. Higher N_e estimates have been obtained for the collared flycatcher (*Ficedula albicollis*). It increased from 500,000 five million years ago to 1,500,000 about 200,000 years ago [Nadachowska-Brzyska et al., 2013]. Then, probably like many species in the temperate part of the world, the population decreased in size to again reach about 500,000 during the last glaciation period.

Final Remarks

It is not unlikely that the majority of bird species in not too many years will have been subject to genome sequencing. This will provide a most comprehensive portray of the genetic diversity within the class Aves and allow us to pinpoint the genetic changes that underlie specific adaptations in different bird lineages. It will help us addressing topics such as convergence and what types of mutations are essential for phenotypic novelties. Genome resequencing of multiple individuals within species will also be important for characterization of avian biodiversity. It probably has its greatest potential when it comes to illuminating the processes that govern avian evolution, like local adaptation, speciation and diversification.

The Use of Avian BAC Libraries and Clones

(Prepared by M.N. Romanov and D.K. Griffin)

High-density gridded libraries of large-insert clones using bacterial artificial chromosome (BAC) and other vectors are essential tools for genetic and genomic research in chicken and other avian species. Earliest chicken BAC libraries [Zoorob et al., 1996; Zimmer and Verrinder Gibbins, 1997; Crooijmans et al., 2000] were applied to solve a range of problems at both genome-wide and chromosomal levels. Uses of BAC libraries and individual clones include physical and comparative mapping, support of whole-genome sequencing projects, positional cloning of quantitative trait loci, and isolation of genes and other genomic regions of interest [e.g. Lee et al., 2003; Ren et al., 2003; Wallis et al., 2004; Huang et al., 2006; Reed et al., 2008; Blagoveshchenskiĭ et al., 2011]. In early studies, BACs were also used as one of the sources for the development of microsatellite markers to characterize genetic diversity in populations [e.g. Morisson et al., 1998; Romanov and Weigend, 1999; Crooijmans et al., 2000]. For the purposes of BAC library screening and cross-species BAC-based physical mapping, one of the most costand time-effective methods is the overgo-based filter hybridization technique [Romanov et al., 2003; Romanov and Dodgson, 2006]. Pools of short sequence-specific overgo probes facilitate integration of linkage and physical maps by high-throughput assignment of genes and markers to BAC contigs.

BAC clones are the probes of choice in FISH mapping for the development of chromosome-level physical maps of genomes and (along with chromosome paints) for comparative cytogenetic mapping. Given that defining avian karyotypes (and hence whole genome assemblies) is impeded by the presence of a large number of nearindistinguishable microchromosomes, development of microchromosome-specific BAC probes has proved to be very useful in microchromosome identification [Masabanda et al., 2004; Griffin et al., 2008; Skinner et al., 2009; Völker et al., 2010], fine mapping, and characterization of inter- and intrachromosomal rearrangements [e.g. Fillon et al., 1998; Romanov et al., 2005; Griffin et al., 2008; Skinner et al., 2009; Völker et al., 2010; Lithgow

Table 12. Major avian BAC libraries showing approximate number of clones, insert size and coverage [after Romanov et al., 2009a, with amendments]

Bird	Library	Clone DB ^a abbreviation	No. of clones	No. of clones in Clone DB ^a	Average insert, kb	Coverage	Reference
Duck (Anas platyrhynchos)	China Agriculture University	-	84,480	-	117.94	9.84	Yuan et al., 2006
Golden pheasant (Chrysolophus pictus)	Zhejiang University reverse-4D	-	89,600	-	106.87	7.421	Ye et al., 2012
Emu (Dromaius novaehollandiae)	VMRC-16 (LIBGSS_011154 ^b)	VMRC16	133,632	-	165	13.5	Kellner et al., 2005
California condor (Gymnogyps californianus)	CHORI-262 ^c	CH262	89,665	-	-	14	Romanov et al., 2006
Chicken (Gallus gallus)	CHORI-261 ^c Texas A&M 031-JF256-BI unspecified ^d Texas A&M 032-JF256-RI Texas A&M 033-JF256-H3 Texas A&M Wageningen 020-CHK-H3 (LIBGSS_009949 ^b) Tohoku University China Agriculture University LIBGSS_003202, LIBGSS_009895 ^b	CH261 TAM31 XXbac ^d TAM32 TAM33 WAG	73,000 38,400 - 38,400 38,400 49,920 49,152 138,240	66,486 108 10 10,792 66 8,611 - -	195 (182 ^a) 150 - 152 171 130 (134 ^a) 149 118	11 5.2 - 5.3 6 5.4 (5.57 ^a) 3.2 13.34	Romanov et al., 2003 Lee et al., 2003 Crooijmans et al., 2000 Hori et al., 2000 Liu et al., 2003
Turkey (Meleagris gallopavo)	CHORI-260 ^c (LIBGSS_010330 ^b) TKNMI	CH260 TKNMI	71,000	10,438 9,519	190 160	11.1	Romanov and Dodgson, 2006 Zhang et al., 2011
Budgerigar (Melopsittacus undulatus)	CHORI-263 ^c	CH263	197,392	-	-	-	CHORI ^c
Crested ibis (Nipponia nippon)	Zhejiang University 4D-PCR Zhejiang University reverse-4D		129,312 1,040,000		86.5 100.9	7.8 35	Lan et al., 2014
Zebra finch (Taeniopygia guttata)	TG_Ba TGMCBa	TG_Ba (TG_Ba) TGMCBa	147,456	205 132,650	-	-	Kellner et al., 2005 Thomas et al., 2008
White-throated sparrow (Zonotrichia albicollis)	CHORI-264 ^c (LIBGSS_011743 ^b)	CH264	196,354	1,872	144	21.1	Romanov et al., 2009b

^a According to information in the NCBI Clone DB (http://www.ncbi.nlm.nih.gov/clone).

^b Library name in the NCBI GSS database (http://www.ncbi.nlm.nih.gov/nucgss).

^c As listed in the Library Resources, BACPAC Resources Center, CHORI, Oakland, Calif. (http://bacpac.chori.org/libraries.php).

^d A subset of the Texas A&M 031-JF256-BI (TAM31) chicken library as deposited in the NCBI Clone DB (http://www.ncbi.nlm.nih.gov/clone).

et al., 2014]. Indeed, the generation of cross-species FISH data, along with bioinformatic analysis of genome sequences, is crucial in understanding the gross genome evolution of avian species and, in particular, reconstructing ancestral karyotypes [e.g. Zhang et al., 2011; Romanov et al., 2014].

In addition to chicken, BAC libraries have been constructed and characterized for several other bird species including representatives of Anseriformes, Casuariiformes, Cathartiformes, Galliformes, Psittaciformes, Pelecaniformes, and Passeriformes. Most of these are publicly available (table 12), enabling BAC-based applications in other important research areas. These include structural analyses of the MHC chromosome (microchromosome 16 in chicken), loci and alleles in chicken [Solinhac et al., 2010; Suzuki et al., 2012], turkey [Chaves et al., 2007; Bauer and Reed, 2011] and golden pheasant [Ye et al., 2012]. These studies have contributed to a more detailed exploring of distinctive organization, function and evolution of the bird MHC, showing conserved synteny of a single microchromosome involved in the immune system in Aves and variation in MHC alleles related to disease resistance/susceptibility.

Comparative screening of BAC libraries and FISH using BAC clones have also assisted in developing genomic resources and tools for the endangered California condor and the white-throated sparrow, a model for behavioral studies in humans [Romanov et al., 2006, 2009a, b, 2011; Thomas et al., 2008; Modi et al., 2009]. Kothapalli et al. [2011] examined California condor BACs harboring the immunoglobulin lambda locus and found a region of high homology to the chicken and zebra finch orthologs. In another highly endangered bird, the crested ibis, 2 genomic BAC libraries were created to target and characterize a cluster of defensin genes known to be a critical component of the innate immune system [Lan et al., 2014]. Moreover, by generating a zebra finch library TG_Ba, Luo et al. [2006] reported a genomic region that was previously difficult to sequence and involved the first bird androgen receptor gene and a conserved regulatory element.

The emu BAC library and clones have proved to be pivotal tools for investigating the structure, organization and evolution of the sex chromosomes in ratite, reptilian and other bird species [Chapus and Edwards, 2009; Janes et al., 2009, 2011]. Indeed, a gene *SubA*, with orthologs being expressed in mammalian gonads, has been isolated from the emu library and assigned to a pair of microchromosomes using BAC-FISH [Janes et al., 2008].

Taken together, these studies demonstrate that applications of large-insert clones and BAC libraries derived from birds are, and will continue to be, effective tools to aid high-throughput and state-of-the-art genomic efforts and the important biological insight that arises from them.

Comparative Genomics

(Prepared by D.M. Larkin, M. Farré, and J. Damas)

Even after the chicken genome sequencing and assembly completion in 2004 [International Chicken Genome Sequencing Consortium, 2004], the avian comparative genome studies were not as advanced as studies of mammal genomes due to a very limited number of sequenced avian genomes, with only 3 of them published by 2013 [International Chicken Genome Sequencing Consortium, 2004; Dalloul et al., 2010; Warren et al., 2010]. In contrast, more than 50 mammalian genomes were published or sequenced by the end of the same year. Because of these unfortunate limitations and a lack of ordered physical maps for avian genomes, until recently, comparative genome studies of avian species were mostly focused on the comparison of karyotype evolution achieved through either a direct DNA-to-DNA hybridization (FISH) [Griffin et al., 2007] or sequence comparison of avian and mammalian genomes [Larkin et al., 2009]. Despite the limited resolution of FISH, these studies provided important insights into the unique structure and evolutionary stability of avian karyotypes, and revealed major patterns of early amniote genome evolution [Griffin et al., 2007; Larkin et al., 2009].

The situation changed drastically in 2014 when de novo sequencing of 45 and parallel comparative studies of 48 avian genomes were published in Science and other journals [e.g. Whitney et al., 2014; Zhang G et al., 2014b]. This became possible due to the Avian Genome Consortium, with members spread across 13 countries. The sequencing of 20 genomes was performed at Beijing Genome Institute with a high level of sequence coverage $(>50\times)$, which resulted in high N50 and long scaffolds for 19 of them; while the additional 25 genomes were sequenced at lower (24-32×) coverage [Zhang G et al., 2014a]. A major difference of this 'flock of genomes' from the vertebrate genomes sequenced and assembled in a traditional way was that the majority of the species still had no physical maps that could be used to verify the quality of the assemblies and anchor scaffolds along chromosomes [Lewin et al., 2009]. Several exceptions are the genomes of chicken, turkey and zebra finch, published previously and the assemblies of budgerigar and ostrich assisted by optical maps [Griffin et al., 2008; Warren et al., 2010; Zhang G et al., 2014a].

Due to the highly fragmented nature of the majority (25 out of 45) of the new avian genome assemblies, most comparative analyses involving this dataset were focused on gene or short sequence feature aspects of avian genome evolution. These comparisons provided new insights into the avian genome evolution, phylogeny, and unique adaptations formed in the avian lineage. Some important aspects of avian chromosome evolution were also uncovered; they are described in the cytogenetic section of this report.

Genome Size Reduction

Among amniotes, birds have the smallest genome size ranging from 0.91 Gb in the black-chinned hummingbird to 1.3 Gb in the common ostrich [International Chicken Genome Sequencing Consortium, 2004]. The small size of avian genomes is often associated with adaptations related to flight (e.g. higher rates of oxidative metabolism) [Hughes and Hughes, 1995]; however, there is evidence that genome size reduction could have already occurred in the dinosaur ancestor of all birds [Organ et al., 2007]. One of the striking differences between the genomes of mammals and birds, which accounts for the smaller genome sizes in birds, is a lower fraction of transposable elements (TEs) and other repetitive sequences in avian genomes compared to those of mammals. In birds, TEs comprise 4-10% of the genome length in most species, while in mammals, TE fractions range from 35 to 52% [Zhang G et al., 2014b]. Among the genomes analyzed by

the Avian Genome Consortium, only the downy woodpecker had a large fraction of TEs (22%) resulting from either a species- or lineage-specific expansion of LINE type CR1 (long interspersed elements; chicken repeat 1) transposons [Zhang G et al., 2014b]. Compared to nonavian reptiles, birds have on average $19 \times less$ SINEs (short interspersed repeat elements), suggesting that this reduction could have happened in the common ancestor of all birds [Zhang G et al., 2014b]. Endogenous viral elements are also rare in avian genomes. Their fraction is $6-13 \times$ lower in avian than in mammalian genome sequences, suggesting that birds either are able to suppress viral DNA insertion in their genomes, or (less likely) are not subjected to viral infections to the same level as mammals [Cui et al., 2014].

In addition to a lower fraction of TEs, the comparative analysis of avian genomes demonstrated that the proteincoding genes and intergenic regions are shorter in birds compared to mammalian and non-avian reptile counterparts [Zhang G et al., 2014b]. Most likely the shorter gene length in avian genomes is a consequence of a lower fraction of TEs in introns and intergenic intervals. A similar compression of the gene and intergenic region length was observed in bats, the only flying mammal [Zhang G et al., 2013]. In both cases (birds and bats), this could be related to the necessity of fast gene regulation associated with flight [Zhang G et al., 2013, 2014b].

Interestingly, avian genomes have a large number of ancestral microdeletions absent from the genomes of other reptiles. Larger-scale structural rearrangements were also found associated with segmental deletion of 1,559 genes in birds, most of which had at least one additional paralog in the lizard or human genomes [Lovell et al., 2014]. These missed genes are not randomly distributed across amniote chromosomes, but tend to cluster into syntenic blocks with the highest fraction of missed genes located in lizard chromosome 2 (human chromosome 19) [Lovell et al., 2014]. About 90% of the genes missed in avian genomes are present in the crocodile genome suggesting that the loss occurred in the lineage leading from the archosaurian to the avian ancestor [Lovell et al., 2014]. This is the highest level of segmental loss of genes found in vertebrates so far. The fact that the majority of lost avian genes are members of gene families suggests that the remaining members of the same gene family in the avian genomes could have compensated for the loss of paralogs.

Chromosome Structure

While a detailed review of the karyotype evolution of birds is covered in the cytogenetic section of this report,

here it is worth mentioning several aspects of this feature. A macro-synteny comparison of 6 avian genomes assembled to scaffolds with N50 > 10 Mb and the remaining genomes with N50 > 2 Mb [Zhang G et al., 2014a] confirmed previous observations based on cytogenetic studies that avian karyotypes are highly conserved [Burt et al., 1999]. An alignment of all avian genomes against chicken chromosome sequences is available from the Evolution Highway Comparative Chromosome Browser [Larkin et al., 2009] (fig. 3). Very few interchromosomal rearrangements were found in the majority of avian species confirming previous observations that the structure of avian chromosomes is stable [Romanov et al., 2014]. A micro-synteny analysis of homologous genes among avian genomes confirms the hypothesis of evolutionary stability of avian genomes, indicating that a higher fraction of genes share the same neighbours in avian genomes than in different mammals [Zhang G et al., 2014b]. However, it was observed that all vocal learning species had significantly more rearranged chromosomes than the closely related non-vocal learning species [Zhang G et al., 2014b]. This could be related to unique adaptations of the vocal learners or alternatively to a larger radiation of highly successful vocal learning clades compared to other avian groups [Zhang G et al., 2014b]. The former hypothesis is supported by findings of Whitney et al. [2014] who report 10 songbird-specific genes with the majority of them being located in the regions of synteny disruption between the chicken and zebra finch genomes. Among these genes, they identified two (YTHDC2L1 and TMRA) that are highly expressed in the vocal learningassociated nuclei of the zebra finch brain [Whitney et al., 2014].

Nucleotide and Gene Evolution

Comparative analysis of avian DNA sequences demonstrated that the nucleotide substitution rates in avian genomes are generally lower than those in mammals [Zhang G et al., 2014b]. However, large interordinal differences in the rates were detected between distinct avian groups. It was found that the substitution rate correlates with the number of species in a clade. For example Passeriformes, the most specious avian order, had a neutral nucleotide substitution rate 2 times higher than any other neoavian species on average. Interestingly, significant differences in the substitution rates were also observed when comparing all landbirds as a group to waterbirds, consistent with observations that landbirds have higher diversification rates [Zhang G et al., 2014b]. Overall, these observations suggest that differences in nucleotide

GGA1:100K	GGA1:500K
Turkéy Pekin duck Rock pigeon Common cockoo Chinney swift Anna's hummingbird Hoatzin Adelie penguin Crested ibis Little egret Downy woodpecker Peregrine falcon Budgerigar Golden-collared manakin American crow Zebra finch Medium ground finch Chinese alligator Western-painted turtle Boa snake Anole lizard Opossum	Turkey Pekin duck Rock pigeon Common cockoo Chimney swift Anna's hummingbird Hoatzin Killdeer Emperor penguin Adelie penguin Crested ibis Little egret Downy woodpecker Peregrine falcon Budgerigar Golden-collared manakin American crow Zebra finch Medium ground finch Chinese alligator Western-painted turtle Boa snake Anole lizard Opossum
1 1	1 1

substitution rates between avian groups could be related to the levels of diversification and the differences in adaptive pressures the diverse groups were subjected to.

In agreement with the hypothesis of evolutionary stasis in avian genomes (compared to other vertebrates and mammals particularly) [Burt et al., 1999], orthologous genomic intervals in bird genomes were found to contain more constrained sequences than orthologous regions of mammalian genomes. It was shown that 7.5% of the avian genome on average is found in sequences that evolve slower than at the neutral substitution rate estimated for birds [Zhang G et al., 2014b]. Only ~12.5% of these sequences were associated with protein-coding genes, while the majority of other sequences were intergenic or located in introns. These sequences are candidates for regulatory elements in avian genomes and often are sites for transcription regulatory factors [Woolfe et al., 2005]. This compares to 2.2% of the human and mouse genome sequences that are constrained in both species after divergence from their common ancestor [Rands et al., 2014]. Interestingly, the avian-specific constrained noncoding sequences are enriched for transcription factors functioning in metabolism, while the ancestral amniote-specific elements (shared by birds and mammals) are associated with factors contributing to signal regulation, stimulus responses and development [Zhang G et al., 2014b]. This indicates that the regulation of genes responsible for metabolism could have changed during the evolution of the ancestral avian genome.

The availability of a large number of avian genomes distributed among the whole avian phylogenetic tree allowed for the first time the identification of signatures of convergent evolution in avian genomes. For example, signatures of selection related to vocal learning, a phenotype that appeared in the avian evolution 3 times in different lineages [Zhang G et al., 2014b], were detected comparing the sequences of protein-coding orthologous genes of vocal learning and non-learning species. Out of 7,909 genes present in all vocal-learning and control non-learning species' genomes, about 200 genes contained signatures of convergent accelerated evolution in vocal learners. Strikingly, 73% of these genes were expressed in the song-

Fig. 3. Syntenic fragments and homologous synteny blocks (HSBs) identified in chicken chromosome 1 (GGA1) at 2 resolutions (100 and 500 kb). Blue and red blocks define syntenic fragments in target genomes in '+' and '-' orientation, respectively, compared to the chicken chromosome, with target species scaffold or chromosome numbers indicated inside the blocks. Only the rows with genomes assembled to chromosomes (turkey, duck, zebra finch, Anole liz-

bird brain with the vast majority of these being expressed in song-learning nuclei [Whitney et al., 2014]. The analysis of accelerated evolution of non-coding sequences revealed 822 accelerated elements shared by all 3 vocal learning groups. Of these elements, 322 were associated with 278 genes with a high proportion (192) expressed in song nuclei. Overall, there was a 2.0–3.5-fold enrichment in accelerated evolution associated with genes expressed in vocal learning brain regions compared to other tissues [Zhang G et al., 2014b].

Adaptive Phenotypes

Comparative analysis of avian genomes shed light on the evolution of avian-specific adaptations. The leading one - ability to fly - requires several major skeleton and bone modifications: bones must be light and at the same time strong. In birds and other tetrapods this was achieved through the reduction of the number of bones and their pneumatization [Zhang G et al., 2014b]. Out of 89 genes related to ossification in birds, 49 showed evidences of positive selection, which is twice as high as in mammals. The highest ratio of non-synonymous to synonymous substitutions (d_N/d_S) was detected for the genes AHSG and P2RX7 related to bone mineral density and bone homeostasis, respectively [Zhang G et al., 2014b]. An increased metabolism related to flight adaptations requires highly-efficient gas exchange in the lungs. Birds have evolved a constant volume lung and a rigid trunk region. Five genes related to the mammalian lung development were apparently lost in birds. The development of feathers (composed of α - and β-keratin proteins) is associated with an almost 2-fold increase in the number of β -keratin genes in birds compared to reptiles, while the a-keratin gene family was significantly contracted in avian species [Zhang G et al., 2014b].

An important phenotypic feature of birds that finally was explained through the comparative genome analysis of avian genomes is the lack of teeth or 'edentulism'. This phenotype has evolved independently in several animal lineages. Fossil records show that several extinct avian lineages had teeth [O'Connor and Chiappe, 2011]. Therefore, edentulism could have evolved independently in avian evolution [Meredith et al., 2014]. A search of avian

ard, and opossum) contain complete HSBs, while blocks in the remaining rows represent either HSBs or syntenic fragments. Evolutionary breakpoint regions are defined as white intervals in between either 2 adjacent syntenic fragments originating from the same scaffold in a target genome or 2 adjacent HSBs. A complete set of all chicken chromosomes compared to other avian and non-avian genomes is available from http://eh-demo.ncsa.uiuc.edu/birds/. genomes for the traces of tooth-related genes recovered multiple pseudogene fossils of enamel and dentin genes with multiple frame-shift or exon-deletion mutations. The majority of these mutations differ in diverse avian genomes, suggesting that they occurred independently in evolution. However, all birds analyzed by the Avian Genome Consortium shared the same deletions in 4 enamel genes (*ENAM*, *AMELX*, *AMTN*, and *MMP20*) and 1 dentin-related gene (*DSPP*) suggesting that the common ancestor of all birds likely had no mineralized teeth [Meredith et al., 2014].

Another amazing feature of birds is their most advanced vertebrate visual system. They exhibit the ability to distinguish colors over a wider range of wavelengths than mammals. Unlike mammals, birds likely have retained the ancestral tetrapod set of cones [Zhang G et al., 2014b]. For the majority of vertebrate visual opsin genes, birds had a higher number of copies compared to mammals. The number of opsin gene classes (4) found in most birds suggests that birds are most likely tetrachromatic [Zhang G et al., 2014b], with the exception of penguins who had only 3 classes of opsin genes [Li C et al., 2014] suggesting a 3-chromatic vision, consistent with earlier observations in aquatic mammals who also lost 1 or 2 of cone pigments [Newman and Robinson, 2005].

In conclusion, the comparative analysis of 48 avian genomes proved to be a powerful tool to reveal multiple signatures of genome adaptations related to avian ability to fly. The evolutionary stability of avian karyotypes is likely related to the reduction of transposable and other repetitive sequences in avian genomes. Avian-specific segmental deletions of gene paralogs together with shorter genes and intergenic regions made gene regulation fast and energy-efficient. The skeleton modifications that resulted in a smaller number of light-weight bones were accompanied by accelerated evolution of genes involved in ossification. Avian genomes tend to show relatively small variation in regulatory gene sequences compared to mammals reflecting the high degree of adaptation and specialization of bird genomes, probably inherited from their dinosaur ancestor.

Avian Cytogenetics Goes Functional

(Prepared by D.K. Griffin, M.N. Romanov, R. O'Connor, K.E. Fowler, and D.M. Larkin)

It is now over 10 years since the first avian genome [International Chicken Genome Sequencing Consortium, 2004] and the first complete avian karyotype [Masabanda et al., 2004] were both published; however, until 2014, avian cytogenetics has focused heavily on descriptive studies [e.g. Griffin et al., 2007, 2008; Skinner et al., 2009; Völker et al., 2010] with less attention to its functional relevance. Last year, however, saw 2 landmark efforts in the chromosomal studies of birds: a special issue of Chromosome Research in April and the announcement of recently completed sequences of multiple new avian genomes in Science and the BMC journals (taking the total number sequenced to over 50) in December. Studying the chromosomes of birds is, perhaps for the first time, telling us more about avian biology, function and evolution than it ever has.

What Do We Know So Far? Karyotypic Stability

The near-unique nature of the avian karyotype has remained a consistently reported feature of bird biology since the first chromosome preparations were made. Although many animal groups have microchromosomes, the small size and abundant number of chromosomes in avian species set birds apart genomically from other vertebrate groups. To the best of our knowledge, there are over 1,000 published avian karyotypes, most comprehensively summarized by Christidis [1990], with several hundred added since this review. All of these karyotypes are partial however, with usually only 5-10 pairs of chromosomes easily distinguished, and the rest homogeneously classified. Moreover, the vast majority of karyotypes hardly differ from each another, with rare exceptions including the stone curlew (*Burhinus oedicnemus*; 2n = 40), the beach thick knee (*Esacus magnirostris*; 2n = 40), several hornbills (2n = 42), kingfishers and hoopoes (Upupa *epops*; 2n > 120) at each end of the numerical spectrum [Christidis, 1990]. Indeed, even since the advent of zoo-FISH, the identification of an interchromosomal rearrangement in a bird is a relatively uncommon event [Griffin et al., 2007].

Central to our understanding of avian biology and evolution is establishing the reasons *why* avian karyotypes are evidently so stable. Clues to such an enquiry might lie in those rare exceptions to the rule. For instance, the Falconiformes (falcons, etc.) and Psittaciformes (parrots, etc.) have noticeably undergone numerous evolutionary changes. Moreover, it is noteworthy that when interchromosomal change occurs, it tends to recur. The best example of this is a fusion of the ancestral chromosomes 4 and 10; an event that appears to have occurred independently throughout evolution in chicken (*Gallus gallus*), greylag goose (*Anser anser*), collared dove (*Streptopelia decaocto*) and probably other species also [Griffin et al., 2007]. In this review, we



Fig. 4. FISH of 2 BACs for chicken microchromosome 19 (green, p arm; red, q arm) to peregrine falcon (*Falco peregrinus*) chromosomes. A fusion to a falcon macrochromosome is apparent [our unpubl. data].

examine some of the latest tools and preliminary solutions that are being used to understand the underlying mechanisms that lead to chromosome rearrangements in birds (and in eukaryotes in general).

If we accept that interchromosomal change occurs only rarely in birds, then it is reasonable to assume that this happens usually only when there is an adaptive value to doing so. In most species, phenotypic diversity is usually associated with wholesale changes in karyotype structure. Aves as a phylogenetic class underwent a series of rapid speciation events beginning ~65 million years ago (Mya) and ending ~50 Mya. Chromosomal change is usually a cause or consequence of speciation (i.e. a species barrier), but until recently, the microchromosomes that constitute the majority of the avian karyotype, have not been amenable to study. The latest studies, however, have paved the way for a flurry of research activity that not only describes the avian karyotype in more detail, but might also provide functional clues as to its nature.

New Molecular Cytogenetic Tools

Lithgow et al. [2014] produced a set of chromosome paints and bacterial artificial chromosomes (BACs) that will start the process of characterizing the microchromosomes and their changes over evolutionary time. They reported the development of chicken microchromosomal paint pools and generation of pairs of specific microchromosome BAC clones with some success in zoo-FISH experiments. For instance, they detected a fusion of the ancestral chicken chromosome 23 orthologue to a macrochromosome in gyrfalcon (*Falco rusticolus*). A FISH image of BACs hybridized to peregrine falcon (*Falco peregrinus*) chromosomes [unpubl. data] is shown in figure 4.

Third Report on Chicken Genes and Chromosomes 2015 McPherson et al. [2014] examined the Japanese quail (Coturnix japonica). Comparing chicken and turkey BAC clones on mitotic and meiotic chromosomes, they demonstrated that high-resolution FISH is practicable. Ishishita et al. [2014] also assessed the distribution of centromeric repetitive sequences on both micro- and macrochromosomes. It is therefore now possible to achieve full, high-resolution characterization of all avian chromosomes in all species studied, including the elusive chromosome 16 and the D-group (smallest) chromosomes. There are several current strategies to fill the gaps; one of these is by the use of PacBio, a novel single-molecule realtime sequencing platform, targeting the sequence of smaller chromosomes using sorted chromosome preps, and assembling contigs into scaffolds and super-scaffolds from optical maps [Ganapathy et al., 2014].

What Have Sequence Assemblies Taught Us?

The progress of genome assembly in birds has been slow in comparison to other animal groups such as mammals. Following chicken [International Chicken Genome Sequencing Consortium, 2004], it took a further 6 years until the second and third avian genome sequences were published, namely those of the zebra finch (Taeniopygia guttata, a model for neurological function, especially learned vocalization) [Warren et al., 2010] and turkey (Meleagris gallopavo) [Dalloul et al., 2010]. More recently, the Pekin duck (Anas platyrhynchos) [Huang et al., 2013] was added along with 2 falcon species (Falco peregrinus and F. cherrug) [Zhan et al., 2013] and many others (table 4). The availability of these assembled genomes provided the opportunity for comparative genomics at a chromosomal level. In 2010, we made the first comparison of 2 species using genome assembly information from the macrochromosomes [Völker et al., 2010]. A similar comparison more recently was made in chicken compared to duck [Rao et al., 2012], and then a 3-way comparison (allowing studies of the direction of change) in chicken, turkey and zebra finch [Skinner and Griffin, 2012; Lithgow et al., 2014]. The principal features of chromosomal change in birds are homologous synteny blocks (HSBs), which are demarked by evolutionary breakpoint regions (EBRs). While analyzing these features, some general patterns have started to emerge. The first is that, although interchromosomal change is rare, intrachromosomal changes are commonplace. Breakpoint reuse is also commonplace, significantly more so than in mammals, and there is some evidence of an association between chromosomal breakage and non-allelic homologous recombination (NAHR) [Völker et al., 2010].

Zhang G et al. [2014b] used a whole-genome shotgun strategy to generate new whole-genome sequences from 45 bird species representing many of the major clades and at least 1 representative from over 90% of all avian orders. Around 20 species had a high (50× or greater) coverage and these were the subjects of further cytogenetic studies. These included the common ostrich (Struthio camelus) and the budgerigar (Melopsittacus undulatus), which were further assembled using data from optical mapping experiments [Ganapathy et al., 2014]. This had the effect of significantly increasing the assembly's N50 scaffold sizes to around 15 Mb, and these were subsequently used with those already assembled by chromosome (chicken, turkey, zebra finch, and duck). Romanov et al. [2014] made use of novel whole-genome sequence information from 21 avian genome sequences available on an interactive browser (Evolution Highway). By focusing on the 6 best-assembled genomes (chicken, turkey, duck, zebra finch, ostrich, and budgerigar), a putative karyotype of the avian ancestor (probably a bipedal feathered dinosaur) was assembled for each chromosome. The evolutionary events were reconstructed that led to each of the 6 species' genome organization. Intra- and interchromosomal changes appear best explained most parsimoniously by a series of inversions and translocations with common breakpoint reuse. Microchromosomes represent conserved blocks of synteny in most of the 21 species, and a series of interchromosomal changes in the ostrich were also described that would not have been predicted by karyotype analysis alone. These results suggest that mechanisms exist to preserve a static overall avian karyotype/genomic structure, including the microchromosomes, with rare interchromosomal change (e.g. in ostrich and budgerigar lineages); this is discussed in depth in the next section. Of the species examined, it seemed that chicken had the least number of chromosomal rearrangements compared to the dinosaur ancestor. From Evolution Highway it is also possible to assess rates of chromosomal evolution in birds. Zhang G et al. [2014b] suggest that birds have a lower chromosomal rearrangement rate than mammals but nonetheless can undergo 'bursts' of rearrangement, e.g. during the evolution of vocal learning. This finding corroborates those of Romanov et al. [2014] that identified the zebra finch and budgerigar as the 2 species with the most chromosomal rearrangements from the avian ancestor.

If we accept that chicken and its galliform relatives underwent the least number of chromosomal changes whilst diverging from the ancestral bird, we also must consider whether they also have undergone the fewest phenotypic changes. In other words: is the dinosaur avian ancestor more like a land fowl than any other bird? The most ancient near-certain fossil representative of modern birds (Neornithes) was almost certainly aquatic (for example, *Vegavis*, a genus of birds from the Late Cretaceous epoch) and has been identified as a Galloanseres. Indeed, the earliest known bird-like creatures in the fossil record (e.g. the Ornithurae Gansus) were either fully aquatic or at least amphibious, and it has been suggested that, due to the fact that they had webbed feet (as well as other traits), they were more like ducks [Romanov et al., 2014]. On the other hand, most authors agree that the dinosaur ancestors of birds were terrestrial, feathered, bipedal, relatively small and with limited flying ability - not unlike a chicken. At best we can determine therefore, the ancestral birds were most likely more phenotypically associated with the Galloanseres, and the confusion of whether they were more akin to water- or land fowl may be due to interpretations based on depositional sampling biases, limited understanding of functional anatomy, and whether the individuals that have been discovered are actually fully representative of the groups to which they belonged. Chromosomal evidence provides an independent record of the functional material of inheritance in living birds and, as such, can complement a fossil record that is always likely to be incomplete.

Of all species studied so far, it seems clear that the rearrangement of chromosomes is non-random [Pevzner and Tesler, 2003; Larkin et al., 2009]. The reasons for this non-random nature warrant deeper investigation. According to mammalian evidence, evolutionarily conserved HSBs appear to evolve in different ways from the dynamic and ever-changing EBRs; whether this is true of birds remains to be seen. In mammals, chromosomal breakpoints are correlated with sequences of segmentally duplicated or repetitive DNA [Bovine Genome Sequencing and Analysis Consortium et al., 2009; Larkin et al., 2009; Groenen et al., 2012; Ruiz-Herrera et al., 2012], and species-specific EBRs are correlated with regions enriched for transposable elements (TEs) [Bovine Genome Sequencing and Analysis Consortium et al., 2009; Groenen et al., 2012]. In mammals, EBRs and HSBs largely contain genes with notably different functional ontologies, e.g. organismal development in HSBs [Larkin et al., 2009] and lineage-specific biology and adaptive features in EBRs [Bovine Genome Sequencing and Analysis Consortium et al., 2009; Larkin et al., 2009; Groenen et al., 2012]. It has been suggested therefore that chromosome rearrangements and the respective gene ontologies con-

tained within HSBs and EBRs help to explain lineage-specific phenotypes in mammals. Mammalian and avian genomes are very different however (not least because of the interchromosomal stability of avian genomes), and thus the question remains about whether the patterns that have been observed in mammals will apply to birds also. Birds have less repetitive DNA through the elimination of repetitive sequences [International Chicken Genome Sequencing Consortium, 2004; Shedlock, 2006; Zhang G et al., 2014b] so that the avian genome is constrained by size, primarily because of gene loss as well as lineage-specific erosion of repetitive elements and large segmental deletions. In addition to their karyotypic stability, bird genomes also have a very high degree of evolutionary stasis at nucleotide sequence and gene synteny levels. Nonetheless, one of the key findings was the detection of nonneutral evolutionary changes in functional genes as well as non-coding regions. Many of these changes coincide with adaptations to different lifestyles and niches and display homoplasy [Zhang G et al., 2014b].

The non-random nature of chromosome rearrangement in birds, the reasons for the apparent interchromosomal (but not intrachromosomal) stability of avian karyotypes (see next section), the role of TEs and NAHR, the relationship to phenotype, the question of whether spatial organization of ancestral gene networks is maintained in bird and other reptile lineages, and the question of whether lineage-specific EBRs alter gene order in networks that had adaptive value, all require further investigation. Harnessing the data from over 50 avian genomes (undoubtedly with many more on the way) and employing tools such as Evolution Highway will give us unprecedented insight into avian chromosome evolution and its relationship to avian biology.

Why Is the Avian Karyotype Structure Conserved Inter- but Not Intrachromosomally?

Burt's 'fission-fusion' hypothesis suggested that most avian microchromosomes became fixed in the common dinosaur ancestor with a karyotype of $2n \approx 60$ including 20 microchromosome pairs [Burt, 2002]. The remainder, including the smallest, probably was created by further fission. Romanov et al. [2014] suggested that a basic pattern of 2n = 80 (~30 microchromosome pairs) was fixed before the Palaeognathae-Neognathae divergence 100 Mya. The subsequent paucity of intermicrochromosomal rearrangements between most Neognathae indicates an evolutionary advantage either to retaining this pattern or a lack of opportunity for change. For instance, an explanation for such evolutionary stasis might be that the underlying mutational mechanisms of chromosomal changes are fundamentally different in birds compared to other amniotes through a lack of adaptive value, rather than purifying selection, slowing down the rate of change. Much of this could be explained, in part, by a paucity of copy number variants (CNVs; including segmental duplications), recombination hotspots, TEs and/or endogenous retroviruses; however, this would not explain why interchromosomal change is rare but intrachromosomal change is common, particularly in groups that have undergone rapid speciation such as Passeriformes.

The rate of chromosome rearrangement (and subsequent speciation) depends on: (1) the mutation rate and (2) the fixation rate [Burt et al., 1999]. The first of these is related to the frequency of homologous sites [Burt, 2002]. Repeat structures in general (e.g. CNVs), and TEs in particular, provide substrates for chromosomal rearrangement. In a genome constrained by size, the opportunity for mutation is reduced and only fission (or intrachromosomal change, e.g. inversion) can occur. This provides an explanation why (1) avian genomes are more fragmented than any other vertebrate (birds have the most chromosomes) and (2) why there have been fewer interchromosomal rearrangements. There might also be advantages to retaining multiple chromosomes in a karyotype through the generation of variation, the driver of natural selection. That is, a karyotype with more chromosomes leads to a greater number of genetic variants that the gametes produce and an increase in recombination rate due to the fact that there needs to be at least 1 obligatory chiasma per chromosome. Burt [2002] proposed that a higher recombination rate has also led to the features that we most associate with microchromosomes (high GC content, low repeats, high gene density, etc.) and resulted in the formation and fixation of the archetypal avian karyotype with both macro- and microchromosomes and little interchromosomal rearrangement. Such a constraint, however, does not preclude rearrangement within the individual chromosomes. Romanov et al. [2014] and King [1995] argue that an increase in intrachromosomal rearrangement correlates with bursts of speciation in birds, perhaps mediated by an increase in localized repeat content.

Some birds nonetheless have a significantly different karyotype from the standard $2n \approx 80$. This can occur within one closely related group, e.g. Adélie penguin (*Py-goscelis adeliae*; 2n = 96) and the emperor penguin (*Ap-tenodytes forsteri*; 2n = 72) (but both associated with high degrees of intermicrochromosomal rearrangement), thereby suggesting that similar mechanisms can both re-

duce or increase chromosome number in relatively short time frames. Comparisons of chromosomal change in the zebra finch and the budgerigar suggest that rearrangement rates are similarly high in both groups to which they belong (Passeriformes and Psittaciformes, respectively) but that the latter is capable of fixing interchromosomal rearrangements, while the former is not. The mechanisms underpinning these differences are, as yet, unknown, but studies of the gene ontology terms of speciesspecific EBRs might provide clues. As more avian genomes with better assemblies are analyzed, this may indicate adaptive phenotypic features associated with specific gene ontologies typical of individual orders, families or genera.

The Sex Chromosomes

Worthy of especial consideration is the conserved sex chromosome ZW system that is present in all birds apart from the Palaeognathae. Their independent origin from the XY system does not escape the fact that similar mechanisms appear to have run in parallel, for instance genes on the Z chromosome (like the mammalian X) have undergone selection for male-advantage functions. Like the Y chromosome, the W is small (albeit medium-sized by avian standards), heterochromatic and gene poor. Graves [2014] suggests that the W chromosome is at a more advanced stage of differentiation than the Y chromosome as it has accumulated more LINEs and lost more genes during its evolution. Pokorná et al. [2014] considered multiple sex chromosomes and meiotic drive in a range of amniotes. This study noted that the single ZW system in birds contrasts with that of other reptile and amniote groups; they raised a very exciting hypothesis that this contrast may possibly be related to the differential involvement of sex-specific sex chromosomes in female meiosis (females being the heterogametic sex). Early in the assembly of the chicken genome, the quality of the build of both the Z and W sex chromosomes was very poor, and limited studies existed on sex determination. Since this, the Z chromosome was painstakingly assembled and sequenced BAC by BAC [Bellott et al., 2010], and is now one of the best-assembled chromosomes in the chicken genome. The same is now expected for the W sex chromosome, which currently is very poorly assembled [Chen et al., 2012]. Zhou Q et al. [2014] conclude that the ancestral sex chromosome organization is closer to that of the Palaeognathae (ostrich and emu) and demonstrated that there is less degradation of the sex chromosomes and a closer synteny with non-avian reptile species.

Copy Number Variation

Redon et al. [2006] first highlighted the impact of CNV in the human genome. This seminal study heralded a new era in cytogenetics and has subsequently been applied to many other species and groups including birds. Skinner et al. [2014] provided a global overview of apparent crossspecies CNVs in birds using cross-species array-CGH. Griffin and Burt [2014] pointed out issues of definition in that 'copy number variation', strictly speaking, refers to polymorphisms within a species. The question arises therefore whether results of cross-species array-CGH represent genuine variation in copies of orthologous genes between species. Skinner et al. [2014] stated that 'difference in gene copy number between species is a question of gene duplication, segmental duplications etc. and may be driven by expansion and contraction of paralogs within different gene families.' Nonetheless, this paper provided a broad appraisal of apparent cross-species CNVs in 16 avian species. Microchromosomes appear to have more apparent CNVs than macrochromosomes. Indeed, in species with microchromosomal fusions such as Falconiformes, the fused 'former microchromosomes' still retained their ancestral features such as a higher degree of cross-species CNVs. Skinner et al. [2014] reported that $\sim 50\%$ of the apparent cross-species CNVs overlap with known chicken-specific CNVs. In terms of gene ontology, there appears to be a general enrichment in immune response and antigen presentation genes as well as 5 CNV regions perfectly correlated with the unique loss of sexual dichromatism. More specifically, there were also suggestions of CNVs involved in diet in turkey (proteolytic digestion/degradation of trypsin inhibitors), and correlation of the unique migratory behaviour of common quail among fowl through the following genes: OBSCN associated with hypertrophy of myofibrils, and MAPK8IP3 implicated in respiratory gaseous exchange [Skinner et al., 2014]. There were also suggestions of an association with muscle activity in falcons through the gain of MYOZ3, preferentially expressed in fast-twitch myofibers and skeletal muscle and an association between immune function in the common quail (Coturnix coturnix) and silver pheasant (Lophura nycthemera) (LEAP2 and ITCH) as well as homeotic genes in common pheasant and California quail (SCML2 and DLX5). Finally, Skinner et al. [2014] identified cross-species CNVs associated with brain development and neuronal function in turkey (e.g. loss of CTXN1), common quail (gain of LRFN5) and duck (e.g. DLGAP2).

Conclusions

The most recent advances in avian cytogenetics have culminated in great promise not only for the study of bird karyotypes, but also for providing insight into the mechanisms of chromosome evolution in general. New avenues for investigation include gene regulation; for instance, it will become necessary to map accurately the physical location of polyadenylation and transcription start sites, important reference points that define promoters and post-transcriptional regulation. It will also become possible to sequence full-length transcripts, to allow accurate identification of alternate splicing events and their controlling elements. The ENCODE (Encyclopedia of DNA Elements) project has helped to define functional elements of the human genome, including those aforementioned as well as other chromatin signals, e.g. active chromatin, enhancers, insulators, methylation domains, etc. An effort of agENCODE is underway to include agriculturally important birds such as chicken, turkey, duck, quail, and perhaps ostrich. The study of cytogenetics will be essential here in helping to define higher-order structures in nuclear organization that show regulatory interactions within and between chromosomes. Finally, reconstruction of evolutionary events allows us to study genome organization and function not only in extant but, by extrapolation, in extinct species also. Reconstruction of avian-reptilian ancestral karyotypes will allow us to define chromosomal rearrangements in long-dead species that have captured the public imagination. Here be dragons!

Hypermethylated Chromosome Regions in Chicken and Other Birds

(Prepared by M. Schmid, C. Steinlein, A.-S. Schneider, I. Nanda, and T. Haaf)

The advent of specific antibodies against the different nucleosides and nucleotides has promoted direct cytogenetic analyses of the various DNA classes along eukaryote chromosomes. These antibodies were first produced by the group of Bernard F. Erlanger some decades ago [Erlanger and Beiser, 1964; Garro et al., 1968; Sawicki et al., 1971; Erlanger et al., 1972]. They were produced by immunizing rabbits to bovine serum albumin (BSA) conjugated to one of the DNA bases. The antibodies are reactive with the BSA conjugate used to induce them and also with single-stranded DNA [Erlanger and Beiser, 1964]. They are highly specific for the base and show little or no cross-reaction with the other bases. Over the years, a se-

specificities for a number of nucleosides, nucleotides and dinucleotides [Dev et al., 1972; Erlanger et al., 1972; Miller, 1973]. In the early 1990s, the first monoclonal antibodies against 5-methylcytosine (5-MeC) and other modified nucleosides were produced [Reynaud et al., 1992] and subsequently used for chromosome staining [Barbin et al., 1994; Miniou et al., 1994; Montpellier et al., 1994; Bernardino et al., 1996].
Of special interest were antisera specific for 5-MeC

ries of such polyclonal antibodies were produced, with

which were initially applied by the group of Orlando J. Miller to the chromosomes of several mammalian species, including human, chimpanzee, gorilla, cattle, mouse, and kangaroo rat [Miller et al., 1974; Schreck et al., 1974, 1977; Schnedl et al., 1975, 1976]. Using an immunofluorescence technique and anti-5-MeC antibodies, they showed that methylated DNA can be detected in fixed metaphase chromosomes after they have been UV-irradiated to generate regions of single-stranded DNA. In these species, the methylated regions corresponded to the locations of repetitive DNA, i.e. to the heterochromatic regions of all or a subset of the chromosomes in the karyotypes. Subsequently, this technique was applied to chromosomes of further mammalian species [Vasilikaki-Baker and Nishioka, 1983; Bernardino et al., 2000] and to human chromosomes [Barbin et al., 1994; Montpellier et al., 1994; Bernardino et al., 1996; Kokalj-Vokac et al., 1998], even including cases of inherited chromosome aberrations [Breg et al., 1974] and leukemia cell lines [Bensaada et al., 1998].

With one exception [Grützner et al., 2001], no immunofluorescence studies on the distribution of hypermethylated regions in bird chromosomes have been published. The present report is a brief summary of the results obtained for avian chromosomes in an ongoing project on the hypermethylation patterns in vertebrate chromosomes [Schmid et al., in preparation].

Mitotic chromosomes of 13 species from 7 orders, belonging to both modern (Neognathae) and primitive (Palaeognathae) birds (table 13), were prepared from embryonic or skin fibroblast cell cultures following standard techniques (colcemid treatment, exposure to hypotonic solution, fixation with methanol:acetic acid). Hypermethylated DNA was detected by indirect immunofluorescence using a monoclonal antibody against 5-MeC. In double-stranded DNA, the methyl groups are hidden in the phosphodiester backbone of the double helix and not accessible to the antibody. The anti-5-MeC antibody recognizes and binds to its target only if the DNA is in the single-stranded configuration. Therefore, the slides with

Order, family	Species	Common name	2n ^a	Hypermethylated heterochromatic regions present in				
				macroautosomes ^{b, c}	microautosomes ^{b, c}	Z chromosome	W chromosome	
Anseriformes Anatidae	Anas platyrhynchos	mallard duck	80	+ (15)	+ (20)	+	+	
Columbiformes Columbidae	Streptopelia risoria	ringneck dove	74	+ (15)	+ (19)	_	+	
Galliformes								
Phasianidae	Gallus gallus Coturnix coturnix Pavo cristatus	chicken Japanese quail Indian peafowl	78 78 78	+ (12) + (1) + (1)	+ (23) + (15)	- - +	+ + d	
	Phasianus colchicus	ring-necked pheasant	82	+(1) + (12)	+ (23)	_	-	
Gruiformes								
Rallidae	Gallinula chloropus	common moorhen	78	+ (3)	+ (11)	-	+	
Passeriformes								
Corvidae Paridae Psittacidae	Garrulus glandarius Parus ater Melopsittacus undulatus	Eurasian jay coal tit budgerigar	78 80 62	+ (17) + (5) + (5)	+ (21) + (6) + (8)	+ +	+ d	
Sittidae	Sitta europaea	Eurasian nuthatch	80	+(17)	+ (22)	_	d	
Rheiformes Rheidae	Rhea americana	nandu	80	+ (1)	+ (20)	_	_	
Struthioniformes Struthionidae	Struthio camelus	ostrich	80	+ (7)	+ (17)	_	-	

Table 13. Hypermethylated heterochromatic chromosome regions detected by 5-MeC antibody in 13 bird species

^a Because of the tiny size of some microautosomes, the exact diploid chromosome number is uncertain in some species.

^b In all species analyzed, the chromosomes continuously decrease in length, and there is no clear-cut size difference between macro- and microautosomes. Therefore, the first 13 autosome pairs were arbitrarily classified as macroautosomes and the remaining pairs as microautosomes.

^c Numbers in parentheses indicate the number of macro- and microautosomes containing heterochromatic regions with distinct immunofluorescence labeling.

^d Because only male (ZZ) specimens were examined, data on W chromosomes are not available.

the chromosome preparations were immersed 1 cm below the level of a buffer solution (PBS) and denatured by UVlight irradiation for 2.5–3 h at a distance of 10 cm from a UV lamp (254 nm). For indirect immunofluorescence, the slides were first incubated in a coplin jar for 1 h in blocking solution (PBS, with 0.3% BSA, 0.1% Tween) and then with 50 µl of a monoclonal mouse anti-5-MeC (primary) antibody (Imprint[®]monoclonal anti-5-methyl-cytosine antibody 33D3, Sigma-Aldrich) diluted 1:1,000 with the blocking solution in a humified incubator at 37°C for 1 h. A non-siliconized coverslip $(22 \times 60 \text{ mm})$ was placed on the 50-µl drop to spread the anti-5-MeC antibody over the complete slide surface. Subsequently, the slides were washed twice in PBS (with 0.3% BSA) for 3 min each and then incubated with 70 µl of the secondary antibody (TRITC-conjugated rabbit-anti-mouse IgG, Sigma-Aldrich) diluted 1:200 with PBS. The incubation conditions were as for the primary antibody. After 2 further washes

with PBS for 3 min each, the chromosome preparations were mounted in Vectashield[®] mounting medium (Vectashield) with DAPI. Image analysis was performed with Zeiss epifluorescence microscopes equipped with thermoelectronically cooled charge-coupled device cameras (Applied Spectral Imaging) using FISHView 6.0 software. The demonstration of constitutive heterochromatin (Cbands) in the chromosomes followed the technique of Sumner [1972]. C-banded metaphases were photographed under a Zeiss Axio ImagerA1 epifluorescence microscope employing the digital imaging system software BandView 6.0 (Applied Spectral Imaging).

The present study shows that, like in mammals, the hypermethylated regions in the karyotypes of birds are largely confined to constitutive heterochromatin (figs. 5, 6; table 13). In some preparations, faint fluorescence signals are also located outside the constitutive heterochromatin, but these are not consistent if many metaphases



Fig. 5. Karyotypes of female chicken (*Gallus gallus*) (**a**–**c**) and male Indian peafowl (*Pavo cristatus*) (**d**–**f**) after indirect immunofluorescence using a monoclonal antibody against 5-MeC (**a**, **b**, **d**, **e**) and C-banding (**c**, **f**). The 5-MeC-rich heterochromatic regions show red fluorescence signals, the chromosomes are stained blue with DAPI.

are analyzed (e.g. macroautosomes 1 in fig. 5a, b). Again, similar to mammals, there seems to be no strict rule on the distribution of hypermethylated chromosome regions in birds. In most instances, hypermethylated heterochromatic regions are located in the centromeric regions of chromosomes, but in sex chromosomes they can also be found in telomeric postitions (fig. 5d, e). In most of the species examined, the centromeric heterochromatin in many, if not all, of the microchromosomes is hypermethylated (figs. 5a, b, 6a, b; table 13), but there are no-



Fig. 6. Karyotypes of female mallard duck (*Anas platyrhynchos*) after indirect immunofluorescence using a monoclonal antibody against 5-MeC (**a**, **b**) and C-banding (**c**). The 5-MeC-rich heterochromatic regions show red fluorescence signals, the chromosomes are stained blue with DAPI.

table exceptions. Thus, in the karyotype of *Pavo cristatus*, the only detectable hypermethylated heterochromatic regions are located in the macroautosome pair 1 and in the Z chromosome. None of the microchromosomes is labeled (fig. 5d, e).

The nearly completely heterochromatic W sex chromosomes of neognathous birds, which are distinctly smaller than their corresponding Z chromosomes, show distinct hypermethylation, although in some species the W chromosomes exhibit only partial immunofluorescent labeling (fig. 5a, b). In contrast, in the extant palaeognathous birds (ostrich, emu, nandu), the primitive ZW sex chromosomes are still largely homomorphic, mostly euchromatic and contain similar genes [Fridolfsson et al., 1998; Ogawa et al., 1998; Shetty et al., 1999; Nishida-Umehara et al., 2007]. No hypermethylated heterochromatic regions are present in the W chromosomes of these species (table 13).

In the female specimens examined, visible differences in the hypermethylation patterns between the 2 Z sex chromosomes are not present. Depending on the species, both Z chromosomes contain either no hypermethylated heterochromatic regions at all, or both have clear fluorescing signals which were present in all metaphases analyzed (fig. 5d, e).

A close analysis of high-quality karyotypes of those species possessing many hypermethylated microchromosomes reveals that, even in the smallest microchromosomes, the immunofluorescence labeling is restricted to the heterochromatic centromere regions and does not extend over the whole microchromosomes (fig. 6a, b). In a previous study on neognathous birds (chicken, quail, pheasant) and palaeognathous birds (nandu, emu), Grützner et al. [2001] visualized the hypermethylated chromosome regions by the technique of indirect immunofluorescence, but used a different, well-characterized antibody against 5-MeC [Miniou et al., 1994; Mayer et al., 2000], and denatured the chromosome preparations in 70% formamide, 2× SSC for 1 min at 90°C. Contrary to the present study, they observed an apparently complete immunofluorescence labeling of the microchromosomes. They correlated this high density of methylated cytosines with the high gene density on the avian microchromosomes. Indeed, there is considerable evidence for an increased density of genes in microchromosomes [Smith J et al., 2000]. They have a

high GC base-pair content, enrichment with CpG islands, histone H4 hyperacetylation, and early replication in the S-phase of the cell cycle [Schmid et al., 1989; McQueen et al., 1996, 1998]. The figures published by Grützner et al. [2001] show dense clusters of microchromosomes, so it is not clearly evident if the immunofluorescence actually extends over the entire microchromosomes. In this context, it is important to emphasize that immunofluorescence patterns do not only reflect inherent chromosomal properties (i.e. distribution and enrichment of 5-MeC) but are also dependent on the different denaturation conditions and antibody specificities used. Moreover, the methylation status does also depend on the cell type and culture conditions. Admittedly, some sites of DNA methylation may not be recognized by the technique of UV-irradiation used here because the anti-5-MeC antibody can attach only to regions of single-stranded DNA. UV-irradiation is particularly effective in generating single-stranded regions in the DNA which is rich in AT base pairs with adjacent pyrimidines so that thymine dimer formation can occur [Schreck et al., 1974; Schnedl et al., 1975]. In DNA regions which are not denatured by the UV-irradiation technique, 5-MeC would not be recognized. However, partially denatured DNA strands in euchromatic chromosome regions do not suffice to explain the differences obtained in the present study and those published by Grützner et al. [2001].

Although karyotypes of a very limited number of avian species were examined with the present technique, it already becomes obvious that more closely related species within the same family do not share similar patterns of hypermethylated heterochromatic regions (table 13). Apparently, these patterns do not represent stable properties reflecting a common phylogenetic ancestry of different taxa. This is to be expected, because the major components of constitutive heterochromatin are repetitive DNA sequences which are characterized by repetition of relatively long monomers (of a few hundred base pairs) over many megabases of DNA [for reviews, see Brutlag, 1980; Long and Dawid, 1980; Singer, 1982; Southern, 1984; Beridze, 1986]. It is not uncommon to find up to 25% of a genome made up of different repetitive DNA families [for review, see Lohe and Roberts, 1988]. Even among closely related species, reiterated DNAs usually differ in quantity, sequence and chromosomal location [Miklos, 1985; Charlesworth et al., 1994]. Since they are not subjected to evolutionary selection pressure, sequence changes in repetitive DNAs can be accumulated and fixed in genomes very much faster than can the changes in functional (transcribed) DNA. As a consequence, constitutive heterochromatin is heterogeneous within and between species [for review, see Verma, 1988]. This heterogeneity of heterochromatin is paralleled by extremely rapid changes of its hypermethylation patterns.

The data obtained indicate that, like in mammals, the hypermethylation patterns of constitutive heterochromatin in birds are species-specific with respect to size, location and staining intensity. Because of this species-specificity, they are extremely useful as cytogenetic parameters for differentiating between closely related species that possess the same diploid number and chromosome morphology. It should, however, be emphasized that the results of this preliminary study were obtained exclusively on mitotic chromosomes prepared from fibroblast cultures. It must be confirmed whether the hypermethylation patterns of heterochromatin actually constitute a stable intraspecific feature, or if there exist tissue-specific patterns. In this context, the examination of meiotic chromosomes, especially those in the lampbrush stage of female bird meiosis is important.

An Overview of Avian Evolution

(Prepared by S.B. Hedges)

Present and Past Diversity

The living birds form a large and diverse group of vertebrates, with nearly 10,000 species in ~2,000 genera, 200 families and 29 orders [Sibley and Monroe, 1990; IUCN, 2014]. New species are discovered each year, but at a low rate indicating that our knowledge of current avian diversity is quite good. The major uncertainty in the total number of species concerns disagreement over the definition of particular species, such as the recognition of subspecies as full species, rather than discovery of new species. In contrast, other groups of tetrapods (e.g. mammals, lizards, snakes, amphibians) have a higher rate of species discovery [Uetz and Hošek, 2014].

The fossil record of birds is relatively poor, probably because the avian skeleton is fragile (hollow bones), and most species are small and occur in environments (humid forests) where decomposition is rapid. However, the early history of birds has become much better known in recent decades with the discovery of exceptionally preserved fossils of birds and dinosaurs from the Cretaceous (142–65 million years ago, Mya), especially in China [Chiappe and Dyke, 2002]. The earliest bird is *Archaeopteryx* from the late Jurassic (~150 Mya) of Ger-

many, and the consensus view is that birds evolved from carnivorous (theropod) dinosaurs called coelurosaurs (e.g. Tyrannosaurus rex), and specifically from a group of relatively small and agile species called dromaeosaurids (e.g. Velociraptor). This was suggested in the 19th century when Archaeopteryx was first classified as a small dinosaur, before the impressions of feathers were noted, and has since been supported by detailed comparisons of anatomy and (recently) egg morphology [Varricchio et al., 1997]. To recognize this derivation from dinosaurs, paleontologists sometimes refer to birds as 'living dinosaurs' and refer to classical dinosaurs (e.g. Stegosaurus) as 'non-avian dinosaurs'. The carnivorous (theropod) ancestry of birds is also illustrated by the fact that Archaeopteryx and many of the other Mesozoic (251-65 Mya) species possessed sharp teeth and raptorial claws.

Discoveries of feathered dinosaurs in the last 2 decades have also bolstered the link between birds and dinosaurs. In the earliest-branching species of coelurosaurs, the feathers were filamentous and probably functioned more like the down of some living birds, providing insulation [Chiappe and Dyke, 2002]. However, the maniraptoran coelurosaurs (including dromaeosaurids), which were the closest relatives of birds, possessed vaned feathers that more closely resembled avian flight feathers. There is no evidence that these dinosaurs could undergo the sustained flapping flight of modern birds, but extensive development of feathers and their configuration indicate that at least some of these dinosaurs were gliders [Xu et al., 2003]. These fossils have helped to blur the distinction between dinosaurs and birds. Nonetheless, the full suite of adaptations for sustained, flapping flight, such as asymmetric feathers, alulas (wing structures for improving the airfoil function), and other aerodynamic structures, are found only in birds. Even Archaeopteryx probably was able to initiate flight directly from the ground [Chiappe and Dyke, 2002].

Debate continues as to how the Cretaceous birds are classified. One arrangement divides them into 2 large groups (Sauriurae and Ornithurae), whereas another places them in a ladder-like tree of lineages leading to modern birds [Chiappe and Dyke, 2002]. In either case, the closest relative of modern birds is believed to be *Ichthyornis*, a small, toothed, tern-like marine bird of the late Cretaceous. Some Cretaceous fossils have been postulated to be representatives of modern orders such as Galliformes (e.g. fowl), Anseriformes (e.g. ducks), Psittaciformes (e.g. parrots), Charadriiformes (e.g. plovers), Procellariformes (e.g. petrels, albatrosses), Gaviiformes (e.g. loons), Gruiformes (e.g. cranes), and Pelecaniformes (e.g. pelicans). However, all of these fossils are considered to be problematic in some way [Chiappe and Dyke, 2002]. With those aside, the fossil record of birds shows a major dichotomy at the Cretaceous-Tertiary boundary (65 Mya), when a large asteroid collided with Earth causing the extinction of the dinosaurs and other life. Almost no unambiguous fossils of modern (neornithine) birds are known before that event and no fossils of non-neornithine birds are known after that event. One important exception is *Presbyornis*, an aquatic neornithine bird apparently related to ducks (Anseriformes) which is known from the late Cretaceous and the early Tertiary [Kurochkin et al., 2002]. Fossils of most orders of modern birds appear in the early part of the Cenozoic (65–0 Mya).

Phylogenetic Relationships

Until recently, the phylogeny of modern birds has been poorly known despite decades of attention from morphologists and molecular phylogeneticists. Although early molecular studies [Sibley and Ahlquist, 1990] identified some obvious cases of convergence in previous morphological classifications and have helped to better organize the higher-level classification of birds, it has, in general, been a slow progression of knowledge. Probably the best explanation of the slowness is data limitation (need for many genes) and closeness in time of divergence among the neoavian orders [Cracraft et al., 2004]. Recent genome analyses have started to close this knowledge gap and have finally begun to resolve the tree of bird orders [Jarvis et al., 2014].

Some of the higher-level clades have been firmly established (fig. 7). For example, there is wide agreement that modern birds (Neornithes) form 3 major clades: Paleognathae (tinamous and ratites), Galloanserae (e.g. ducks, fowl), and Neoaves (all other birds). In addition, DNA sequence analyses, primarily of nuclear genes, indicate that Galloanserae and Neoaves are closest relatives [Groth and Barrowclough, 1999; van Tuinen et al., 2000; Hackett et al., 2008; van Tuinen, 2009; Jarvis et al., 2014]. Although earlier DNA hybridization studies had correctly identified the 3 groups [Sibley and Ahlquist, 1990], those data were ambiguous regarding their relationships. Initially, even data from complete mitochondrial genomes (~16 kb) provided a conflicting signal, indicating that Passeriformes was the most basal clade of modern birds and that the paleognaths and galloanserines were close relatives [Härlid and Arnason, 1999; Mindell et al., 1999]. Later mitochondrial analyses with additional taxa found support for the paleognath-basal tree [Paton et al., 2002; Garcia-Moreno et al., 2003].

Schmid et al.



Fig. 7. Relationships and divergence times of birds and other amniote vertebrates (see text). The closest living relatives of birds are the crocodilians [Hedges, 2012], but the split with turtles is similar in time. The shaded area in Neoaves represents a time when most ordinal and superordinal lineages diverged based on molecular clocks. The lineages of Mesozoic birds and *Archaeopteryx* are shown ending arbitrarily at the Cretaceous/Tertiary boundary, although fossil data are sparse and some lineages may have disappeared earlier.

The living ratites (flightless paleognaths) include the ostrich of Africa, rheas of South America, the emu of Australia, the cassowaries of Australia and New Guinea, and the kiwis of New Zealand. Extinct species occurred on Madagascar (elephant bird), New Zealand (moas), and elsewhere. The phylogeny of these birds has garnered considerable interest, in part because of their primarily southern distribution and the expectation that the breakup of Gondwana influenced their biogeographic history. Although the moas are now extinct, complete mitochondrial genome sequences have been obtained from fossil remains, and phylogenetic trees have been constructed with large data sets of DNA sequences. Nonetheless, the relationships have proven to be difficult to resolve. Three of the Australasian species (kiwi, emu and cassowary) form a group in most analyses, but the moa usually branches basally when present in the tree [Lee et al., 1997; van Tuinen et al., 1998; Cooper et al., 2001; Haddrath and Baker, 2001; Paton et al., 2002]. In contrast, morphological trees place the kiwi as a basal lineage, separate from the other Australasian species [Lee et al., 1997]. More recently, a nuclear gene study found tinamous to be nested

among ratites rather than as closest relatives [Hackett et al., 2008]. For these reasons, firm biogeographic conclusions must await additional data and analyses.

The Galloanserae includes 2 orders of primarily robust-bodied birds. The Galliformes consists of mostly non-aquatic, ground-dwelling birds such as game fowl, megapodes, guans, and chachalacas. The chicken (Gallus gallus; common name, domestic fowl) belongs to this order. Its wild counterpart (also G. gallus) goes by the common name red junglefowl and is native to the lowlands (<2,000 m) of southern Asia and the Malay archipelago [Sibley and Monroe, 1990]. It is placed in the family Phasianidae along with other widely domesticated species such as the common quail (Coturnix coturnix), turkey (Meleagris gallopavo) and pheasant (Phasianus colchicus). Other members of the order Galliformes include the New World quails (Odontophoridae) and guinea fowl (Numididae). The other order within Galloanserae, the Anseriformes (ducks, geese, swans, screamers), is primarily aquatic and comprises about 450 species in ~125 genera [Sibley and Monroe, 1990].

Neoaves is the third major clade of living birds and accounts for 95% of the species. Although some of the 20 or so neoavian orders are well-defined, others are not [Cracraft et al., 2004]. For example, the diving birds, wading birds and marine birds are placed by most morphologists into about 8-9 different orders, including Charadriiformes (e.g. gulls, plovers), Ciconiiformes (e.g. storks), Gaviiformes (loons), Gruiformes (e.g. cranes), Pelecaniformes (e.g. pelicans, boobies, tropicbirds), Phoenicopteriformes (flamingos), Podicipediformes (e.g. grebes), Procellariiformes (albatrosses, shearwaters), and Sphenisciformes (penguins). Most of these orders are recognized today in avian field guides and other reference sources. However, DNA hybridization data [Sibley and Ahlquist, 1990] and DNA sequences [Hedges and Sibley, 1994; van Tuinen et al., 2001; Cracraft et al., 2004; Jarvis et al., 2014] have revealed relationships that place into question the recognition of those orders as classically defined. For example, the pelicans are most closely related to the shoebill stork and hammerkop and not to the boobies or tropicbird, and the grebes are most closely related to the flamingos, not to the loons [Hedges and Sibley, 1994; van Tuinen et al., 2001; Cracraft et al., 2004]. Those surprising findings have been obtained with multiple genes in different laboratories, and they have strong statistical support. Other controversial findings among the waterbirds, such as a clustering of storks with New World vultures [Sibley and Ahlquist, 1990], have not been corroborated with genomic data [Jarvis et al., 2014].

Besides the water birds, progress has been made with DNA sequence analyses in understanding relationships within other neoavian groups, including the Passeriformes [Johansson et al., 2001, 2002; Barker et al., 2002; Yuri and Mindell, 2002; Cracraft et al., 2004]. One particularly problematic bird has been the hoatzin (Opisthocomus hoazin) of South America, sometimes placed in its own order (Opisthocomiformes). It is the only bird that uses microbial foregut fermentation to convert cellulose into simple sugars, as in some mammals (e.g. ruminants). It feeds on young leaves and twigs of marsh plants and has a large muscular crop for fermentation. Most earlier phylogenetic analyses have placed the hoatzin in the Galloanserae, but DNA hybridization studies and DNA sequence studies have agreed that it is a neoavian species [Sibley and Ahlquist, 1990; Hedges et al., 1995; Hughes and Baker, 1999; Sorenson et al., 2003]. However, its particular relationship within Neoaves has been hard to resolve, even in the recent complete genome analysis [Jarvis et al., 2014].

Similarly, some other nodes in the phylogeny of neoavian bird orders have been difficult to resolve. However, large numbers of nuclear genes and complete genomes [Hackett et al., 2008; Jarvis et al., 2014] have succeeded in resolving most nodes. The results show an unexpected clade of birds (Columbea), comprising flamingos, grebes, and pigeon relatives, as the closest relative of all other neoavian orders, and – among the remaining orders (Passerea) – 2 large clades comprising birds of similar ecological zone: a land clade and a water clade [Jarvis et al., 2014]. Other orders of Passerea do not fit neatly into the land and water clades, and more work will be needed to better resolve the branching pattern of the orders and understand the important traits involved in their evolutionary history.

Molecular Clocks and Biogeography

A literal reading of the fossil record of birds is that the modern orders evolved and radiated in the early Cenozoic, after the Cretaceous-Tertiary extinctions [Feduccia, 1995, 2003]. Even the problematic Cretaceous fossils of Neornithine birds noted above are mostly from the final stage of the Cretaceous (Maastrichian; 71–65 Mya). However, molecular clocks have instead indicated that the orders of modern birds branched more deeply in the Cretaceous [Hedges et al., 1996; Cooper and Penny, 1997; Kumar and Hedges, 1998; Haddrath and Baker, 2001; van Tuinen and Hedges, 2001; Paton et al., 2002]. If these molecular clocks are correct, why is there such a large gap in the avian fossil record? Local clock methods (e.g. Bayes-

ian) have been used in addition to global clocks, and some studies [Hedges et al., 1996; Kumar and Hedges, 1998; van Tuinen and Hedges, 2001] have used only rate-constant genes calibrated from outside of the avian fossil record to avoid potential calibration biases. Despite the use of these different methods, and carefully controlling for rate, the resulting time estimates have been fairly consistent in supporting relatively deep divergences among modern birds (fig. 7).

The divergence of the chicken (Galliformes) and duck (Anseriformes) has proven to be an important anchor point for avian molecular clocks. Anseriformes is about the only order of modern birds that can be confidently placed in the Cretaceous, with the earliest fossils from the Campanian (84-71 Mya) [Chiappe and Dyke, 2002]. Calibrating outside of birds and using 12 rate-constant nuclear genes, this divergence was estimated to be 90 \pm 7 Mya [van Tuinen and Hedges, 2001]. A similar date of 85 ± 17 Mya was obtained with different data (mitochondrial DNA), calibrations (internal avian calibrations), and clock methods ('rate smoothing' local clock) [Haddrath and Baker, 2001]. The advantage of establishing an anchor point within birds is to provide a robust calibration for obtaining other time estimates among birds [van Tuinen and Hedges, 2004]. Using that chicken-duck calibration point, several data sets were analyzed (mitochondrial DNA, DNA-DNA hybridization, and transferrin immunological data), resulting in the following mean divergence times for major nodes: 119 ± 5 Mya (paleognath-neognath), 104 ± 3 (Galloanserae-Neoaves), and 89 ± 9.6 Mya (base of neoavian radiation). Similar dates for the earliest nodes were obtained in a more recent review [van Tuinen, 2009]. Dates among paleognaths ranged from 50 (kiwi-emu) to 83 Mya (tinamous-ratites), between orders from 76 to 80 Mya, and within orders from 39 (chicken-quail) to 80 Mya (kingfisher-hornbill). However, sequence data were limited and most neoavian orders were not represented. Recent reviews of molecular time estimates found Cretaceous dates for all nodes among ratites and tinamous [Baker and Pereira, 2009] and most nodes among Galloanserae, with a mean split time of 107 Mya between ducks and fowl [Pereira and Baker, 2009].

In another study [Paton et al., 2002], divergence time estimates were obtained using mitochondrial sequence data, local clock methods, and calibrations of 85 Mya (chicken-duck) and 35 Mya (emu-cassowary). The resulting times were similar to an earlier study in finding deep splits (Cretaceous) among neoavian orders and a deep divergence between paleognaths and neognaths (123 Mya; 156–108 Mya). However, their dates for divergences among paleognaths were slightly older: 105 Mya (tinamous vs. ratites), 89 Mya (rhea vs. ostrich), and 81 Mya (kiwi vs. emu). More recently, Pacheco et al. [2011] constructed a timetree of neoavian orders using mitochondrial genomes and found that most divergences occurred in the Cretaceous, corroborating earlier work. As more sequences are obtained, especially from nuclear genes, time estimates from molecular clocks should become more stable until the point where most or all error is from the fossil record and calibration points [Hedges and Kumar, 2003, 2004].

A timetree was presented with the recent multi-genome analysis of bird evolution [Jarvis et al., 2014]. It showed a split between Palaeognathae and Neognathae of about 100 Mya, the split of Neoaves and Galloanserae of 88 Mya, and the split of most neoavian orders about 65 Mya. These times are considerably (16%) younger than found in earlier studies, reviewed above [e.g. van Tuinen, 2009]. It should not be concluded that the more recent study [Jarvis et al., 2014], based on more data (complete genomes), is correct and the earlier studies are incorrect. The difference, instead, is likely from an error (calibration bias) in the methodology used in the recent genome study [Jarvis et al., 2014]. Those authors used a suite of fossil minimum calibrations and one maximum calibration. That maximum calibration, the split of Palaeognathae and Neognathae, was set to 99.6 Mya for an arbitrary reason: it is the boundary between the Early and Late Cretaceous, with the authors qualifying that it 'far exceeds the age of paleontological evidence for the existence of Neornithes.' However, because the resulting analysis returned a date for that maximum calibration node equal to the calibration for that node, it suggests that the true time is earlier and that the resulting times throughout the tree are likely underestimates. Otherwise, it would be a remarkable coincidence that the maximum calibration used, thought to be much older than the true date for the node, would exactly match the resulting time. This data set should be reanalyzed using a better calibration methodology.

With this current timescale and phylogeny of avian evolution (fig. 7), and recognizing its limitations, it is possible to draw some general inferences concerning the niche and morphology (habitus) of birds during their early history. For example, the heavy-bodied habitus and primarily ground-dwelling, non-marine habits of the Galloanserae and Palaeognathae, and their dinosaurian ancestors, suggest that these features were common among the stem neornithine birds of the Cretaceous [van
Tuinen et al., 2000]. Also, considering that shallow, marine environments are favorable for fossilization, this (the predominance of non-marine species) may in part explain the sparseness of the early fossil record of modern birds. Nonetheless, fossils indicate that Cretaceous birds also occupied other niches (e.g. perching, diving, etc.).

The early (Cretaceous) branching of ordinal lineages of modern birds found with molecular clocks 'is compatible with the extensive speciation that occurred within orders following the sudden availability of niches in the early Tertiary period' [Hedges et al., 1996]. Those niches were vacated by dinosaurs, pterosaurs, and other vertebrate groups that became extinct following the Cretaceous-Tertiary asteroid impact. The near simultaneous appearance of so many major lineages of neoavian birds, adapted to different environments, in the early Tertiary supports the niche-filling hypothesis.

One explanation for the early branching of ordinal and superordinal lineages of birds is that they reflect the splitting of landmasses in the Cretaceous, the Continental Breakup hypothesis [Hedges et al., 1996]. The same hypothesis was proposed for the orders of placental mammals [Hedges et al., 1996], based on molecular clock evidence, and subsequent phylogenetic analyses have provided support [Springer et al., 1997; Stanhope et al., 1998]. In birds, it has proven more difficult to associate particular orders with geographic regions (e.g. continents) because of their wider distributions and ease of dispersal. As noted above, ratites have been singled out as having an evolutionary history associated with continental breakup [Cracraft, 1973, 2001; van Tuinen et al., 1998], but biogeographic details will remain sketchy until their relationships are better resolved. A good case has been made that modern (neornithine) birds in general are Gondwana in origin, based on a review of fossils, phylogeny, and molecular divergence times [Cracraft, 2001]. However, the extent that plate tectonics and the divergence of continents have had on avian evolution remains unanswered.

Completion of the chicken genome, and recently other bird genomes, will be a major benefit for evolutionary studies. For example, orthology determination, which is necessary for assembling phylogenetic data sets of nuclear genes, will be greatly facilitated by having avian genomes for comparison. Also, genomic comparisons will now be able to include many representative birds and therefore increase precision in analyses and questions being addressed, providing a better evolutionary context across amniote vertebrates [Hedges, 2012]. Finally, detailed studies within birds, involving molecular clocks and phylogenies, will have the benefit of avian genomes for primer design, clock calibration, and better genetic comparisons in general.

I thank Jennifer Hines for drawings of animals.

An Update on Chicken Sex Determination and Gonadal Sex Differentiation

(Prepared by C.A. Smith)

Since the publication of the Second Report on Chicken Genes and Chromosomes 2005 [Schmid et al., 2005], there have been some significant advances in our understanding of avian sex chromosomes and sex determination [Smith CA et al., 2000; Zhao et al., 2010; Chue and Smith, 2011; Ayers et al., 2013c]. Sex in chicken and other birds is governed by the inheritance of sex chromosomes, with male being ZZ and female ZW. It is still unclear whether sex is determined by the presence/absence of the W sex chromosome or whether it depends upon Z dosage or the Z to autosome ratio (2 Z's in male; 1 Z in female). This question could be solved once and for all by determining the sexual phenotype of birds with sex chromosome aneuploidy (e.g. 2A:Z0 or ZZW), but such birds have not been definitively identified, and indeed these genotypes could be embryo lethal in avians due to potential gene dosage effects [Graves, 2003]. Although some recent studies on sex determination look beyond the gonads (discussed below), most researchers in the field still equate 'sex determination' with embryonic gonadal sex differentiation. While sex is set at fertilization with the differential inheritance of the sex chromosomes, one or more genes on these chromosomes direct differentiation of the gonads into testes or (unilateral) ovary during embryonic life. Such genes - or different sex-linked genes may also direct sexual differentiation outside the gonads in a cell-autonomous fashion (discussed below). At present, most evidence favours Z dosage underlying chicken sexual differentiation, at least in the gonads [Chue and Smith, 2011; Ayers et al., 2013c].

Figure 8 shows the genes currently known or postulated to play a role in gonadal sex differentiation in the chicken embryo. The gonads are of mesoderm origin, forming on the surface of the mesonephric kidneys. Sexual differentiation begins at the morphological level on embryonic day 6.0–6.5 [Hamburger and Hamilton (HH) stage 29–30], when seminiferous cords of the inner medulla condense in males (ZZ), and the outer epithelial layer thickens in females (ZW) [Hamburger and Hamilton,



Fig. 8. Gonadal sex differentiation in the chicken embryo. The undifferentiated gonad comprises a medulla (blue) and outer epithelial layer (pink). Germ cells are shown as black circles. The medulla proliferates in males during testis formation, while the epithelial layer proliferates in females to generate the cortex of the ovary. Male expressed genes are shown in blue, female expressed genes in pink. The size of the circles represents relative expression levels compared to the opposite sex. *DMRT1* is expressed in both sexes, but more highly in males. *GATA4* is also expressed in both sexes and may activate DMRT1. In males (ZZ), DMRT1 can acti-

vate *HEMOGEN* (*HEMGN*), *AMH* and *SOX9* expression, either directly or indirectly. *HEMOGEN* can activate *SOX9* and also *DMRT1*. *AMH* mRNA is expressed prior to *SOX9*, and may be activated by SF1 together with DMRT1. Prostaglandin D₂ (PGD₂) can stimulate *SOX9*. DMRT1 can repress *FOXL2* and aromatase (*CYP19A1*). In females (ZW), *MHM* may contribute to lower *DMRT1* expression. FOXL2 likely activates aromatase (*CYP19A1*). SF1 may also contribute to *CYP19A1* expression. R-SPO1 and WNT4 stabilize β -catenin. Both aromatase (estrogen) and β catenin regulate ovarian development.

1992; Guioli et al., 2014]. The gonadal asymmetry characteristic of female gonads (only the left becomes a functional ovary) has been shown to be driven by the asymmetric expression of the *PITX2* gene and, downstream, differential activity of retinoic acid signaling and cell cycle regulators [Guioli and Lovell-Badge, 2007; Ishimaru et al., 2008; Rodríguez-León et al., 2008]. There are some interesting differences between gonadal development in mouse and chicken. The key preSertoli cell lineage, for example, derives from the outer epithelial layer of the gonad in the mouse embryo, but this layer appears to generate interstitial cells in the chicken, indicating the preSertoli cell population likely comes from the mesonephric kidneys (hence a mesenchymal origin) [Sekido and Lovell-Badge, 2007].

The best candidate master regulator of chicken gonadal sex differentiation is the Z-linked gene, *DMRT1* (doublesex and mab-3 related transcription factor, #1) [Chue and Smith, 2011; Matson and Zarkower, 2012]. This gene encodes a zinc-finger-like transcription factor, and it has a conserved role in testis development after birth in mammals [Matson and Zarkower, 2012]. This role includes maintenance of Sertoli and germ cells [Minkina et al., 2014]. It the mouse, DMRT1 prevents the ovarian developmental pathway from being activated, even in adults [Matson et al., 2011]. In the chicken, DMRT1 is expressed specifically in the embryonic urogenital system, where it is more highly expressed in male compared to female gonads [Raymond et al., 1999; Smith et al., 1999; Shan et al., 2000]. Knockdown of DMRT1 expression using RNA interference results in feminization of the gonads, characterized by loss of seminiferous cord structure and male markers, and activation of the female marker, aromatase [Smith et al., 2009]. Most recently, using in ovo electroporation of DNA constructs into developing gonads, we over-expressed DMRT1 in the embryonic chicken urogenital system and found that it can induce male pathway genes (SOX9 and AMH) in genetically female (ZW) go-

nads [Lambeth et al., 2014]. This further strengthens the proposal that Z-linked DMRT1 initiates testis development in the chicken (fig. 8). Downstream of DMRT1, the conserved AMH and SOX9 genes are male-upregulated and likely play a role in testis development, as in mammals (fig. 8). In mammals, SOX9 activates AMH during testicular development, but in the chicken, AMH mRNA expression precedes that of SOX9, indicating that the AMH gene is activated by other factors in birds. However, at the protein level, we detect SOX9 before or at the same stage as AMH protein, suggesting delayed translation of AMH mRNA until the onset of SOX9 expression [Lambeth et al., submitted]. In mammals, prostaglandin D_2 is synthesized male-specifically, and it augments SOX9 expression in preSertoli cells, and this process has been shown to be conserved in the chicken embryo [Moniot et al., 2008].

Recently, another Z-linked gene, HEMOGEN, has been implicated in chicken testicular development [Nakata et al., 2013]. As for DMRT1, the gene encodes a transcription factor. It is involved in haematopoiesis in mammals, but is not expressed in the embryonic mouse gonad. In contrast, it is expressed in male but not female embryonic chicken gonads. Its overexpression induces both DMRT1 and SOX9 expression in female (ZW) embryos, and the gonads appear masculinized (sex cords, maletype arrangement of germ cells). Given its location on the Z sex chromosome, HEMOGEN could lie upstream or act codominantly with DMRT1. If it lies upstream, then a Wlinked factor would be implicated in its suppression in females (directly or indirectly). However, there is still no evidence for a W-linked ovary determinant in chicken, despite our recent transcript profiling analysis of embryonic chicken gonads using RNA-seq [Ayers et al., 2013a]. More likely is the possibility that Z-linked DMRT1 activates the HEMOGEN gene during testis development. This is supported by the fact that DMRT1 is expressed before HEMOGEN in developing male (ZZ) gonads, and overexpression of DMRT1 in female gonads can induce HEMOGEN expression [Lambeth et al., 2014]. As shown in figure 8, DMRT1 may activate HEMOGEN, which in turn activates SOX9 during testicular development. Knockdown of HEMOGEN expression would clarify this point, but this experiment has not yet been reported.

Unilateral ovary development occurs in female chicken embryos (ZW). It has been known for over 30 years that estrogen synthesis is essential for ovarian development in birds [Smith et al., 1997; Vaillant et al., 2001; Lambeth et al., 2013]. The terminal enzyme responsible for estrogen synthesis, aromatase (encoded by the CYP19A1 gene), is expressed female-specifically from the earliest stage of ovarian differentiation (HH29), while inhibition of aromatase causes female-to-male sex reversal (fig. 8). The regulation of gonadal CYP19A1 is therefore a pivotal event in chicken gonadal sex differentiation. The conserved FOXL2 transcription factor colocalizes with aromatase, and evidence from chicken and other vertebrates indicates that it activates aromatase during ovarian differentiation [Govoroun et al., 2004; Pannetier et al., 2006; Wang DS et al., 2007; Chue and Smith, 2011]. Other genes may also participate in the regulation of CYP19A1 in chicken, such as SF1, GATA4 and DAX1 (fig. 8). Recently, epigenetic modifications of key gonad genes such as CYP19A1 have come under scrutiny, in chicken and in other species. Chicken CYP19A1 shows sexually dimorphic methylation in a region of its gonadal promoter (45% methylated in ZZ males vs. 20% in ZW females), a pattern that is partially reversed when gonads are feminized with exogenous estrogens [Ellis et al., 2012]. This suggests that part of the mechanism of chicken gonadal development may involve differential gene methylation. A similar idea has been put forward for reptiles with temperature-dependent sex determination, in which temperature influences gene methylation status [Matsumoto et al., 2013]. The estrogen synthesized by aromatase enzyme must act locally in the embryonic gonads, activating estrogen receptor a, leading to transactivation of target genes. Those targets in embryonic chicken gonads have not been defined, despite the key role for estradiol.

In the past 5 years, several lines of evidence have shown that the cell adhesion and signaling factor, β -catenin, is required for ovarian development in mammals [Maatouk et al., 2008; Nicol and Yao, 2014]. In the developing mouse ovary, but not the testis, the signaling molecules R-spondin1 and Wnt4 stabilize β -catenin, leading to its nuclear translocation and activation of target genes. All 3 molecules are required for proper ovarian differentiation in mammals [reviewed in Chassot et al., 2008, 2014]. This pathway appears to be conserved in the chicken embryo, as R-spondin1, Wnt4 and β-catenin are all enriched in the cortical region of the female gonad (fig. 8) [Smith et al., 2008; Ayers et al., 2013b]. In mammals, loss of both FOXL2 and components of the R-SPO1/WNT4/β-catenin pathway lead to potent female-to-male sex reversal [Ottolenghi et al., 2007; Auguste et al., 2011]. Knockdown and overexpression of these genes in chicken will confirm a role in gonadal sex differentiation. How the R-SPO1/ WNT4/β-catenin and FOXL2/CYP19A1 pathways intersect to control ovarian differentiation is currently not known. Indeed, in the chicken embryo, there are currently few proven functional links between the known or putative gonadal sex differentiation genes.

Teranishi et al. [2001] described a reiterated sequence on the chicken Z sex chromosome that they called MHM (male hypermethylated). As the name indicates, this sequence is hypermethylated and transcriptionally silent in ZZ male cells (in the gonad and throughout the body). It is hypomethylated in female cells (ZW), where it encodes a long noncoding RNA that coats the Z close to the site of its transcription. This is reminiscent of the mammalian Xist RNA, which coats one X chromosome as part of the X-inactivation phenomenon (dosage compensation). MHM may be involved in localized dosage compensation in chicken (although widespread dosage compensation appears to be absent, discussed below). In chicken, this would involve upregulation of neighbouring genes in the single Z of females, as opposed to repression by Xist on the mammalian X. However, MHM RNA coats the female chicken Z sex chromosome very close to the DMRT1 locus, and it has been suggested that it may contribute to the lower level of DMRT1 expression in females as part of the avian sex (gonadal) determination mechanism (fig. 8) [Teranishi et al., 2001]. Indeed, more recently, we found that overexpression of MHM in male chicken embryos could disturb embryonic development, including reducing the expression of gonadal DMRT1 [Roeszler et al., 2012]. The potential role of MHM in localized dosage compensation and perhaps gonadal sex differentiation requires further study. The fact that the sequence is expressed from the single Z of females and in ZZW triploid birds, not in males nor in ZZZ triploid males, points to the existence of a W-linked factor in triggering its demethylation and expression. No such factor has been identified on the W chromosome as yet [Ayers et al., 2013al.

One of the major issues that has arisen in the field is the definition of sex determination and the relative contribution of the sex chromosomes and sex steroid hormones [Arnold et al., 2012]. The traditional view of sex determination in birds and other vertebrates is that the sex chromosome complement controls embryonic differentiation of the gonads into ovaries or testes, and these organs then secrete hormones that feminize or masculinize the rest of the body (estrogen and progesterone in females, and testosterone and anti-Müllerian hormone in males). However, a growing body of evidence challenges this view, supporting the idea that the sex chromosomes can have direct, cell-autonomous effects upon sexual phenotype. This has been described in mice, in which the sex chromosome constitution, independent of gonadal sex, can cause sex differences in metabolism. It has been argued that the 'gonad-centric' view of sex determination is inappropriate and that multiple parallel pathways may regulate sexual development in different tissues, integrating to yield the overall sex of an individual [Arnold, 2012]. Such pathways might be controlled by a sex chromosome that is present in one sex and not the other. An example is the Y chromosome of therian mammals, expressing the Sry gene that induces testis formation during embryonic life. Other genetic differences between the sexes can come from other sources, such as X-linked genes that escape Xinactivation in mammals, or epigenetic differences such as genomic imprinting. Sex determination may therefore occur in different ways in different tissues, with gonadal sex differentiation being one aspect - one of the most visible aspects - of sexual differentiation [Arnold et al., 2013].

This idea is especially relevant to avian sex, because there is no chromosome-wide dosage compensation mechanism, such that males (ZZ) have, on average, higher levels of Z-linked gene expression compared to females [Ellegren et al., 2007; Itoh et al., 2007; Melamed et al., 2009]. However, this varies on a gene-by-gene basis, so that there is variable dosage compensation or no compensation of Z-linked genes that differs between tissues and life cycle stage [Mank and Ellegren, 2009; McQueen and Clinton, 2009]. The chicken Z sex chromosome is enriched for genes associated with sex and reproduction [Storchová and Divina, 2006], while genes associated with testis expression are overrepresented on the chicken Z chromosome [Ellegren, 2011]. These observations could explain avian sex determination being dependent upon sex-biased expression of different Z-linked genes in different tissues, consistent with the modern view of sex determination outlined above and expounded by Arnold et al. [2012]. Recent findings support the idea that sex determination in the chicken may be at least partly cell autonomous, involving direct effects of the sex chromosomes in different tissues, and supporting the idea of multiple parallel pathways underlying sexual development. Zhao et al. [2010] described 3 gynandromorphic chickens showing striking bilateral asymmetry, with male features on one side of the body (large wattle, spur and large muscle mass) and female features on the other (small spur and wattle, less breast muscle mass) (fig. 9). The 'male side' contained primarily ZZ cells, while the female side contained 40-60% ZW cells. The gonads of these birds reflected the relative contributions of ZZ and ZW cells, so that gonads with predominantly ZZ cells were testis-like, while those with predominantly ZW cells were ovarian-like [Zhao et



Fig. 9. Gynandromorphic chicken, showing male features on the left (large spur, wattle and breast muscle) and female features on the right (smaller spur, smaller wattle and smaller breast muscle). Reproduced with permission from Zhao et al. [2010].

al., 2010]. These observations suggest direct cell-autonomous effects of the sex chromosomes within the various tissues. The authors also generated chimeric embryos by explanting GFP-labeled presumptive gonadal tissue into opposite sex recipients and found that donor cells retained their sexual identity (expressing male or female marker genes) and were not incorporated into the gonadal tissues of the hosts. This again supports the notion that avian somatic cells have an inherent cell-autonomous cell identity (labeled CASI) that is largely independent of hormonal signaling. The factor/s driving CASI are not known, but it cannot involve DMRT1 in all tissues, as the expression of this gene is limited to the urogenital system. It could be driven by one (or more) of the hundreds of Z-linked genes in chicken that are not subject to dosage compensation and therefore show sexually dimorphic expression. Alternatively, a W-linked gene/s could be involved. Early chicken embryos show sexually dimorphic gene expression (from both sex chromosomes and autosomes) well prior to gonadal sex differentiation [Lee et al., 2009; Zhao et al.,

2010; Ayers et al., 2013a]. This supports CASI, and the idea that sexual differentiation pathways are engaged in avian embryos very early in development. As mentioned above, there is still no clear evidence that the W sex chromosome encodes a 'female' or ovary determinant, although genes associated with female fertility are reported to be enriched on the chicken W chromosome [Moghadam et al., 2012].

It seems clear that hormones can override CASI, because exogenous estrogens or estrogen antagonists can quite extensively sex reverse avian embryos [reviewed in Vaillant et al., 2001]. This suggests a potential interaction between cell-autonomous factors and hormonal signaling. In the case of the gynandromorphic birds, it is possible that factors responding to sex hormones are sexlinked, hence accounting for the different development of sex characters on the left and right sides of the body, which had mainly ZZ or ZW genotypes (fig. 9). Androgen and estrogen receptors are not sex-linked in chicken, although other factors influencing the response to hormonal signaling might be located on the Z and/or W sex chromosomes.

In summary, the past 10 years have seen several advancers in our understanding of sex determination and gonadal sex differentiation in the chicken embryo. Although the basic mechanism (Z dosage or dominant W) remains elusive, an increasing body of evidence points to the importance of the DMRT1 gene for testicular differentiation. Some new genes have emerged such as HEMO-GEN, and some interactions in the pathway have been developed (DMRT1 inhibition of CYP19A1, for example). Despite several efforts in several labs, no clear Wlinked female/ovary determinant has been identified, beyond those previously found to be expressed but for which a function has not been proven (HINTW, FET1, FAF) [Ayers et al., 2013a, b]. We carried out a thorough RNA-seq study of very early chicken blastoderms and day-4.5 (HH25) gonads, but did not detect a clear candidate ovary determinant [Ayers et al., 2013a]. Most of the 40 or so W-linked genes have Z homologues, to which they have very high sequence homology. The so-called dominant W hypothesis therefore remains open-ended. The discovery that birds lack a chromosome-wide system of dosage compensation together with the discovery of gynandromorphic chickens points to the possibility that different sex-linked genes, in different combinations, may underlie sexual differentiation in different tissues during development. The lack of dosage compensation provides a good basis for CASI, if one or a combination of Z-linked factors tips sexual development towards the male versus female pathways.

Schmid et al.

Avian Epigenetics

(Prepared by H. Zhou)

The draft reference sequence of the chicken (*Gallus gallus*) genome was completed in 2004 [Wallis et al., 2004]. Spatial, temporal and state-specific gene expression differences are a major contributor to phenotypic variation in organisms [Pastinen, 2010]; however, a decade later, like in humans and mice, a mechanistic understanding of gene expression regulation (when, where, and how much) in chickens remains to be elucidated. While genetic variations across individuals in a population may partially contribute to the gene expression differences among them, a major question remains: how can an almost identical copy of the genome within an individual be able to have cell-specific expression profiles in more than 200 different cell types in the chicken?

It became clear that the epigenome plays a significant role in directing the unique gene expression programming in each cell type at different development stages. The epigenome, a second dimension to the genome, provides pivotal insights to gene regulation besides the genomic sequence [Rivera and Ren, 2013]. The epigenome refers to sequence-independent processes that can modulate the functional output of the genome [Sarda and Hannenhalli, 2014]. It consists of chemical changes to DNA or covalent modifications of histone proteins, chromatin accessibility, noncoding RNA localization, and higher-order chromatin architecture (nucleosome positioning and occupancy, 3D chromatin structure) [Bernstein et al., 2007; Berger et al., 2009; Bonasio et al., 2010].

Fueled by the rapid development of next-generation sequencing (NGS) and the significant drop of cost rate in the past decade (even faster than Moore's Law used in computer technologies) [Koboldt et al., 2013], the rate of growth in datasets and publications of epigenomics have drastically increased, especially in humans, mouse and model organisms. Consequently, a few international consortia in the field of epigenomics have been established including the Encyclopedia of DNA Elements (ENCODE; http://encodeproject.org/) [ENCODE Project Consortium, 2012], the Roadmap Epigenomics Project (http://www.epigenomebrowser.org/) [Bernstein et al., 2010], the International Human Epigenome Consortium (IHEC; http://ihec-epigenomes.org), and modEN-CODE [modENCODE Consortium et al., 2010]. Although progress in the avian epigenome is very limited, recent international efforts on the functional annotation of animal genomes (FAANG) including chickens have

been initiated [The FAANG Consortium, 2015]. It is expected that epigenome data generated in the next few years will greatly contribute to our understanding of the genome-phenome relationship and eventually enhance genome information, enabling approaches to improve the efficiency, sustainability and biosecurity of poultry production systems.

In this review, recent advancements of technologies, advantages and disadvantages of each technology and progress in different areas of avian epigenomics as well as challenges and future directions in this field will be discussed.

DNA Methylation

One of the well-studied DNA modifications is methylation at its 5th carbon (5mC), which plays an important role in imprinting, retrotransposon silencing and X chromosome inactivation in mammals [Bird, 2002]. There are 3 main techniques in measuring DNA methylation: (1) methyl-sensitive restriction enzyme digestion (MREseq), (2) affinity enrichment of methylated DNA fragments (MeDIP-seq with an antibody and MBD-seq with a methyl-binding domain), and (3) bisulfite sequencing (MethylC-seq) and reduced representation bisulfite sequencing (RRBS) with enzyme digestion [Bock, 2012]. Both MRE-seq and MeDIP-seq and MBD-seq have a limited resolution and relatively low cost compared to MethylC-seq. The resolution of MRE-seq is dependent on the frequency of enzyme cut sites and can be improved with multiple enzymes, while the resolution of MeDIP-seq is highly related to the DNA fragment size, CpG density and quality of antibody. MethylC-seq with single-base resolution is generally accepted as the gold standard for DNA methylome profiling, although potential PCR biases may skew 5mC quantitation [Laird, 2010].

Chicken methylomes at different tissues have been characterized using MeDIP- and MRE-seq [Li Q et al., 2011; Hu et al., 2013; Tian F et al., 2013]. Liver and muscle methylomes in both red junglefowl and broiler were generated using MeDIP. The hypermethylations were observed in gene body regions and repetitive sequences, while promoter regions and 5' end regions were hypomethylated [Li Q et al., 2011]. Whole-genome methylation profiling between non-infected and Marek's disease virus (MDV)-infected thymus in 2 genetically distinct genetic lines that confer different resistances to MDV infection were studied using MRE-seq [Tian F et al., 2013]. A higher DNA methylation level was observed in the MD-susceptible line, while a lower DNA methylation level was found in the MD-resistant line with MDV

infection [Tian F et al., 2013]. Recently, MethylC-seq has been used to generate single-base resolution methylome in the lungs of 2 distinct genetic lines (Fayoumi and Leghorn). Like in other species, DNA methylation was found throughout the chicken genome but rarely occurred in CpG island promoters [H. Zhou, pers. commun.]. More than 90% CpG sites were detected using MethylC-seq compared to MRE-seq (32%) [Tian F et al., 2013].

In addition, the role of DNA methylation on dosage compensation has been investigated. Genome-wide methylation and gene expression analysis revealed few significant differences in the promoter DNA methylation, although hundreds of genes were differentially expressed between male and female brains. This result suggests that, unlike mammals, DNA methylation was not associated with dosage compensation in chickens [Natt et al., 2014].

DNA methylation is also crucial in primordial germ cell (PGC) reprogramming. Jang et al. [2013] revealed more than 200 differentially methylated regions within mammalian homologues of imprinting and X-linked genes between male and female PGCs. By comparing PGCs with chicken embryonic fibroblasts (CEFs), the promoter regions of a dozen of differentially expressed genes between PGCs and CEFs were either hyper- or hypomethylated, indicating the potential role of methylation in lineage differentiation [Jang et al., 2013].

DNA methyltransferase genes (DNMTs) play important roles in the epigenetic regulation of DNA methylation at CpG sites [Okano et al., 1999]. The methylation levels of 3 chicken DNMTs (DNMT1, DNMT3a and DNMT3b) and their genetic variations across different genetic lines, generations, tissues, and age were studied, and the results suggested that a CpG to TpG transition in DNMT3b may be associated with genetic susceptibility to MDV infection in chickens [Yu et al., 2008].

Recent studies have shown that there are several intermediates during demethylation of 5mC to cytosine [Tahiliani et al., 2009]. The 5mC can be oxidized to 5-hydroxymethylcytosine (5hmC), 5-formylcytosine (5fC) and 5-carboxylcytosine (5caC) before being converted to cytosine by the ten-eleven translocation (TET) family of proteins (TET1, 2, and 3) [Pastor et al., 2013]. Subsequently, several sequencing methods have been developed to distinguish these intermediates. Oxidative bisulfite sequencing (oxBS-seq) and TET-assisted bisulfite sequencing (TAB-seq) were used to identify and quantify 5hmC [Booth et al., 2012; Yu et al., 2012]; fC-seq, fCABseq and 5fC-DP-seq for measuring 5fC; and 5caC-seq to detect 5caC [Raiber et al., 2012; Shen et al., 2012; Song et al., 2013]. It is expected that these new techniques will contribute new insights into the avian epigenome in coming years.

Chromatin Modification

Chromosomal DNA is wrapped around histone octamers comprised of H2A, H2B, H3, and H4 subunits. These histone tails at specific residues are subject to covalent modifications. More than 130 post-translational modifications and 700 distinct histone isoforms have been identified in humans [Tan et al., 2011; Tian et al., 2012]. There are a few well-studied modifications including methylation, acetylation, phosphorylation, and ubiquitination. These modifications can play both activation and repression roles in transcription regulation by controlling DNA accessibility or recruitment/elimination of other protein complexes [Kouzarides, 2007]. The Mouse ENCODE generated a map of nearly 300,000 murine cisregulatory sequences from 19 tissues and cell types [Shen et al., 2012]. Analyses that have integrated all this information together indicate that histone H3 lysine 4 trimethylation (H3K4me3), H3 lysine 27 trimethylation (H3K27me3), H3 lysine 27 acetylation (H3K27ac), and H3 lysine 4 monomethylation (H3K4Me1) are the most informative histone modifications [ENCODE Project Consortium, 2012]. A high abundance of H3K4me3 correlates with promoters of active genes and transcription starts [Shen et al., 2012], while H3K27me3 is a repressive mark, and increased levels of H3K27me3 are associated with promoters of inactive genes [ENCODE Project Consortium, 2012]. H3K27ac is a mark of active regulatory elements and may distinguish active enhancers and promoters from their inactive counterparts. H3K4me1 is a mark of regulatory elements associated with enhancers and other distal elements, but is also enriched downstream of transcription starts [ENCODE Project Consortium, 2012]. High levels of H3K27ac and H3K4me1 are associated with enhancer regions, and also correlate with areas of DNase I hypersensitive sites (DHSs) [Greer and Shi, 2012; Shen et al., 2012].

Chromatin immunoprecipitation followed by sequencing (ChIP-seq) has been employed to map genomewide DNA-associated proteins with modified histones by specific antibodies. Only a couple of studies have been conducted in chickens. Tissue-specific genome-wide landscapes of H3K4me3 and H3K27me3 in chicken liver, spleen and breast muscle were generated, and the correlation between enrichment of these 2 marks and mRNA expression in 3 tissues was analyzed. The results suggested a conserved relationship like in mammals: H3K4me3 positively correlated with gene expression, while H3K27me3 negatively correlated with gene expression [Saelao and Zhou, 2015]. Luo et al. [2012] provided H3K4me3 and H3K27me3 profilings in MD-resistant and -susceptible chicken lines. Line-specific changes of these 2 histone modification marks with MDV infection were identified, which support the role of these 2 marks in the MD resistance in chickens.

Nucleosome Positioning and Occupancy

Chromatin is tightly packaged into basic repeat units of nucleosomes (147 bp of DNA and linker DNA). Nucleosome positioning can prevent or display specific DNA sequences for the regulation of transcription such as transcription factor recruitment and progression of RNA polymerase II [Rivera and Ren, 2013]. The most common method of genome-wide characterization of nucleosome positioning is micrococcal nuclease (MNase) for the digestion of chromatin followed by high-throughput sequencing (MNase-seq) [Bell et al., 2011], which can reveal the areas of the genome occupied by nucleosomes and other regulatory factors. MNase-seq has been used in a variety of organisms from yeast to humans to map nucleosome occupancy [Kaplan et al., 2009; Cui and Zhao, 2012; Gaffney et al., 2012]. However, there are several limitations to this method due to the nature of the assay: (1) MNase is AT-sequence biased; (2) resolution is low (not single-base); and (3) it is difficult to reproduce the experiment as the data is affected by the extent of chromatin crosslinking and the level of digestion [Tsompana and Buck, 2014].

Chromatin Accessibility

Chromatin openness is a prerequisite for the binding of transcription factor proteins to specific biochemically active regulatory elements in the genome such as promoters, enhancers, silencers, and insulators. A few highthroughput assays have been developed to map open chromatin including DNase hypersensitivity sequencing (DNase-seq) [Boyle et al., 2008] or formaldehyde-assisted identification of regulatory elements sequencing (FAIREseq) [Giresi et al., 2007]. Both assays require 100,000-10,000,000 cells. DNase-seq is time-consuming, labor-intensive, and requires a complicated sample preparation process, and FAIRE-seq has the disadvantage of a low signal-to-noise ratio [Tsompana and Buck, 2014]. Recently, a high-quality DHS map was generated in the MSB1 line, an MDV-transformed CD4+ T-cell line in chickens [He et al., 2014]. There were fewer DHS enrich-

Third Report on Chicken Genes and Chromosomes 2015 ments in CpG island-abundant regions. Furthermore, DHSs were strongly correlated with highly expressed genes in MSB1 cells [He et al., 2014].

To circumvent the weaknesses of MNase-seq and DNase-seq, Buenrostro et al. [2013] recently developed a novel method, ATAC-seq, that has several advantages over MNase-seq and DNase-seq. ATAC-seq is primarily based on hyperactive Tn5 transposase in probing open chromatin. First, it only requires 500–50,000 unfixed nuclei; furthermore, it has single-base resolution. Finally, it can map nucleosome positioning, chromatin accessibility, and TF footprints. The FAANG consortium is planning to use ATAC-seq as a standard for measuring chromatin accessibility, although the protocol in tissues using this assay requires more optimization.

3D Chromatin Architecture

Besides 1-dimensional linear genomic sequence, chromatin also has high-order structures such as the 30nm fibers and 700-nm mitotic chromosomes, which have more functions than simple compaction. Orchestrated activities among linearly distant elements can be accomplished through short- and long-range DNA interactions. Like histone modifications and chromatin accessibility, 3D conformation/looping can regulate gene expression [Rivera and Ren, 2013]. Several assays have been developed in mapping high-order chromatin architecture depending on the number of loci examined and selected. Dekker et al. [2002] first introduced the chromosome conformation capture (3C) assay, then the circular chromosome conformation capture (4C), the chromosome conformation capture carbon copy (5C), the tethered conformation capture (TCC), a derivative of 3C, called Hi-C, and a variation of Hi-C, namely chromatin interaction analysis by paired-end tag sequencing (ChIA-PET) [van Steensel and Dekker, 2010; de Wit and de Laat, 2012; Sajan and Hawkins, 2012]. 3C and 5C are based on priori loci of interest. 3C detects the interaction frequency of one locus with another (one vs. one), while 4C measures the genome-wide interaction frequency of a single locus (one vs. all) [Zhao et al., 2006]. 5C measures thousands of anchors and baits (many vs. many) [Dostie et al., 2006]. Hi-C measures genome-wide interaction frequency without loci-specific enrichment (all vs. all) [Lieberman-Aiden et al., 2009]. Hi-C is considered as one of the most powerful and unbiased assays in measuring 3D structure.

Genomic Imprinting

As one of the most important germ-line-specific epigenetic modifications that usually results in imbalanced allelic expression between 2 parental alleles, genomic imprinting has been intensively studied in mammals. However, this phenomenon in chickens is still in debate. Several known imprinting genes such as insulin-like growth factor type 2 (IGF2) and IGF2 receptor (IGF2R) in eutherians and marsupials have been tested in chickens and revealed that both genes were biallelically expressed [Yokomine et al., 2001]. Subsequently, a 0.5-Mb ortholog of the mammalian imprinting region in the chicken was investigated and further showed that several imprinting regulatory elements in mammals such as H19 imprinting center and local regulatory elements were not identified in chickens, which suggests that imprinting may have evolved after the divergence of mammals and birds [Yokomine et al., 2005]. A recent study of transcripts of 20 embryos in 2 genetically distinct chicken populations by RNA-seq identified 79 potential imprinting SNPs. Unfortunately, none of them could be confirmed [Frésard et al., 2014].

Perspectives and Challenges

In summary, avian epigenomics is still in its infancy. With more advanced high-throughput technologies, robust assays developed from human, mouse and model organism ENCODE projects, and significant cost reductions in NGS, in addition to the launching of the international FAANG consortium and great interest in functionally annotating, the chicken genome will prompt the next wave of exponential growth of epigenome data in the chicken. However, significant challenges remain. Firstly, data analysis is still one of the most difficult hurdles that the community must face and work on. In order to decipher functional information underlying the genetic blueprint in the genome, new bioinformatics tools need to be developed to integrate a variety of data types such as ChIP-seq, RNA-seq, DNase-seq, MethylC-seq, Hi-C, etc. Computational algorithms and novel statistical models also need to be implemented to appropriately analyze these 'omic' data. More importantly, few scientists with the required expertise available have signified an urgent need to train next-generation computational biologists, especially in the agricultural field. Secondly, new techniques need to be invented to address some limitations of each of the assays discussed above. An assay can be developed to characterize more than one mark. For example, nucleosome occupancy and methylome sequencing (NOMe-seq) provides dual epigenomic information in

both nucleosome position and DNA methylation abundance from a single molecule [Kelly et al., 2012]. Singlemolecule and single-cell assay is another direction of technology development. A specific cell type has a distinct epigenomic state. In order to dissect cell-specific regulatory mechanisms, single-cell populations with limited quantity need to be purified from tissues usually consisting of heterogeneous cell populations. A couple of new techniques are already underway to tackle this issue. Readers can have more details in a recent excellent review by Hyun et al. [2015]: ChIP-nano (nano-scale ChIP, only requires less than 50,000 cells) [Adli and Bernstein, 2011], SCAN (single chromatin molecule analysis in nanochannels) [Murphy et al., 2013], SMRT (single-molecule realtime without amplification step) sequencing [Flusberg et al., 2010]. Thirdly, many studies have supported the histone code hypothesis 'that multiple histone modifications, acting in a combinatorial or sequential fashion on one or multiple histone tails, specify unique downstream functions' [Strahl and Allis, 2000]. Re-ChIP or SeqChIP (sequential ChIP) has been developed to map the genomic locations of combinations in chromatin features [Furlan-Magaril et al., 2009; de Medeiros, 2011]. Finally, great coordination within the international scientific community is crucial to maximize 'outputs' with minimum 'inputs' as each group has limited resources and funding support. Indeed, a couple of projects under the umbrella of the FAANG consortium are already underway to profile more epigenomes from different tissues at different development stages in different breeds (UC Davis group funded by the USDA NIFA, and FR-AgENCODE funded by the INRA). We expect hundreds of epigenome maps with features of key epigenetic marks from a variety of chicken tissues will be generated in the coming years, which will contribute vital insights into the second and third dimensions of the chicken genomic sequence, and fulfill our promise in fully understanding phenotypes and mechanisms of inheritance not yet revealed by primary genomic sequences.

Structural Variation and Copy Number Variation in Poultry

(Prepared by R.P.M.A. Crooijmans and M.A.M. Groenen)

To understand phenotypic variation in farm animals and in poultry in particular, it is essential to define all potential genomic variation within a genome. Most of the emphasis has been on SNP variation [Wong et al.,

Table 14. Overview of the different studies performed for CNV of	detection
--	-----------

Study types	Platform			Analysis			Population un	Reference		
	NimbleGen	Agilent	Illumina	ref genome build	coverage	detected CNVs/CNVRs	breeds/lines	animals/breed	total No. of animals	
SNP			60K	3	WG	204	2	385/361	746	Jia et al., 2013
SNP			60K	3	WG	137	1	1,310	1,310	Zhou W et al., 2014
SNP			60K	4	WG	271-188	2	203/272	475	Zhang H et al., 2014
aCGH	385K			3	WG	96	3	2-4	10	Wang X et al., 2010
aCGH	180K			3	GGA1 (60 Mb)	477	3	32/39/39	110	Abe et al., 2013
aCGH	385K			3	WG	45	4	1-2	6	Luo J et al., 2013
aCGH	385K			4	WG	281	5	2	10	Han et al., 2014
aCGH		400K		3	WG	130	4	6	18	Wang et al., 2012
aCGH		244K		3	WG	1,156	15	3-8	64	Crooijmans et al., 2013
aCGH		244K		3	WG	273	2	6	12	Abernathy et al., 2014
NGS			120-bp paired-end	3	WG (23-24×)	8,839	2	1	2	Fan et al., 2013
NGS			100-bp paired-end	4	WG (8.4-12×)	8,840	12	1	12	Yi et al., 2014

2004] especially in chicken, the first farm animal whose genome was sequenced [International Chicken Genome Sequencing Consortium, 2004]. This resulted in the development of medium-high density SNP chips such as the Illumina 60K beadchip [Groenen et al., 2011] and, more recently, the 600K Affymetrix Axiom Genome-Wide Chicken Genotyping Array [Kranis et al., 2013]. In addition to single base alterations, genomes contain numerous other variations collectively referred to as structural variants (SVs). SVs include small insertions and deletions (InDels) of a few base pairs in size, large InDels of thousands to sometimes hundreds of thousands base pairs in size, inversions and translocations. Rather arbitrary, copy number variation (CNV) has been defined as duplications and losses of DNA segments of 50 bp and longer. CNVs have been implicated to play a role in a wide variety of phenotypic traits in many species including chicken as discussed below.

Detection of SVs

There are basically 3 different approaches to analyze CNVs, which are based on (1) medium-high density SNP arrays, (2) high-density arrays of genomic probes (aCGH) and (3) next-generation sequencing. For the first method, so far, only the Illumina chicken 60K SNP beadarray has been used to detect CNVs by identifying groups of adjacent SNPs by means of the software PENNCNV [Wang K et al., 2007]. The number of CNVs detected in these studies (table 14) varied from 137 to 271. The limited number of CNVs is mainly due to the low SNP density (60K) of this SNP array. A more direct

way to analyze CNVs is by using specifically designed comparative hybridization arrays (aCGH), where the difference in hybridization signal between a test sample and a reference sample is scored. Two commercial aCGH platforms are available, the NimbleGen and Agilent platforms, respectively. The NimbleGen arrays available for chicken have a probe density of 180K or 385K, and the Agilent arrays for chicken have a probe density of 244K or 400K, respectively (table 14). The number of CNVs detected not only depends on the number of probes on the array but also on the number of samples and the diversity of populations under study. The number of detected CNVs with aCGH varied from 45 to 1,156 and is higher than that obtained with the 60K SNP chip analysis (table 14). The third method uses whole-genome re-sequencing data, and is a very powerful tool to detect CNVs. However, this method is very dependent on the quality of the reference genome and the available depth of the sequence data. The number of CNVs detected by the 2 studies that used this approach exceeds 8,000, a number that is mainly the result of an increase in the smaller size-range fraction of CNVs, due to the higher resolution of this method (table 14). One of the main advantages of the re-sequencing method is that all the different types of variation (both SNPs and SVs) can be identified simultaneously.

Impact of SVs on Phenotype

In many species, the impact of SVs and CNVs on specific phenotypes is still poorly studied. Nevertheless, a number of studies have reported CNVs as the underlying variation for specific qualitative traits. In pigs, the white

Trait	Gene	CNV, kb	Kind	Reference
Late feathering Dermal hyperpigmentation Pea-comb Dark brown plumage color	PRLR and SPEF2 EDN3 SOX5 SOX10	176 129 and 172 3.2 8.3	duplication and rearrangements duplication and rearrangements partial duplication deletion	Elferink et al., 2008 Dorshorst et al., 2011 Wright et al., 2009 Gunnarsson et al., 2011
Rose-comb	MNR2	7.4	inversion	Imsland et al., 2012

Table 15. Structural variation affecting traits in chicken

color is related to a CNV in the *KIT* gene [Wiseman, 1986; Giuffra et al., 2002], and in chicken several CNVs have been identified that affect specific traits. The partial duplication of a 180-kb fragment containing the genes *PRLR* and *SPEF2* on the Z chromosome is responsible for the late feathering phenotype [Elferink et al., 2008], while the Pea-comb phenotype is associated with a CNV in the *SOX5* gene [Wright et al., 2009]. Besides these 2 examples, 3 more CNV-trait relations have been identified (table 15) and many more are expected to be identified in the future.

SNPs and InDels – The Most Abundant Sources of Genetic Variations

(Prepared by A.A. Gheyas, C. Boschiero, and D.W. Burt)

Single nucleotide polymorphisms (SNPs) constitute the most abundant source of genetic variation in the vertebrate genome followed by short-length (≤100 nucleotides) insertions and deletions (InDels). These genetic variants are important drivers of genomic evolution. Since the publication of the Second Report on Chicken Genes and Chromosomes [Schmid et al., 2005], the number of SNPs and InDels reported for chicken has increased many folds. This rapid increase in the discovery rate of variants has been possible thanks to the development of a reference genome assembly for chicken and the advancements in high-throughput sequencing technologies with simultaneous development of bioinformatics approaches to mine large volumes of sequence data. Nevertheless, not only the discovery but also the characterization of these variants is essential to understand their physiological impacts, if any. In the following sections, we discuss the progress in the discovery and characterization of SNP and InDel variants and their current and future applications in research and poultry breeding.

Surge of Discovery of SNPs and InDel Variants from the Chicken Genome

During the write-up of this article, the dbSNP (build 140) reported about 11 million SNPs and InDels from the chicken genome. Major proportions of these variants were contributed by a few studies only. The first major genome-wide discovery of variants in chicken was reported by the International Chicken Polymorphism Map Consortium (ICPMC), and the seminal paper [Wong et al., 2004] was published in the same issue of the journal Nature that released the draft assembly of the chicken genome based on the sequencing of a red junglefowl [International Chicken Genome Sequencing Consortium, 2004]. The ICPMC reported a genetic variation map of 2.8 million SNPs and short InDels discovered by comparing the sequences of 3 domestic chicken breeds – a male broiler, a female layer and a female Chinese Silkie - with the reference genome from red junglefowl. To facilitate the application of these variants to avian genetics, genomics and evolutionary studies, Beijing Genomics Institute created a Chicken Variation Database (ChickVD) to display these in relation to various genomic contexts such as genes, cDNAs, chicken orthologs of human disease genes, quantitative trait loci (QTLs), and raw sequence traces [Wang et al., 2005].

While the ICPMC study performed the sequencing with shotgun method with coverage of only 0.25×, later studies reporting the discovery of a large catalogue of variants applied next-generation sequencing (NGS) approaches to re-sequence many more individuals with much higher depth of coverage. One major study reported over 7 million SNPs and 1,300 InDels by re-sequencing individuals from 8 different breeds – 2 commercial broiler lines, 1 high-growth line, 1 low-growth line, 1 line of Rhode Island Red, 1 obese strain, and 2 White Leghorn lines of which one was used for commercial purpose [Rubin et al., 2010]. To allow sequencing of multiple individuals from each of these lines and thus to obtain the frequency spectrum of the variants, the study adopted a

pool-seq approach where DNAs from several individuals (n = 8-11) within lines were pooled prior to sequencing. The sequencing was performed with coverage of 44.5×. The variants detected in this study were used to investigate signatures of domestication selection in chickens.

In order to develop a moderate-density genotyping array (60K), a USDA-funded project created reduced representation libraries from 4 distinct chicken populations - 2 broiler, 1 white-egg layer (WEL) and 1 brownegg layer (BEL) – by pooling DNAs from 25 individuals within each line [Groenen et al., 2011]. Variant discovery was performed based on Illumina sequencing of these reduced representation libraries at an intended coverage of $25-40\times$ and resulted in the detection of over 352,000 SNPs. Another recent study reported the discovery of about 7.6 million SNPs by whole-genome re-sequencing of only 2 domestic chickens (a Silkie and a Taiwanese native chicken) using Illumina sequencing technology with sequence coverage of 23-25× [Fan et al., 2013]. The identified variants were characterized in relation to their position and potential effects on genes and known QTLs and were used to identify regions of selective sweep.

So far, the largest re-sequencing and variant discovery project on chicken has been undertaken by the Roslin Institute of the University of Edinburgh that sequenced 243 chickens from 25 commercial and experimental lines from diverse breeds including 4 commercial broiler lines of composite breed origin; 6 WEL lines of White Leghorn origin; 6 BEL lines originating from several breeds such as White Plymouth Rock, Rhode Island Red and White Rock; 8 experimental inbred lines from the Institute of Animal Health (currently known as the Pirbright Institute); and 1 unselected experimental line of Brown Leghorn origin from the Roslin Institute [Kranis et al., 2013]. This study also applied the pool-seq approach to combine DNAs from several individuals (n = 10-15) within lines and performed the sequencing on Illumina platform with coverage of 8-17×. From the millions of SNPs discovered in this work, a high-density (600K) genotyping array was developed and made commercially available to facilitate genome-wide analyses in chicken. The project has also characterized over 15 million SNPs adopting a number of approaches with the goal to delineate potentially functional variants [Gheyas et al., 2015].

Although the advent of NGS approaches has accelerated the pace of variant discovery, the discovery of InDels is still lagging behind that of SNPs due to a number of intrinsic difficulties. Most of these difficulties are associated with mapping reads against the reference genome. For instance, the short reads covering InDel sites from



Fig. 10. Size distribution of InDels in chicken (source: dbSNP; Yan et al. [2014]; Boscheiro et al. [in press]).

NGS are often difficult to map to the correct genomic locations and require complex algorithms for gappedalignment [Li et al., 2008; Neuman et al., 2013]. Moreover, since the mapping algorithms align each read independently, variant callers tend to consider InDels as stretches of SNPs unless a local realignment of reads around the suspected InDel sites is performed [Albers et al., 2011; GATK, https://www.broadinstitute.org/gatk/ gatkdocs/org_broadinstitute_gatk_tools_walkers_indels_IndelRealigner.php]. Furthermore, the majority of the InDels occur in short tandem repeats which are difficult to map [Messer and Arndt, 2007; Madsen et al., 2008]. Due to these issues, along with the relatively low frequency of InDels, a higher sequencing coverage is needed to detect these variants [Mullaney et al., 2010].

The current dbSNP release (build 140) reports only about 439,000 InDels of which ~273,000 were generated from the study by the ICPMC [Wong et al., 2004]. Only recently, a study has reported about 1.3 million InDels by NGS analysis of single individuals from 12 diverse breeds including the red junglefowl, 7 Chinese indigenous breeds, and 4 commercial breeds (Cornish, Rhode Island Red, White Leghorn and White Plymouth Rock) [Yan et al., 2014]. In order to minimize the rate of false positives in InDel detection, this study retained only the consensus In-Dels called by 2 different variant callers, viz. SAMtools and GATK. Another recent study [Boschiero et al., in press] has characterized over 883,000 InDels based on the sequenced data from 16 commercial and experimental layer lines described in Kranis et al. [2013]. The same approach of taking consensus from 2 variant callers, SAMtools and Dindel, was adopted by this study as well. A large proportion of the



Fig. 11. Mean SNP and InDel densities across chicken chromosomes based on 17 million SNPs and 2.5 million InDels detected till date.

InDels (80%) detected in this study overlapped with the variants from Yan et al. [2014] serving as a validation for the majority of the InDels. Unlike the previous study, however, this work screened multiple individuals from each line by sequencing pooled DNA and is the first study to report the frequency spectrum of detected InDels. The study observed that with low sequencing coverage, NGS tends to detect mostly high-frequency variants. This result, along with a large proportion of common InDels across these 2 studies, indicates that most of the known InDels are high-frequency old variants, and it would take sequencing of many more individuals with higher coverage to detect most of the low-frequency variants. Analysis of the size distribution of the InDels suggests that most of the known InDels (89% insertion and 88% deletion) are small in size $(\leq 5 \text{ nucleotides; fig. 10}).$

Genomic Distribution of SNPs and InDels

The chicken genome consists of 38 pairs of autosomes and 2 sex chromosomes. The current build of the chicken reference genome (Gallus_gallus_4.0), however, represents only 29 autosomes, 2 linkage groups, the sex chromosomes and 14,093 unplaced scaffolds (http://www.ensembl.org/Gallus_gallus/Info/Annotation/assembly, accessed 10/11/2014). Since variant discovery in most cases was performed by aligning sequence reads against the reference genome, the detected variants mostly represent regions covered by the reference genome. Combining the chicken variants from different sources – i.e. the ones already in the dbSNP and those yet to be released in public domain – we found about 17 million SNPs (n = 16,890,433) and 2.5 million (n = 2,540,719) In-Dels. These numbers are comparable to those reported for other species in the public domain such as human (~63 million), mouse (~71 million) and cattle (~74 million), taking into account the fact that the chicken genome is about 2.5–3 times smaller than the genomes of these species. Counting the number of variants in each 1-kb non-overlapping window over the genome reveals the average densities of variants to be 16.22 ± 10.01 SNPs per kb and 2.43 ± 2.14 InDels per kb. About 4% of the 1-kb windows or ~51 Mb of the reference genome, however, did not have any variants at all.

Discovery of millions of variants has improved our understanding of the patterns of their genomic distribution across chromosomes. In chicken, the sizes of the autosomes are highly variable with 5 macro-, 5 intermediate and 28 microchromosomes [Burt, 2005]. The study by the ICPMC [Wong et al., 2004] suggested that SNP and InDel rates are independent of chromosome size, except for chromosome 16. However, when we recalculated the densities based on the known list of SNPs and InDels, a different picture emerged. This showed that the microchromosomes had the highest mean SNP density but the lowest InDel density among the 3 groups of chromosomes (fig. 11). The higher SNP density in microchromo-

	Wong et al., 2004	Fan et al., 2013	Gheyas et al., 2015	Yan et al., 2014	Boschiero et al., in press
Number of total variants, in millions Number of variants from individual lines, in millions	2.8 0.9-1.2	7.6 5.1–5.4	15.3 3.6-5	1.3 0.37-0.52	0.9 0.17-0.39
Proportion, %					
in genic regions	37	43	47	45	48
in coding regions	1.2	1.1 - 1.2	2.3	0.13	0.12
of nonsynonymous and stop-gain/loss SNPs	0.4^{a}	$0.3 - 0.4^{a}$	1.2 ^b	_	-
of frameshift and stop-gain/loss InDels	_	_	_	0.07	0.09
of non-frameshift InDels of nonsynonymous SNPs and non-frameshift InDels	-	_	_	0.05	0.06
with potential radical effects	26 ^c	11 ^c	49 ^{c, d}		29 ^d

Table 16. Summary of functional annotation of SNP and InDel variants from major studies

^a The estimates are based on variants detected within single lines.

^b The estimate is based on all the variants detected, many of which are private to single lines.

^c The predictions were performed with SIFT package.

^d The predictions were performed with PROVEAN package.

somes is expected due to their higher recombination rates [Burt, 2005] as studies on other species revealed that the polymorphism rate was positively correlated with the recombination rate [Begun and Aquadro, 1992; Nachman, 2001]. The lower rate of InDels in microchromosomes, however, is contrary to this paradigm but has been observed by several studies on chicken. This has been attributed to the high gene content of these chromosomes [Brandström and Ellegren, 2007; Yan et al., 2014; Boschiero et al., in press]. Since presence of InDels in the coding regions can have a more detrimental effect than that of SNPs, purifying selection against InDels possibly counteracts the effects of high recombination rates in microchromosomes. Chromosome 16 is another exception to the 'recombination-polymorphism rates' paradigm. This chromosome has one of the highest SNP densities (26 SNPs/kb) (fig. 11) and the lowest InDel density among all the chromosomes. The presence of highly variable major histocompatibility complex (MHC) genes is the probable cause of the high SNP rate [Wong et al., 2014]. On the other hand, the potential effect of purifying selection due to the presence of several important gene families such as MHC genes and the nucleolus organizer region possibly explains the low rate of InDels [Yan et al., 2014].

Compared to the autosomes, the sex chromosomes have ~3 times less SNP and InDel densities (fig. 11). The low polymorphism rate in the Z chromosome is possibly a function of multiple factors including skewed reproductive success among males leading to low male effective population size, selection on sex-linked characters, and lower recombination rates compared to autosomes [Sundström et al., 2004]. On the other hand, the W chromosome is the sex-limiting chromosome in chicken, and its reduced genetic diversity is the result of selection and the complete lack of recombination outside the pseudoautosomal region [Berlin and Ellegren, 2004].

Functional Characterization of SNPs and InDels

The first tier of functional characterization of variants often entails investigating their location in relation to coding and noncoding functional elements in the genome. This, however, demands a comprehensive annotation of the reference genome for functional elements. Although the chicken genome is well-annotated for protein-coding genes, the annotation of functional noncoding elements (fNCEs) is still poor. Even though protein-coding genes cover ~45% of the chicken reference genome, the actual coding part constitutes only ~1.5%. For human, a massive research consortium (ENCODE) has recently predicted that >80% of the human genome is involved in some biochemical functions, and most of these elements belong to noncoding regions, thereby emphasizing their importance [ENCODE Project Consortium, 2012]. Until a similar database of fNCEs for chicken is developed, only a small proportion of variants can be annotated.

Most studies on chicken have, therefore, attempted to annotate variants in the functional context of proteincoding genes and predict the effects of only coding vari-

ants. The study by ICPMC [Wong et al., 2014] showed that ~37% of the variants fell within genes – mostly protein coding (based on Ensembl database) – while only 1.2% fell within coding regions. Later studies, however, predicted a larger proportion (43–48%) of detected variants to be present within genic regions, possibly reflecting the continual improvement of annotation of the chicken genome (table 16). Only about 1.1–2.3% of the SNPs and 0.12–0.13% of the InDels were reported to be within coding regions by different studies (table 16).

The SNPs within coding regions can be categorized as synonymous (does not alter amino acid), nonsynonymous (alters amino-acid sequence) and stop-gain/loss (causing the gain or loss of stop codons) types. The estimated proportion of amino-acid altering variants ranged between 0.3 and 1.2% in different studies (table 16). In comparison, the proportion of InDels within coding regions has been found to be very small irrespective of their categories such as: non-frameshift InDels, which are multiples of 3 nucleotides and do not change the protein-coding frame, constituted only 0.05-0.06% of variants; frameshift InDels, which are not multiples of 3 nucleotides and hence disrupt the reading frame, account for about 0.07-0.09% along with stop-gain/loss types. This indicates that the InDels in coding regions can be much more disruptive compared to SNPs and are therefore selected against.

Amino-acid altering variants are not always deleterious or radical in impact. Numerous approaches have been described to classify these variants, particularly the nonsynonymous SNPs, on both quantitative and discrete scale, by taking into account the evolutionary conservation and other properties of the proteins [Cooper and Shendure, 2011]. Major variant discovery studies on chicken have predicted the effects of nonsynonymous SNPs using various tools such as SIFT [Ng and Henikoff, 2003] and PROVEAN [Choi et al., 2012], and these studies predicted about 11–49% of the nonsynonymous variants may have radical impacts on protein functions (table 16).

One recent study by Boschiero et al. [in press] has also predicted the effect of non-framehsift InDels using the PROVEAN package and has shown that ~29% of these InDels were expected to be radical or deleterious in their effects. Such categorization can be very useful in shortlisting candidate functional mutations in studies involving specific disease or traits.

To go beyond the functional characterization of coding variants only, different studies have also utilized other resources and methods to delineate potentially functional mutations. Several studies, for instance, have attempted to analyse signatures of selection in the genome using the detected variants, not only to identify regions under selective sweep but also as a means to delineate variants that may be functionally important [Rubin et al., 2010; Qanbari et al., 2012; Fan et al., 2013; Gheyas et al., 2015]. Yan et al. [2014] have annotated InDels in the context of known QTLs. Few studies have overlaid the variants with most-conserved elements in the chicken genome as mutations within these elements are likely to have functional effects [Gheyas et al., 2015; Boschiero et al., in press]. Furthermore, the study by Gheyas et al. [2015] also attempted to predict the effects of variants from different categories, i.e. UTR, intronic, ncRNA and coding, on the secondary structures of RNAs - both mRNA and ncRNAs - as these may have functional implications. Several studies have looked for enrichment of the potentially functional variants within gene clusters/ networks to associate these variants with specific physiological functions [Fan et al., 2013; Yan et al., 2014; Gheyas et al., 2015; Boschiero et al., in press]. With rapid advancements in the transcriptome profiling (e.g. RNAseq) and continuous improvements in our understanding of fNCEs in the genome, it will soon be possible to perform a comprehensive annotation of noncoding variants.

Current and Future Applications of SNP and InDel Variants

The millions of SNP and InDel variants discovered over the last few years from chicken genome have found many applications in both research and breeding. One major outcome of the availability of a large catalogue of variants has been the development of SNP genotyping arrays to facilitate high-throughput screening of many individuals. Although genotyping arrays of various densities have been developed by different groups [Gheyas and Burt, 2013], detailed descriptions are available for only 3 arrays: a low-density 3K Illumina array [Muir et al., 2008a, b], a medium-density 60K Illumina array [Groenen et al., 2011], and a high-density 600K Affymetrix[®] Axiom[®] array [Kranis et al., 2013]. Only the 600K array is commercially available, while all others are proprietary and are not available for wider use. Below, we describe some of the major areas where high-throughput variant data, either from genotyping arrays or directly from re-sequencing, has been or can be used.

Phenotype-Genotype Association

A major use of molecular markers has been in the detection of QTLs associated with various traits or diseases. Traditionally, QTL analysis in chicken has been performed using only a handful of markers, generally micro-

satellites. Since most of these studies used low-density marker maps, the QTLs were often detected with very large confidence intervals, making it difficult to identify important genes or causal mutations [Zhang H et al., 2012]. Availability of a dense set of markers has opened up the opportunity to map QTLs or locate putative causal mutations with greater accuracy by fine mapping of QTLs or applying genome-wide association studies (GWAS). As a result, we have seen a surge of research papers on GWAS published just in the last few years on a wide range of traits in chicken, e.g. growth [Xie et al., 2012], fatness [Abasht and Lamont, 2007], meat quality [Sun et al., 2013], egg production and quality [Wolc et al., 2014], egg shell strength [Ghebrewold et al., 2014], resistance towards diseases and pathogens [Luo CL et al., 2013, 2014; Wolc et al., 2013], pigmentation [Park et al., 2013], etc.

Although genotyping arrays have so far been used for GWAS, they often fall short in identifying the actual causal mutations. This is because the genotyping arrays only search for association from among pre-defined panels of markers, which generally includes only common SNPs leaving out the low-frequency variants or mutations of other types that might be important [Gheyas and Burt, 2013]. As a consequence, it has been observed that the regions detected through GWAS often explain only a small part of the genetic variance of a trait [Maher, 2008]. With increasing affordability of NGS, it is now possible to perform association analysis on all the variants detected from whole-genome or targeted sequencing of candidate regions. A number of recent studies have used NGS to identify candidate functional mutations from known QTL regions such as those affecting adiposity and fatness [Roux et al., 2014; Moreira et al., 2015], meat quality [de Koning et al., 2012] and muscle deposition [Godoy et al., 2015] in chicken.

A number of recent studies have used the variant data either from genotyping arrays or from NGS platforms to identify genomic regions that are under selection pressure [Rubin et al., 2010; Elferink et al., 2012; Qanbari et al., 2012; Fan et al., 2013; Gholami et al., 2014]. Although such analyses do not require the use of phenotypic information, they can generate valuable understanding about genotype-phenotype relationship.

Genomic Selection

Genomic selection is the latest advancement in marker-assisted selection technology where genotypes from dense marker panels are used to estimate total genomic breeding values (GEBVs) of animals for various traits, and these GEBVs are used for selection of breeding candidates [Meuwissen et al., 2001]. Genomic selection promises to increase the rate of genetic gains substantially by capturing all or most of the genetic variance associated with a trait, allowing accurate estimation of breeding values without phenotypic records or progeny information, and facilitating selection at an early stage leading to shortening of the generation interval [Meuwissen et al., 2001]. Several studies have reported the application of genomic selection in commercial poultry breeding using medium- to high-density genotyping arrays [Avendaño et al., 2010; Chen et al., 2011; Fulton, 2012; Preisinger, 2012].

Population Genetics

High-throughput genotype data on a large panel of markers can be very useful in answering a wide range of questions related to population genetics. A number of studies have utilized dense marker sets to investigate the linkage disequilibrium (LD) profiles in chicken [Aerts et al., 2007; Andreescu et al., 2007; Megens et al., 2009; Elferink et al., 2010; Qanbari et al., 2010]. The LD and haplotype maps are very useful resources as they can provide unique insights into the genetic characteristics of a population, shed lights on the recombination patterns in the genome, help identify regions under positive selection, and guide the design and analysis of GWAS and GS [International HapMap Consortium, 2005]. For example, a comprehensive study on several layer populations revealed that the magnitude of LD in WEL populations was greater and extended over a longer distance compared to that in BELs [Qanbari et al., 2010]. This was attributed to the intense long-term selection pressures on WEL lines and their narrow genetic base in contrast to the BELs. Another study on commercial broiler lines observed a significant overlap of LDs of marker pairs between related populations suggesting that haplotypes are shared among these lines [Andreescu et al., 2007]. Such information has important implications as it can help predict the transferability of genetic parameters (e.g. GEBVs, or QTL effects) from one population to another. Further studies have shown that LD in macrochromosomes extended over longer genomic distances compared to that in microchromosomes, implying that a denser panel of markers would be needed to identify QTLs on micro- than on macrochromosomes by association mapping [Megens et al., 2009; Qanbari et al., 2010].

SNP markers have also been reliably used to investigate genetic diversity within populations and assess biodiversity. Using about 3K SNPs distributed over the genome, Muir et al. [2008a, b] showed that pure chicken

lines from commercial stocks lacked 50% or more of the genetic diversity observed in ancestral breeds and suggested that SNP markers can be used as sentinels of biodiversity. A number of other studies on chicken have confirmed the usefulness of SNPs in biodiversity analyses by showing their ability to reliably cluster related individuals into groups or in expected phylogenetic orders [Hillel et al., 2007; Twito et al., 2007; Kranis et al., 2013; Mekchay et al., 2014].

Concluding Remarks

With the availability of the chicken reference genome and advancements in sequencing technologies, we have seen a surge of discovery of SNP and InDel variants in just few years. These variants are being used for a wide range of purposes, both in research and breeding. The full benefits of these variants, however, can only be reaped with their better characterization such as having knowledge about their frequency profiles in different populations and possible functional effects. With the rapid improvements of the annotation of the chicken genome, interpretation of functional impacts of variants, especially of the noncoding ones, would be possible in the very near future.

Genetic Diversity of Village Chickens

(Prepared by T.T. Desta, R.A. Lawal, and O. Hanotte)

Domestic chickens are represented by 3 main gene pools: village, commercial and fancy (traditional) chicken. Population size of village chicken comes second to commercial ones with a total of at least 3 billion birds [Roberts, 1997]. Across the world, they represent a major economical and sociocultural asset making the livelihood of millions of households. Village chickens are the most ancient types of domestic chicken. They are found and adapted to nearly all agro-ecological systems. Village chickens represent the main repository of domestic chicken genetic diversity. Untangling this diversity is increasingly contributing to our understanding of the history of our agricultural societies, while the breeding structure and ancient history of village chicken make them ideal resource populations to unravel at genome level the impact of natural and human selections and to identify the genetic control of Mendelian and quantitative traits.

Village chickens have been named under a range of synonyms. They are commonly referred as native, local, scavenging, free range, non-descript, smallholders' or simply indigenous fowl. These synonyms invariably indicate the traditional and free-range management systems under which they have been maintained since domestication. In this respect, village chickens are mainly reared by subsistence or smallholder farmers across Africa, South America and Asia where they fulfil a range of sociocultural and economic roles [e.g. Desta and Wakeyo, 2012]. In Europe, village chickens largely do not exist anymore having been replaced by traditional chicken populations with specific plumage and/or morphological standards [e.g. Dana et al., 2011].

Following domestication, village chickens have been shaped by natural and human selection, with survival (e.g. following infectious diseases and predation challenges) a major adaptive trait. Scavenging is the main feed strategy, and mating is polygamous and often uncontrolled. Population structure of village chicken is therefore largely panmictic in contrast to commercial and fancy breeds. Within village chickens gene flow is expected to be high but influenced by several demographic factors including sex ratio, density, flock size, and foraging behaviour. Genetic divergence between village chicken populations in the same geographic area might be low due to the easiness of transporting such a small livestock species. Also, the use of common marketsheds (villages sharing a common trading network) facilitates genetic exchange between populations [Desta, 2015]. It has been shown that genetic divergence and diversity of village chicken populations within a geographic region is rather shaped by historical, ecological and sociocultural factors than geographic distances [e.g. Leroy et al., 2012].

Village Chickens and Their Phenotypic Diversity

Increasingly, a number of studies are documenting the phenotypic diversity of village chicken [e.g. Dessie et al., 2011; Desta et al., 2013; Bett et al., 2014]. Within populations, village chickens often display a vast array of phenotypes with difference among birds in plumage colour and pattern, size, comb shape, earlobe colour, skin colour, etc., even in a typical family flock that contains a few chickens (fig. 12). These phenotypes are often shared across populations over a large geographic distance, and while village chicken may be referred as distinct ecotypes or landraces, following the agro-ecological systems where they are kept, they are often not distinguishable based on phenotypes. Village chicken will often display morphological characteristics of fancy chicken breeds [Scrivener, 2006, 2009]. These phenotypes may therefore be of ancient origin, and they were likely selected from village chicken. Productivity of village chicken is lower compared to commercial breeds, but variations in quantitative traits (e.g. egg-laying characteristics, body weight and



Fig. 12. Example of plumage diversity in domestic chicken observed in a village household (Ethiopia).

growth rate) have been reported indicating their potential for breeding improvement [e.g. Dana et al., 2010]. This is well illustrated by the commercial breeds, which were initially developed from village chicken populations (e.g. the commercial white egg chicken lines developed from the Mediterranean Leghorn) or their crossbreeds (e.g. commercial broiler) [Crawford, 1990].

The main ancestor of domestic chicken, the red junglefowl Gallus gallus, shows uniformity in its plumage and morphology with however a strong sexual dimorphism [Delacour, 1977]. Indeed, only subtle morphological and plumage differentiation are distinguishing the different subspecies of the red junglefowl Gallus gallus ssp [Delacour, 1977; McGowan and Madge, 2010]. So the extensive plumage and morphological phenotypic diversity found in village chicken is not observed in the red junglefowl. It may be expected that the process of domestication and selection for behavioural traits may have 'relaxed' the natural selection constraints on the morphology and plumage of the birds with human selection for phenotypic novelties allowing survival of birds displaying non-wild phenotypes. Increasingly, the genetic control of this diversity is being unravelled. It includes nonsynonymous mutation in coding regions (e.g. lavender plumage) [Vaez et al., 2008], mutation with regulatory effect (silky-feather) [Feng et al., 2014], chromosomal inversion (e.g. rosecomb) [Imsland et al., 2012], retroviral insertion (oocyan egg phenotype) [Wang et al., 2013; Wragg et al., 2013], etc. Whether or not shared phenotypes observed across

Third Report on Chicken Genes and Chromosomes 2015 village chicken populations, including from different continents, do have a common genetic control ('founder phenotype') [Megens and Groenen, 2012] remains unknown, but it may be expected that the same loci are involved. However, in one case, unique so far, it has been shown that a domestic chicken phenotype (oocyan; blue/ green egg shelf colour) has evolved independently on the Asian and South American continents [Wang et al., 2013; Wragg et al., 2013]. Also for the yellow skin phenotype, commonly found in village chicken but also in commercial and fancy chicken, we know now that it originates from the genetic introgression of the grey junglefowl G. sonneratii [Eriksson et al., 2008]. This result is in agreement with earlier observations that reported that F1 birds might be obtained from crosses among all junglefowl species and domestic chickens [Crawford, 1990]. Also, F1 females between grey junglefowl males and red junglefowl females produce fertile offspring when backcrossed to either male grey or red junglefowl [Morejohn, 1968], and accordingly Nishibori et al. [2005] have shown the introgression of red junglefowl mtDNA into the grey junglefowl. Whether non-red junglefowl species introgression makes the genetic basis of domestic chicken phenotypes beside the yellow skin remains unknown. So far, there are no documented genetic evidences of introgression from the green junglefowl G. varius or the Sri Lanka junglefowl G. lafayettii into village chicken. However, worth mentioning is the common practice of breeding of F1 hybrids (called bekisar) between male green junglefowl and female domestic chicken on the Indonesian island of Java [Delacour, 1977], as well as some recent evidences of genetic introgression of domestic chicken into green junglefowl [Sawai et al., 2010].

Origin of the Genetic Diversity of Village Chicken

The publication of the chicken genome has opened the door to genome-wide analysis of DNA polymorphism in the species [International Chicken Genome Sequencing Consortium, 2004]. In a companion paper, ~2.8 million single nucleotide polymorphisms (SNPs) were identified comparing only 3 domestic chickens (a broiler, a layer and a Chinese Silkie) to a red junglefowl hen [International Chicken Polymorphism Map Consortium, 2004]. The mean nucleotide diversity reported was about 5 SNPs per kb between domestic chickens and red junglefowl as well as within chickens. It is 6–7 times larger to what has been observed, for example, in human or in dog and in contrast to the expectation that domestic animals are inbred compared to their wild ancestors [International Chicken Polymorphism Map Consortium, 2004]. Moreover, the segre-

Cytogenet Genome Res 2015;145:78–179 DOI: 10.1159/000430927

gation of these polymorphisms across breeds indicated that most of these were present before the divergence of modern breeds and might have originated before domestication [International Chicken Polymorphism Map Consortium, 2004]. These findings underline the importance of village chicken as a reservoir of diversity.

Today, it is possible to reconcile the high genetic diversity of village chicken with our current but still fragmentary knowledge on the history of the species. Mitochondrial DNA studies indicate that the main maternal ancestor of domestic chicken is the red junglefowl G. gallus. The today geographic range of the red junglefowl extends from the north of the Indian subcontinent to western China, reaching in the south the Indonesian islands of Sumatra and Java [Delacour, 1977; McGowan and Madge, 2010]. It comprises 5 subspecies (G. g. gallus, G. g. jabouillei, G. g. murghi, G. g. bankiva, G. g. spadiceus). All or several of these might have contributed to the genetic background and the diversity of the domestic chicken at the time of domestication assuming several domestication centers [Liu et al., 2006; Kanginakudru et al., 2008] or subsequently through introgression into the domestic gene pools. Even today, wild red junglefowls are regularly visiting villages and crossbreedings between domestic chicken and wild red junglefowl have been documented [Berthouly et al., 2009]. Any introgression from other Gallus species will also increase the diversity of the village chicken genetic pool.

Outside the geographic ranges of the wild ancestor(s), the genetic diversity of village chicken will depend on the demographic history of the population, including founding events, multiple introductions, crossbreeding, bottleneck, genetic drift, selection, and mutation. The largely uncontrolled breeding system of village chickens is making them less vulnerable to genetic erosion compared to fancy and commercial chicken breeds. Commercial and improved chicken breeds, which are being introduced worldwide in attempts to improve village chicken productivity, may now impact at short term the genetic diversity of village chicken.

Characterization of the Genetic Diversity of Village Chicken

Besides the phenotypic diversity, several studies have now reported the genetic diversity of village chicken. In particular, many studies have now addressed the genetic diversity of village chicken at country or population level [e.g. Silva et al., 2009], sometimes including commercial birds as references [e.g. Leroy et al., 2012]. Large-scale studies involving different geographic regions and/or breeds using autosomal markers are few [e.g. Muir et al., 2008a; Lyimo et al., 2014]. However, several studies using modern and/or ancient mtDNA sequence information have now examined diversity of populations across several countries [Liu et al., 2006; Storey et al., 2012; Miao et al., 2012; Thomson et al., 2014]. No studies have reported so far the genome-wide functional diversity of village chicken, but a few have assessed polymorphism at candidate genes [e.g. Downing et al., 2009a, b].

From a breeding and a conservation point of view, we are interested in 3 main objectives: (1) The assessment of local village chicken genetic diversity, as indigenous chicken populations represent a reservoir of local adaptation to environmental challenges (e.g. climate, diseases) and therefore an indispensable genetic resource for local chicken production; (2) the assessment of village chicken diversity in comparison to the commercial lines, with indigenous chicken populations representing a reservoir of genetic variation for further improvement of productivity; and (3) the mapping of village chicken genetic diversity to develop short- and long-term strategies for the conservation of chicken genetic resources and to assist in population choices for crossbreeding programs.

The most extensive study that assessed the genetic diversity of commercial lines in comparison to other chicken breeds is the one of Muir et al. [2008a]. They did a genome-wide assessment of chicken SNP genetic diversity using 2,551 autosomal markers. More than half of the birds were of commercial origin including broiler, white and brown egg-layers, while the remaining ones included experimental and standard breeds of chicken. Reconstructing allele frequencies from a hypothetical ancestral population, they showed that >50% of the genetic diversity in ancestral breeds is absent in commercial lines. The missing diversity is primarily associated with founding effect attributed to small numbers of breeds and populations used to develop the commercial lines, rather than selection performed by the breeding companies. This loss of diversity observed in commercial chickens is primarily due to loss of rare alleles. The adaptive implication of this loss of diversity was not directly investigated but it has been argued that such loss might be of particular importance for local adaption (e.g. disease resistances). More recently, the study of Lyimo et al. [2014] included 9 populations of commercial lines, 3 populations of red junglefowl as well as a large number of populations from the African, Asian and European continents. They used 29 autosomal microsatellites. The highest diversity (heterozygosity and high mean numbers of alleles) was observed

Schmid et al.

in the red junglefowl populations and village chicken populations from Africa and Asia. From the latter, Southeast Asia is the region with the highest domestic chicken genetic diversity. At the opposite, chicken populations from Northwest Europe show a genetic diversity comparable to commercial lines, with the notable exception of the white egg-layer chicken lines which have the lowest diversity of all populations.

The most exhaustive studies so far in terms of geographic coverage of village chicken diversity have examined mtDNA diversity, and in particular D-loop sequence polymorphisms, with some recent studies including also full mtDNA sequences [e.g. Miao et al., 2012]. These studies primarily aim to unravel the origin of domestic chicken and their dispersion across the world as a proxy to our understanding of human past migrations and trading contacts. Village chicken diversity is an important source of information assuming that their modern mtDNA still carries genetic signatures of past demographic events. To which extent this is true may be examined through ancient DNA studies. Although restricted to a maternally inherited and a small portion of the genome, with no recombination, these studies provide an important framework to assess the origin and the today geographic distribution of village chicken diversity.

Liu et al. [2006] were the first to perform a large-scale study on mtDNA diversity of village chicken. Through the analysis of more than 800 birds across Eurasia, they revealed 9 different clades (haplogroups) with different geographic distribution patterns and expansion signatures suggesting that they originated in different geographic regions. Their findings led them to propose multiple origins of domestic chicken involving South and Southeast Asia as opposed to the previous findings of Fumihito et al. [1994, 1996] which suggested that domestic chicken originated from a single domestication event in Thailand and/or adjacent regions. Subsequently, several studies confirmed the phylogeographic substructure of mtDNA in village chickens both within and outside of the geographic range of the wild ancestor [e.g. Kanginakudru et al., 2008; Muchadeyi et al., 2008; Silva et al., 2009; Mwacharo et al., 2011; Thomson et al., 2014]. These studies indicate that the diversity of village chicken mtDNA is distinct from one geographic region to another and that it has been shaped by human migration and trading networks. It is exemplified by the most likely Indonesian origin, through the Austronesian expansion of a specific mtDNA subhaplogroup in Madagascar, Southern and Eastern Africa [Razafindraibe et al., 2008; Muchadeyi et al., 2008; Mwacharo et al., 2011], as well as by the haplogroup diversity of village chicken from Sri Lanka, an ancient trading crossroad island on the Indian Ocean trading network [Silva et al., 2009]. These maritime dispersion events resulted in mtDNA diversity bottlenecks, which might have been compensated by subsequent admixture and gene flow between chicken populations originating from different dispersion events [e.g. Mwacharo et al., 2011, 2013]. Importantly, all these mitochondrial studies illustrate that village chicken diversity is distinct from one region to another with genetic drift being a major factor in the shaping of this diversity.

Village Chicken as a Reservoir of Genetic and Phenotypic Diversity for Genomics Studies

The high and ancient diversity as well as the breeding history of village chicken make them a treasure-trove for genome association studies of Mendelian and quantitative traits [Megens and Groenen, 2012; Wragg et al., 2012]. Linkage disequilibrium extends to short physical distances [Desta, 2015], and comparison among village chicken populations may offer the opportunity to finemap the regions of interest through partial overlaps of haplotype block structures. Currently, the chicken genomics community has access to 2 genome-wide screening tools, a genotyping assay with around 55,000 SNPs [Groenen et al., 2011] and a high-density one including 600K SNPs [Kranis et al., 2013], well above the recommended number of 100,000 markers for genome-wide association studies [Wragg et al., 2012]. Traits like earlobes, skin colour, egg pigmentation, and comb shapes have now been mapped using the 55,000 SNP chips [Wragg et al., 2012; Siwek et al., 2013]. A study investigating the genetic control of plumage colours and patterns using the high-density 600K SNP chips shows promising results [Desta, 2015]. In the future, it is expected that full-genome sequence data will become increasingly available to unravel in-depth the genome diversity of chickens. Also, beside mapping and identification of the genetic control of morphological and production traits, village chicken populations will be increasingly studied as a model to understand how adaptation to environmental challenges may have shaped the diversity of an avian genome.

Mendelian Traits

(Prepared by D. Wragg)

Mendel's 'Versuche über Pflanzenhybriden' [Mendel, 1866] detailed his experiments on inheritance in peas, the results of which provided evidence for his Principles of

Segregation and of Independent Assortment. In eukaryotes, both of these principles occur in meiosis during which homologous chromosomes exchange genetic information through recombination. However, the Principle of Independent Assortment does not always hold true for genes located proximal to one another, as these are often inherited together during meiosis, and so there exists a measure of genetic linkage known as recombination frequency. This in turn gives rise to the phenomenon of linkage disequilibrium (LD) - the non-random association of alleles at multiple loci along a chromosome. The extent of LD can be influenced by several factors, including for example genetic linkage, selection, and population structure, which combine to form combinations of alleles that occur more frequently than would be expected under a random distribution - in which case they would be considered in linkage equilibrium. Thus, closely related individuals, having witnessed few recombination events, exhibit a greater extent of LD than might be observed in distantly related individuals. When a characteristic has been selected, for instance a particular plumage variant, then the alleles proximal to the loci responsible for the variant will exhibit high LD relative to regions of the genome subject to little or no selection.

These concepts, together with the availability of highdensity single nucleotide polymorphism (SNP) arrays [Groenen et al., 2011; Kranis et al., 2013] and the declining costs of next-generation sequencing, have enabled many phenotypic characteristics of the chicken to be mapped, and subsequently to have their causal genetic mutation identified. As of September 2014, the Online Mendelian Inheritance in Animals (OMIA) database (http://omia.angis.org.au) lists a total of 127 Mendelian traits/disorders in the chicken; for 40 of which the key mutation has been identified. As might be expected, a little over one third of these (35%) relate to plumage variants. Presented here is a brief review of some of the phenotypes that have been characterized to date.

Plumage

The locus responsible for sex-linked barring (*B*) was mapped to the Z chromosome in an F2 population using microsatellites [Dorshorst and Ashwell, 2009], and subsequently refined by Hellström et al. [2010] leading to the identification of 2 SNPs in *CDK2NA* in nearly complete association with the phenotype. Although the 2 SNPs are located in putative E2F-1 and NF κ B transcription factor binding sites, attempts to study the functional significance of the 2 polymorphisms by electrophoretic mobility shift and luciferase reporter assays did not reveal any significant differences between any allele pair [Hellström et al., 2010]. Hellström et al. [2010] propose that the *CDK2NA* mutations might lead to premature cell death of melanocytes in the feather resulting in a lack of pigment, and that a new wave of melanocytes migrates to colonize the feather follicle and produce melanin, giving rise to the barring pattern, characteristic of the Plymouth Rock chicken.

In addition to variations in plumage pigmentation, some chicken breeds are characterized by the structure of their feathers, and the most striking of these phenotypes are the autosomal incompletely dominant frizzle (F) and recessive silkie (h) traits. Linkage analysis by Ng et al. [2012] mapped the F locus to an as yet unplaced linkage group (E22C19W28 E50C23) containing a number of keratin genes. Subsequent sequencing identified an 84-bp deletion in *KRT75* in complete segregation with the trait. The deletion, which spans a splice site, results in the loss of 23 amino acids from the K75 protein and consequently affects the development of the feather. The h locus was mapped by Dorshorst et al. [2010] to chromosome 3 using a mapping resource population, and subsequently refined by Feng et al. [2014] using the 60K SNP array to a 380-kb interval with an overlap of around 182 kb with the earlier mapping by Dorshorst et al. [2010] leading to the identification of a *cis*-regulatory mutation of PDSS2. Sequencing by Feng et al. [2014] identified an 18.9-kb haplotype fixed in chickens possessing *h* alleles, within which a single SNP was found to be in complete segregation with the trait, located 103 bp upstream of the initiator codon ATG of PDSS2. Subsequent expression analysis revealed the mutation to significantly decrease expression of PDSS2 during feather development.

Another plumage variant, the autosomal dominant naked neck (Na), affects the distribution of feathers. Na was mapped to a 200-kb region identicaI-by-descent (IBD) on chromosome 3 [Mou et al., 2011]. Tiling the region with overlapping PCRs revealed that one specific region could not be amplified in naked-neck individuals, suggesting a possible genomic rearrangement. This was subsequently confirmed by sequencing to be a large insertion of sequence from chromosome 1, resulting in a longrange cis-regulatory effect on BMP12. The inserted sequence, originating from between WNT11 and UVRAG, contains conserved noncoding elements, but no sequence that is predicted to be transcribed. The precise mechanism of the mutation remains to be determined, although the authors suggest that BMP12 upregulation might be due to possible WNT11 enhancers in the inserted sequence. This is supported by the nature of the phenotype

Schmid et al.

and the observation that *WNT11* expression was found to be significantly stronger on the neck than the body.

In addition to the absence of neck feathers, another mutation exists resulting in the near complete absence of scales and feathers, known as scaleless (*sc*) [Abbott and Asmundson, 1957]. Using the 60K SNP array, a significant association for *sc* was mapped to a 1.25-Mb region on chromosome 4 containing 11 genes [Wells et al., 2012]. Sequencing of these genes identified a SNP in exon 3 of *FGF20* leading to a premature stop codon and the predicted subsequent loss of receptor and HSPG binding sites. In situ hybridization demonstrated that *FGF20* was expressed in the developing feather placodes, confirming its role in early feather development, with qRT-PCR indicating expression to be specific to the epidermis.

Combs

Following the rediscovery of Mendel's results, the pea comb was one of the first traits to have its mode of inheritance documented [Bateson, 1902]. Prior to the sequencing of the chicken genome, linkage analysis had mapped the autosomal dominant pea comb (P) to chromosome 1 [Bitgood et al., 1980; Bartlett, 1996]. After further refining the localization of the locus through SNP genotyping, Wright et al. [2009] detected a copy number variation in intron 1 of the SOX5 gene. Bateson [1902] also described the dominant inheritance of the rose comb (R), and together with Punnet [Bateson and Punnet, 1908] presented the first case of an epistatic interaction between genes, whereby individuals possessing both P and R alleles exhibit a walnut comb. The R locus was mapped to chromosome 7 [Dorshorst et al., 2010; Wragg et al., 2012] and subsequently identified to be the result of a 7.4-Mb inversion [Imsland et al., 2012]. The inversion results in the re-localization of MNR2, leading to its ectopic expression during comb development. In addition, Imsland et al. [2012] postulate that the reduced fertility observed in homozygotes for R might be due to one of the breakpoints of the inversion disrupting CCDC108, resulting in a truncated transcript in the testis. Another common comb type is the autosomal incompletely dominant duplex (D), which has been mapped to chromosome 2 [Dorshorst et al., 2010; Wragg et al., 2012], although the causal mutation remains to be reported.

Skin/Shank Colour

Despite being recessive, the yellow skin phenotype is abundant throughout commercial chicken. The localization of the yellow skin locus (W) to chromosome 24 was published in the First Report on Chicken Genes and Chromosomes [Schmid et al., 2000]. Eriksson et al. [2008] used a back-cross pedigree to refine its mapping, revealing close linkage at the distal end of the chromosome. IBD mapping across a range of breeds revealed a single SNP in BCO2 in complete LD with the trait, and a minimum haplotype size of 23.8 kb. Subsequent sequencing and RT-PCR analysis suggest that tissue-specific regulatory mutation(s) alter BCO2 expression in the skin. However, the precise causal mutation remains to be determined. Also of note, arising from the same study was a phylogenetic analysis of the different junglefowl species using sequences derived from BCO2. The results of this analysis suggest that yellow skin most likely originates from the grey junglefowl, indicating a hybrid origin of the domestic chicken, although the ubiquity of the phenotype did not emerge until much later during the rapid worldwide spread of commercial chickens [Flink et al., 2014].

Black skin, also known as fibromelanosis and dermal hyperpigmentation, is a defining characteristic of the Silkie breed, amongst several others. Dorshorst et al. [2010] mapped the autosomal dominant locus (Fm) to a 2.8-Mb region of chromosome 20 using a back-cross mapping population, and subsequently refined the region to 483 kb in the same population using additional markers [Dorshorst et al., 2011]. IBD analysis identified a 75kb haplotype within this region which contained 5 SNPs observed to be heterozygous in all black skin chickens. This observation prompted an analysis of copy number variation from the Log R ratio data from the 60K SNP array. A group-wise analysis comparing *Fm* and wild-type individuals revealed 2 putative duplicated regions, each larger than 100 kb and separated by 417 kb on the reference genome. Further analyses by PCR confirmed the duplications, and revealed each duplicated region to be joined to the other in an inverted orientation. The result is an increased expression of EDN3, located within the first duplicated region, leading to dermal hyperpigmentation. The duplication and overexpression of EDN3 was independently discovered and reported by Shinomiya et al. [2012].

Green shanks are the result of an interaction between the sex-linked incompletely dominant inhibitor of dermal melanin (*Id*) and the *W* loci [Smyth, 1990], and are characteristic of the green-legged partridgelike fowl (GP) of Poland. The GP was subject to a study by Siwek et al. [2013] to assess its genetic history using mtDNA and the 60K SNP array. In particular, significant associations were found for both *Id* and *W*, indicating that the reseda green leg phenotype is a result of recessive alleles at these 2 loci. The results of the study also highlighted the pos-

sibility of multiple alleles for Id, as noted by Smyth [1990]. As the causal mutation underlying Id has not yet been identified, the mechanisms behind the interaction between this and the W locus, leading to green shanks, remain to be established.

Blue-Shelled Eggs

As with Fm, the molecular basis underlying the autosomal dominant blue eggshell (O) phenotype has been independently discovered and reported by 2 groups, namely Wang et al. [2013] and Wragg et al. [2012, 2013]. In summary, Wang et al. investigated the trait through classical linkage analysis in an F2 resource population of Dongxiang chickens, mapping the locus to a ~120-kb region on chromosome 1. Whereas Wragg et al. focussed on a diversity of chicken breeds and the Mapuche fowl of South America, from which the modern Araucana breed descends, applying an across-breed mapping strategy using the 60K SNP array to identify a 313-kb haplotype that was IBD in blue-shelled egg breeds. Both groups confirmed the tissue-specific overexpression of SLCO1B3 in blue-shelled hens as a result of an endogenous retrovirus (EAV-HP) insertion upstream of the promoter; together with independent integration events for the South American and Asian continents, supported by different integration sites and target-site duplications. SLCO1B3 is involved in bile acid transport, and its overexpression in the uterus of blue-shelled hens likely enhances the transportation of the bile salt biliverdin to the egg shell; biliverdin being the principal pigment in blue-shelled eggs.

Endogenous retroviruses have also been found to affect other phenotypic traits. For instance, an insertion of avian leukosis virus (ALV) in intron 4 of *TYR* results in the recessive white plumage of some breeds [Chang et al., 2006]. In addition, late-feathering, which is widely used to sex the chicks of some breeds, was originally linked to the promoter activity of a retroviral long-terminal repeat [Matsumine et al., 1991]; however, more recently it has been associated with a 176-kb tandem duplication resulting in the partial duplication of *PRLR* and *SPEF2* [Elferink et al., 2008]. In particular, experiments in mice have shown *PRLR* to influence the timing mechanism of the hair follicle [Craven et al., 2001], and it is therefore a strong candidate for late-feathering.

Foot Development

Several mutations affecting limb development have been documented, from which two that affect foot development have been characterized: oligozeugodactyly (*ozd*) and preaxial polydactyly (*Po*). Birds possessing an allele for Po have an additional digit on one or both feet and/or wings; however, the expression is variable and temperature-sensitive [Robb and Delany, 2012a]. A highly conserved long-range cis-regulatory SHH enhancer, known as the zone of polarizing activity regulatory sequence (ZRS), is present in intron 5 of *LMBR1*. In the case of *Po* in the Silkie breed, an association with a SNP within the ZRS has been reported in several studies [Dorshorst et al., 2010; Dunn et al., 2011; Maas et al., 2011] and found to result in an allelic imbalance of SHH [Dunn et al., 2011]. In contrast to Po, ozd results in the absence of the ulna and fibula and all digits with the exception of one in the foot. However, in common with Po, ozd is the result of a mutation in the ZRS, in this case a 1,654-bp deletion causing the loss of all but the first 135 bp of the ZRS [Maas et al., 2011]. These results demonstrate the utility of the chicken as a valuable model system for the study of both limb development and gene regulation.

Model for Disease

A review on the chicken as an animal model for eye disease [Hocking and Guggenheim, 2013] outlines a number of recessive mutations associated with blindness. Those for which the causal variants have been identified include chicken albino (*ca*), arising from a 6-bp deletion in *TYR* [Tobita-Teramoto et al., 2000]; retinal degeneration (*rd*), caused by an InDel in *GCI* [Semple-Rowland et al., 1998]; retinal dysplasia and degeneration (*rdd*), resulting from a premature stop codon in *MPDZ* [Ali et al., 2011]; and retina globe enlarged (*rge*), which is due to a 3-bp deletion in *GNB3* [Tummala et al., 2006]. The extensive genetic resources available and the large eye size make the chicken an important ophthalmologic model.

Photosensitive reflex epilepsy has been characterized as an autosomal recessive trait in the Fepi chicken strain [Douaud et al., 2011]. In this instance, a mutation in the second intron of *SV2A* results in abnormal splicing, significantly reducing its expression. Upon identifying the mutation, the chickens were treated with an anti-epileptic drug (levetiracetam) which was found to significantly reduce the number and duration of seizures, indicating its anticonvulsant effect in the chicken. The mechanism of action of *SV2A* and its interaction with levetiracetam remain to be fully elucidated, and so the Fepi chicken provides an important model for future research.

The degenerative muscle disorder muscular dystrophy has been identified as an autosomal co-dominant trait in the chicken, for which a missense mutation in *WWP1* has been implicated [Matsumoto et al., 2008]. Comparative genomics indicates the missense mutation to be present only in dystrophic chickens, and that the region is highly conserved across the tetrapods analyzed (7 mammals, 4 reptiles, 1 amphibian, 2 birds). The mutation itself causes an amino acid replacement from arginine with a basic side chain, to glutamine with an uncharged polar side chain, and may therefore affect the function of the *WWP1* protein; although no significant difference in the expression levels of *WWP1* was observed between mutant and wildtype chickens. Further investigation is required to clarify the mechanism by which *WWP1* triggers muscular dystrophy in chicken, the results of which could provide new insights for understanding human muscular dystrophies.

Reproductive Biology

In addition to its utility as a model for disease, and for characterizing the genetics of phenotypic diversity, the chicken is also a valuable resource for understanding reproductive biology. A review of research on the sex-linked recessive restricted ovulator (ro) phenotype [Elkin et al., 2012] indicates it to result from a naturally occurring point mutation in VLDLR located on the Z chromosome. The ovaries of R/O chickens lack follicular hierarchy and contain many small preovulatory follicles of various colours, shapes and sizes; hens also exhibit hyperestrogenemia, hypoprogesteronemia, elevated circulating gonadotropins, and upregulated pituitary progesterone receptor mRNA and isoforms. Another model of compromised female fertility, the Watanabe heritable hyperlipidemic (WHHL) rabbit [Shiomi and Ito, 2009], is widely used for studies of human autosomal dominant familial hypercholesterolemia. WHHL rabbits and R/O hens each possess defective LDL receptor supergene family members; however, the etiology of reproductive dysfunction in the 2 models is likely to be different. Nonetheless, further studies of the R/O model to improve an understanding of the molecular physiology and biochemistry concerned, may lead to new analogies between avian and mammalian systems and help to address important questions in reproductive biology.

Future Opportunities

As indicated from the number of Mendelian traits in OMIA for which the causal mutation has not been identified, many opportunities remain for further investigation. For instance ear-tufts, muffs and beard, and vulture hocks to name but a few. The Araucana and Mapuche fowl, which have already provided an important model for studying blue-shelled eggs, could also present a valuable resource for investigating ear-tufts, muffs and beard. Indeed, the Araucana has recently been the subject of study to identify candidate genes for rumpless and eartufts using the 60K SNP array, mapping them to chromosomes 2 and 15, respectively [Noorai et al., 2012]. The chicken is also an important model for studies concerning reproductive biology and disease, as illustrated in the small number of examples presented here, and is a potential model for numerous human diseases including for example hypothyroidism [Cole, 1966] and systemic sclerosis [Beyer et al., 2010] amongst many others. The recent availability of a 600K SNP array and the declining costs of whole-genome re-sequencing present opportunities to economically map traits at a finer resolution than previously possible. Furthermore, the widespread use of these technologies, combined with accurate phenotypic records, facilitates the incorporation of data from different studies with little difficulty.

Treasure the Exceptions: Utilizing Chicken Mutant Lines and Advanced Genetic Technologies to Uncover Genes Involved in Developmental Processes (Prepared by E.A. O'Hare and M.E. Delany)

Much of the knowledge underpinning cellular processes obtained by biologists to date has been contributed by the study of mutations, either natural variants or induced. Such discoveries shed light on many developmental mechanisms and pathways and in turn provide immeasurable benefits to the world through impacts in health and medicine.

The chicken was one of the first animals used for the analysis of human development (aka animal model) when it was studied by Aristotle (ca. 400 B.C.) to understand chicken embryo formation. Considerable use has been made of chicken mutants/genetic variants in developmental biology. Similarly, the chicken also served as an early genetics model during the seminal studies of Bateson (ca. 1900), wherein by studying comb inheritance, he proved that Mendel's laws operated in animals.

The beauty of the classic chicken model is in its in ovo embryogenesis allowing for easy access and analysis of all stages of development in addition to a relatively short generation interval and large number of progeny. Given this, it was possible to identify a series of developmental abnormalities involving morphogenesis; this in turn led to the discovery and establishment of hundreds of unique genetic resources across the world by both university and government institutions during the 20th century [Somes, 1990a, b; Pisenti et al., 1999; Delany, 2004; Delany and Gessaro, 2008]. To date, many chicken genetic lines have been well-characterized for phenotype and mode of inheritance and were developed into inbred and/or congenic lines, thereby presenting mapping opportunities, candidate gene analysis and the determination of causative gene(s) and involved pathways [Pisenti et al., 1999; Delany, 2004; Robb et al., 2011].

The advent of the chicken genome sequence [International Chicken Genome Sequencing Consortium, 2004], high-throughput technologies, and genetic manipulation techniques provide an opportunity to quickly study priority candidate elements in order to better understand chicken, avian and vertebrate biology, the developmental mechanisms and pathways involved, and overall contribution to phenotype and/or disease onset [Muir et al., 2008a; Robb et al., 2011; Robb and Delany, 2012a, b; Stern, 2004]. Moreover, since many of the malformations observed in the chicken show similarity with human conditions, identification of the causative variant(s) is relevant to human health and development. Thus, the chicken is a valuable model for the scientific community and as such has been recognized as a model organism for biomedical research by the National Institutes of Health in 2009 (http://www.nih.gov/science/models/gallus/). In this section, we discuss the use of advanced technologies for the identification of those genetic variants perturbing normal development through the use of unique chicken genetic resources.

Mechanisms of 'Gene Hunting'

The resounding success of the chicken as the premier non-mammalian vertebrate model can be attributed to the tools, strategies and genetic lines created by poultry researchers dating back over a century. The release of the chicken genome sequence and subsequent genome browser interfaces marked a historical point in this research timeline by allowing scientists to quickly and easily find a gene of interest and identify its corresponding nucleotide or protein sequence [International Chicken Genome Sequencing Consortium, 2004]. The pairing of genomic tools and sequencing strategies with unique, naturally-occurring developmental mutations in the chicken allows one to hone in on the causative gene shedding light onto the mechanisms regulating development. To date a variety of methods have been utilized to identify the causative variant, gene function and biological pathways and processes of a number of developmental mutations in the chicken. Below we describe how different techniques and technologies were utilized to uncover genes involved in development.

Mapping Techniques

The chicken chromosome map has evolved greatly over the past century to become a more fine-tuned genetic map with 97-98% of the total genomic sequence identified and assigned to specific chromosomes. However, even with the current status of the chicken genome, finding the causative variant for a particular phenotype would be nearly impossible without the knowledge of basic genetics and inheritance. As early as the beginning of the 20th century, chicken researchers showed that barring is sex-linked [Spillman, 1908]. Since that time numerous developmental traits have been pinpointed to a particular chromosome type (i.e. autosomal vs. sex) and in many cases, linkage analyses and complementation studies have helped to further map phenotypes to unique chromosomes [Bitgood and Somes, 1990] aiding in the fine-mapping of these unique developmental mutations. In fact, the well-known *talpid*³ polydactylous mutation was localized to chromosome 5 using linkage analyses, and additional sequencing revealed a frameshift mutation in KIAA0586 (aka TALPID3) as the causative mutation [Davey et al., 2006].

Analysis of chromosome structure is also utilized to pinpoint the region harboring causative developmental mutations. Langhorst and Fechheimer [1985] identified an autosomal recessive skeletal mutation, designated shankless (shl), which is characterized by malformed metacarpals, the lack of tarsometatarsal shanks, increased tibia length, extra bones in the digits, and fusion of the phalanges. These skeletal malformations were the result of an X-ray-induced pericentric inversion in chromosome 2 visualized by karyotype analyses [Langhorst and Fechheimer, 1985]. More recently, the causative element responsible for the classic monogenic trait rose-comb was determined to be the result of a chromosomal inversion. Imsland et al. [2012] elegantly utilized FISH and labeled BACs to show the presence and location of the chromosomal rearrangement, while Robb and Delany [2012a] employed the same technique to show that an inversion event was not responsible for the polydactyly phenotype. Thus, although basic in technique, classical methodologies (cytogenetics) combined with genomics (e.g. BACs) provide valuable opportunity in understanding the basis of genetic variants.

Genotyping Arrays

A single nucleotide polymorphism (SNP) array is an extremely valuable tool for studying variation between whole genomes. Exploiting this technology can shed light on linkage associations and fine-mapping, hetero-

zygosity, signatures of selection, and can even pinpoint causative variants for complex and simple genetic traits. Over the past decade, the poultry research community has successfully employed this technology to hone in on the gene(s) causing developmental mutations in the chicken. The availability of the 3K chicken SNP array in 2008 [Muir et al., 2008a] paired with specialized inbred lines allowed researchers to, for the first time, associate a specific chromosomal region with several developmental mutations. Dorshorst et al. [2010] used the 3K array to map 8 different traits in the Silkie chicken including the identification of a casual SNP present within a highly conserved cis-regulatory region of Sonic hedgehog for the polydactyly phenotype. Furthermore, despite the small number of SNPs, the 3K chicken SNP array successfully localized 7 of the 9 University of California, Davis-held developmental mutations to a unique chromosome (see suppl. table 1 in Robb et al. [2011]). Several years later, the utilization of the 60K SNP chip [Groenen et al., 2011] and an updated genome analysis allowed for further fine-mapping of those mutations with subsequent assignment of priority candidate genes [Robb et al., 2011] and the gene of interest for one, Polydactyly [Robb and Delany, 2012a]. The 60K SNP array was also useful in fine-mapping the region of the scaleless mutation to chromosome 4. After exome sequencing and subsequent analyses of the 11 genes in the region, a missense mutation in the FGF20 gene was identified as causative [Wells et al., 2012].

Copy number variants (CNVs) are small deletions and duplications of genomic DNA that impact gene expression usually identified using array-CGH. Jia et al. [2013] successfully employed the 60K SNP chip to identify new CNVs. As CNVs are commonly strongly associated with developmental malformations, utilization of SNP arrays in such a manner opens the door to investigations of associations of CNVs causing developmental defects. In 2012, a new commercially available high-density 600K SNP array was released [Kranis et al., 2013] and is being used by poultry researchers worldwide. This new chip will prove to be useful for identifying trait-associated loci, fine-mapping genomic regions, and detecting CNVs amongst all the other useful applications. With currently 3.5 million SNPs identified in the chicken to date, the chicken arrays will no doubt expand to rival those arrays used in human genome-wide association studies, extending their usefulness and allowing for a finer mapping of causative gene loci.

Sequencing

Massively parallel sequencing systems serve as a means to rapidly generate whole or targeted-region genomic sequence, gene expression profiles (e.g. RNA-seq), and protein-DNA/RNA interactions (e.g. ChIP-seq). Each application has been uniquely developed to fulfill different biological purposes. The use of these advanced technologies has revolutionized our understanding of genetics, genomics, and biology as a whole and has helped to accelerate the fields of health and disease in animals, plants and human.

Exome sequencing seeks to identify coding variants in specific, targeted genes of interest or of all known genes in the genome [Mamanova et al., 2010]. Although cheaper than whole-genome sequencing (WGS), targeted sequencing of only coding regions or lists of genes could miss genetic variants in intergenic and intronic regions or within an unknown gene. Targeted-exome sequencing has, however, been successful in the chicken in identifying the causative element for a number of developmental mutations. As previously noted, Wells et al. [2012] identified the causative element of the scaleless mutation after sequencing 11 genes on chromosome 4. Although variant identification through WGS is now becoming more costeffective, zooming in on the region of interest, when possible, remains an attractive and cost-saving option. Robb and Delany utilized a targeted sequence capture enrichment array to uncover the causative element for 3 unique inherited mutations (coloboma, diplopodia-1, wingless-2) disrupting limb, craniofacial and/or organ development [Robb, 2012; Robb and Delany, 2012b; Robb et al., 2013].

WGS is the most all-encompassing method for identifying elements causing developmental defects in any organism. This method can detect genetic variation in a genome including SNPs, large and small insertions and deletions, CNVs, and chromosome rearrangements [International Chicken Genome Sequencing Consortium, 2004; Yan et al., 2014]. Once the data is sorted and possible causative elements are prioritized, various techniques (e.g. qRT-PCR, in situ hybridization, microarrays, RNA-seq, siRNA, protein analysis, ChIP-seq, relevant functional assays) can be utilized to confirm or eliminate variants. Chang et al. [2014] successfully paired both the 60K chicken SNP array with WGS to identify C2CD3 as the causative gene for the long-studied *talpid*² mutation and provided additional evidence as to its mechanistic role using RNA-seq. Similarly, WGS and analyses have been employed for a number of the developmental mutations maintained at the University of California, Davis including diplopodia-3, diplopodia-4, eudiplopodia, limbless, and stumpy [Delany, unpubl. data].

Future Directions for Discovering the Underlying Factors and Functions of Developmental Mutations

The many naturally-occurring developmental mutations in the chicken offer opportunities for the study of unique aspects of developmental biology. Although many of these mutations were spontaneously derived (naturally-occurring variants), mutations can also be induced by mutagens (e.g. chemical, radiation) or targeted (e.g. TALEN, CRISPR). Gene function can also be assessed by repression of gene transcript (e.g. morpholino technology). Technology-driven mutagenesis provides a unique opportunity to conduct reverse genetics by assessment of specific gene function and will help revolutionize the field of functional genomics by allowing for the functional analysis of specific genes in vivo.

Transient Gene Perturbation

Morpholinos (MOs), antisense oligonucleotides used to downregulate a target gene, are a quick and easy way to assess the function of a gene, in a specific tissue, during embryonic development (GeneTools, LLC; http://www. gene-tools.com). MOs can be uniquely designed to inhibit translation (by blocking the translation initiation complex) or by modifying a normal splicing event resulting in an abnormal protein or splice-modification-triggered nonsense-mediated decay. This technology has been used in the chicken to show gene function in ovo for over 10 years [Kos et al., 2001]. Briefly, chick embryos are incubated to the developmental stage of interest, the egg is windowed on its side, and India ink is utilized to visualize the embryo. A micropipette containing the fluorescein-labeled MO (at a pre-determined concentration) is microinjected into the tissue of interest followed immediately by electroporation [Kos et al., 2003]. Uptake of the MO can be confirmed visually by fluorescent microscopy and effect of gene function can be assessed by morphological analysis [Kos et al., 2001, 2003]. This method of gene assessment can be particularly useful in eliminating possible causative genes and determining 'the gene' in those regions fine-mapped by other advanced technologies (e.g. SNP arrays, capture arrays, WGS). Shepard et al. [2008] elegantly showed the function of the gene versican (VCAN) in limb formation by using gene-specific MOs. This research helped to settle the gene's long-controversial role in embryogenesis. Assessment of the developing embryo indicated that VCAN is involved in skeletogenesis as microinjection of the VCAN MO into the precartilage limb core and distal limb mesenchyme resulted in abnormal development [Shepard et al., 2008]. To date, MOs have been used widely in targeted loss-of-function

experiments, and in many cases, the use of MOs has allowed for researchers to circumvent early lethality issues and identify function of genes in vivo (in ovo). Studies in the chick utilizing MOs will allow researchers to assess the effects of temporal loss-of-function thereby contributing to our knowledge of gene function in development. Assessment of the resulting morphology will elucidate the function and resulting phenotype of a gene of interest allowing researchers to discover new gene-interaction pathways, developmental processes, etc. Moreover, coinjection of a MO and the corresponding mutant transcript [containing variant(s) of interest] will shed light onto the functional role of unique variants in development allowing researchers to gain a better understanding of single variants and gene function.

Genetically Modified Animals

Targeted genome editing of any vertebrate system is an extensive process which involves the introduction or deletion of nucleotides in genomic DNA. Gene targeting/transgenic technology is a valuable approach to aid in the investigation of specific gene function in vivo. Transgenic and knockout animals can be generated using a number of different technologies including zinc finger nucleases (ZF), transactivator-like effector nuclease (TALEN), or RNA-guided mutagenesis using the CRISPR/Cas (clustered regularly interspaced short palindromic repeats-Cas) system; however, each strategy has its complications and limitations [Sanjana et al., 2012; Hsu et al., 2014]. Unfortunately, the difficulty in obtaining 1-celled embryos for injection and the lack of germ-line-competent embryonic stem cells has made conventional gene-targeting techniques established in other model systems difficult to employ in the chicken. However, these technical difficulties can be overcome through the use of primordial germ cells (PGCs), the precursors of eggs and sperm. Genetically modified chickens have been successfully developed by poultry researchers for over a decade. This has been accomplished through the transferring of specific genes to cultured PGCs with subsequent transplantation or microinjection to the developing embryo. Germline chimeras can then be used to produce knockout mutant offspring for future studies [van de Lavoir et al., 2006; Park and Han, 2012; Park et al., 2014a].

Genome editing, through the introduction of doublestranded breaks, is an effective way of introducing mutations into the genome. There are several new genomeediting tools available for use in the chicken. The first method is TALEN utilization, which employs an artifi-

cial restriction enzyme to generate DNA double-stranded breaks in vivo ultimately leading to the introduction of an error or new DNA sequence into the gene of interest. This enzyme was generated by fusing a DNA sequence recognition domain called transcription activator-like effector (TALE) with the catalytic DNA cleavage domain of the FokI nuclease (fused complex: TALEN) [Christian et al., 2010]. Briefly, the TALE domain sequence, engineered to identify a specific DNA sequence, is inserted into a plasmid and then introduced into a target cell (e.g. PGC), wherein it is translated to produce a functional TALEN enzyme. The enzyme then enters the nucleus and, upon recognition, binds to and cleaves the target sequence. TALEN-mediated gene targeting has been successfully utilized in the chicken to mutate a number of genes including those on microchromosomes, macrochromosomes and sex chromosomes [Park et al., 2014b].

The CRISPR/Cas is a fast and cost-effective system for gene modification. This new genome-editing system exploits the immune defense mechanism used by bacteria for destroying foreign genetic material [Barrangou et al., 2007]. The highly specific, genetically engineered RNAguided CRISPR/Cas nuclease system utilizes Watson-Crick base pairing to target genomic DNA. Upon identification of the complementary sequence, the Cas nuclease mediates a double-stranded break at a targeted genomic locus initiating subsequent error-prone cellular DNA damage repair pathways which commonly result in a gene knockout caused by frameshift mutations, premature stop codons, or even large deletions [Ran et al., 2013]. Although CRISPR/Cas system usage has not been reported in chicken to date, it is speculated to be more successful in generating genetically modified birds through injection into PGCs with subsequent transplantation.

Genome modification through use of either TALEN or CRISPR technologies will help shed light on the mechanisms behind developmental anomalies. Moreover, the development of germline mutant embryos will mimic the complete development and growth (i.e. from fertilization to hatch) of those naturally-occurring developmental mutations in the chicken as compared to assessment of gene function using MOs. Any of these tools, however, will serve as a powerful method for gene function identification in the chicken.

Conclusions

The chicken has long been a treasured resource for the scientific community contributing to advances in many fields, in particular, developmental biology. This valuable model is used to gain insight into the processes and mechanisms regulating normal development as well as shed light onto the etiology of both human and animal defects and syndromes and provide insight into possible therapeutic targets. Moreover, the advancement of gene-mapping, sequencing and mutagenesis technologies will further our knowledge of gene function in vivo. The scientific benefits of the classic chicken embryo vertebrate model system also include an expansive literature on developmental processes created by a large community of developmental, genetic and biomedical scientists. These benefits, coupled with the occurrence of inherited mutations affecting the body plan and organ systems [Pisenti et al., 1999; Delany, 2004; Robb et al., 2011, 2013; Robb and Delany, 2012a], now provide the opportunity to enhance our knowledge of the genes and sequences involved in vertebrate development. William Bateson [1908] coined the phrase 'Treasure your exceptions', emphasizing that naturally occurring variations, i.e. 'exceptions' to normal phenotypes, provide insight into the fundamental processes regulating growth and development. We now have the advantage of advanced tools to employ with either naturally-occurring or technology-induced variations to better understand processes regulating vertebrate development.

Genomic Landscape of the Chicken DT40 Cell Line

(Prepared by A. Motegi and M. Takata)

DT40 is a chicken B cell lymphoma cell line that is widely used as an excellent genetic model system with high gene-targeting efficiencies [Buerstedde and Takeda, 2006]. Along with its relatively stable karyotype and cytogenetic characters, DT40 cells provide a unique opportunity to study various cellular processes in vertebrate cells. In particular, DT40 cells contributed to our understanding of the molecular mechanisms of genome maintenance and DNA damage responses by allowing extensive genetic analyses [Motegi and Takata, 2014]. In this section, we summarize the latest state of knowledge of DT40 karyotype, genome, transcriptome, and possible molecular steps of DT40 cell development. We then briefly review the major insights into molecular mechanisms of preserving genomic integrity obtained through the reverse genetic analyses of DT40 cells.

DT40 Karyotype and Genome

DT40 cells were derived from an avian leukovirus (ALV)-transformed lymphoma experimentally devel-

oped in a domestic bred female chicken *Gallus gallus domestica* [Baba et al., 1985]. By the time of the Second Report on Chicken Genes and Chromosomes [Schmid et al., 2005], the release of reference genome sequences of red junglefowl *G. gallus* had much improved our understanding of the chicken genome, while strain- and cell lineagespecific sequence diversities in DT40 cells remained to be determined. During in this period, several genome-wide analyses of DT40 cells, including recent whole-genome shotgun sequencing and single nucleotide variation (SNV) array hybridization [Molnár et al., 2014], had been reported. By combining observations from these genomewide analyses and from conventional cytogenetic studies, we have extracted several important cytogenetic and genomic features of DT40 cells as listed below.

First, by evaluating the read depth of whole-genome sequencing, the original DT40 cells (from the Pirbright Institute, UK) were shown to have an almost normal chicken karyotype (38 pairs of autosomes and the ZW sex chromosomes [Masabanda et al., 2004]), only except for the trisomic macrochromosome 2 and possibly tetrasomic microchromosome 24 [Molnár et al., 2014]. We consistently observe the same karyotype of macrochromosomes and encounter aneuploidies at most less than several percent in our routine inspections into the mitotic spreads [Sonoda et al., 1998], albeit somewhat higher cytogenetic mosaicism of macrochrosomes was also reported in different stocks/conditions [Chang and Delany, 2004]. Tetrasomy of microchromosome 24 agrees with the existence of 4 alleles of the Cbl locus [Yasuda et al., 2000], which is now allocated on chromosome 24. We also note that wild type 'clone 18' from Sale Laboratory (MRC, UK) was found to have additional trisomies of microchromosomes 14 and 20 [Molnár et al., 2014], while the 'clone 18' stock in Kyoto was not trisomic in chromosome 14 [Yamamoto et al., 2011], suggesting that some level of microchromosomal copy number variation does exist among so-called 'clone 18' stocks.

Second, frequencies and spectrums (transitions and transversions) of all as well as unique SNVs in DT40 cells were very similar to those in normal blood cells from 2 domestic chicken breeds, L2 and Silkie [Fan et al., 2013] (different from Hyline SC breed, from which DT40 cells were derived), in comparison with the *G. gallus* reference genome [Molnár et al., 2014]. These observations substantiate the long-held notion that the genome of DT40 cells is fairly stable at a nucleotide level and that no nucleotide instability mechanisms have been operating in this cell line. In contrast, the numbers of insertions/deletions

(InDels) unique to DT40 cells were modestly higher than those in L2- and Silkie-derived cells (156.5% compared with the average from L2 and Silkie), while those in common among the 3 strains were indifferent [Molnár et al., 2014]. Closer inspection revealed that deletions of unit(s) in the 6~16-bp range at repeat sequences, but not insertions in all ranges, were evidently higher, although no obvious mutations or defects were identified in the mismatch repair pathway [Campo et al., 2013; Molnár et al., 2014], mostly responsible for deletion of this range. Interestingly, the ratio of homozygous versus heterozygous In-Dels unique to DT40 cells was somewhat higher, conceivably reflecting the enhanced level of loss of heterozygosity (LOH) in the 0.1~0.2-Mb range by homologous recombination (HR).

Third, integration sites of ALV were identified at 4 different loci [Molnár et al., 2014], including the previously known c-myc locus [Baba et al., 1985]. No apparent contribution of ALV integration at 3 new loci on B-cell transformation, if any, can be dictated from the currently available information. There were no gross chromosomal rearrangements involving the c-myc locus, which is frequently translocated with immunoglobulin loci in human B-cell lymphomas [Janz, 2006]. Since the c-myc locus is located within the region of extensive LOH, and all reads covering the integration site of *c-myc* were identical, ALV seems to have been integrated at one of the c-myc alleles first, and then LOH and whole copy number gain events occurred. Consistent with the previous cDNA microarray and array-comparative genomic hybridization (array-CGH) analyses [Neiman et al., 2001, 2006], the c-myc gene itself does not involve gene amplification, but integration of ALV at the first intron of c-myc juxtaposes the strong promoter in the long terminal repeat of ALV immediate upstream of exon 2 (the first coding exon) and drives overexpression of the c-myc mRNA (estimated 5~18-fold).

Lastly, the telomeric status is one of the critical features of cell lines with stable genomic transmission and infinite proliferation [Murnane, 2012]. DT40 cells exhibit telomerase activity comparable to that seen in chicken embryos, adult germline tissues and other transformed cell lines [Venkatesan and Price, 1998; Swanberg and Delany, 2003]. The average length of telomeres in DT40 cells was estimated to be 3~20 kb [Wei et al., 2002; Swanberg and Delany, 2003; Cooley et al., 2009; O'Hare and Delany, 2009], although some extent of range variations exists in reports at least partly due to the clone-to-clone and end-to-end differences and methodological biases [Aubert et al., 2012]. This range of telomeric repeat length is similar to that observed in normal chicken tissues and chicken embryonic fibroblasts [Venkatesan and Price, 1998], suggesting that telomerase expression in DT40 cells might be only reactive to maintain telomeres above a critical threshold. Cooley et al. [2009] estimated the lengths of telomeres in macrochromosomes and microchromosomes to be 17–43 and 70–1,000 kb in size, respectively, by using quantitative fluorescent in situ hybridization. Further studies would be required addressing whether the estimated ~3-fold increase in telomeric signals of microchromosomes reflects the actual difference in physical lengths or some biased factor(s) such as interstitial telomeric repeats, which were reported to be enriched in many avian microchromosomes [Nanda et al., 2002].

DT40 Transcriptome

The gene expression profile of DT40 cells has been studied by using custom-made microarrays [Neiman et al., 2001, 2006; Koskela et al., 2003]. One study focused on a limited number of bursa-specific ESTs and demonstrated that DT40 cells are more similar to normal bursal cells than B cells at other developmental stages [Koskela et al., 2003]. This observation matches the idea that DT40 cells are likely blocked in B cell differentiation at a stage of bursal cells, both of which already express rearranged cell surface IgM and retain stem cell function. Another study compared DT40 cells with experimentally induced chicken B-cell lymphomas and showed that the transcriptional signature of DT40 cells is more similar to those in bursal preneoplastic transformed follicles than to those in late stage metastatic tumors [Neiman et al., 2001]. In this study, the numbers of upregulated genes in myc-induced lymophomas, including DT40 cells, were positively correlated with the expression levels of myc, and those changes were diminished as myc expression was reduced in metastatic tumors. These observations are consistent with the notion that extensive myc-driven transcriptional changes in transformed follicles likely promote additional mutagenic events that fix transformed phenotypes [Neiman et al., 2001]. DT40 cells persistently express high levels of myc comparable to those in ALV-transformed lymphomas and another lymphoma-derived cell line, RP9 (5-18-fold compared with normal bursa cells), suggesting that the DT40 genome could be effectively protected from the myc-driven genomic stress [Hayward et al., 1981; Baba et al., 1985; Neiman et al., 2001, 2006]. How aberrant expression of myc is tolerated in DT40 cells, however, remains to be solved. Since the above microarrays contained limited numbers of genes with least

annotations, many genes might have escaped from attention. Further studies by using more standardized microarrays such as 33K Affymetrix GenChip and 44K Agilent microarray [Gheyas and Burt, 2013] could allow finer delineations of transcriptional signatures of B-cell lymphomas and DT40 cells.

MicroRNAs (miRNAs) constitute another layer of transcriptional regulations during B-cell development and differentiation [Fernando et al., 2012; de Yébenes et al., 2013; Baumjohann and Ansel, 2014]. miRNA expression profiles of naïve embryonic B cells, CD40L-stimulated B cells and DT40 cells have been studied by deep sequencing [Yao et al., 2013]. Around 20 miRNAs were identified to be upregulated (miR-18a, miR-18b, miR-20b, miR-21, miR-101, miR-103, miR-106, miR-148a, miR-221, and miR-222) or downregulated (miR-15b, miR-16, miR-26a, miR29a, miR-30a-5p, miR-30b, miR-30c, miR-30d, miR-30e, and miR-147) in both CD40Lstimulated and DT40 cells. Although most of miRNAs in these groups have not been well characterized in B-cell lineage, those are likely involved in cell proliferation and plasma cell differentiation, respectively. The overall expression profile of DT40 cells, however, was quite distinct from that of naïve and CD40L-stimulated cells. In particular, the most striking difference was observed in miR-155 (bic), which has been shown to facilitate ALV-induced lymphomagenesis [Tam et al., 2002]. Although the miR-155 locus was originally identified as a frequent target of ALV integration [Clurman and Hayward, 1989], no integration was found in DT40 cells [Molnár et al., 2014]. While miR-155 was the most upregulated miRNA in CD40L-activated B cells, it was rather the most downregulated one in DT40 cells [Neiman et al., 2006; Yao et al., 2013]. Intriguingly, miR-155 has been shown to repress expression of activation-induced deaminase [Dorsett et al., 2008; Teng et al., 2008], a master enzyme controlling immunoglobulin gene diversifications by somatic hypermutation and gene conversion [Arakawa and Buerstedde, 2009]. Therefore, it could be interesting to address the genomic status of miR-155 and to what extent the absence of miR-155 could influence the expression level and function of activation-induced deaminase in DT40 cells.

Insights into the Potential Molecular Steps of DT40 Cell Development

In addition to the *myc*-induced modulation of the transcriptome, DT40 cells are supposed to have developed by additional steps, including inactivation of the pro-apoptotic pathway(s) and activation of the pro-sur-

vival pathway(s). Repression of p53 is one of the earliest observations [Takao et al., 1999] that fits to the former idea and was regarded to be a good explanation for the weak G1 DNA-damage checkpoint and the relatively normal G2/M checkpoint in DT40 cells [Rainey et al., 2006]. We also did not detect p53 in wild-type DT40 cells and were not able to obtain stable clones expressing *p53* transgene [M.Takata, unpubl. data]. However, another line of study showed that the topoisomerase II inhibitor adriamycin did induce p53 expression and that p53 appeared to be functional at least in suppressing the expression of heat shock proteins [Tanikawa et al., 2000]. Therefore, the functionality of p53 in DT40 cells may need further evaluations.

The best clue that fits to the latter idea would be a mutation found in PIK3R1, which encodes the regulatory subunit a of the phosphatidylinositol 3-kinase (PI3-kinase) [Molnár et al., 2014]. A deletion of 6 bp in exon 11 of the only allele of PIK3R1 on the Z chromosome resulted in loss of 2 amino acids, E451 and Y452, located at the well-conserved interSH2 domain. Mutations in this domain in Burkitt's lymphoma, a human germinal centerderived B-cell lymphoma, were shown to be competent in binding to the catalytic subunit of PI3-kinase p110, yet specifically deficient in restricting the catalytic activity of p110 [Mellor et al., 2012]. Interestingly, transgenic cooverexpression of MYC and the constitutive active mutant of p110 was recently shown to suffice to develop Burkitt's lymphoma-like tumors in mice [Sander et al., 2012]. This raised the possibility that activation of the PI3-kinase signaling pathway by mutation in PIK3R1 had been one of the earliest events in DT40-cell development. However, this activation could be only transient, because DT40 cells were also shown to be competent in inducible activation of PI3-kinase by oxidative stresses [Qin and Chock, 2003]. One possible explanation for this could be the counterbalancing upregulation of the PTEN phosphatase [Neiman et al., 2001], which directly dephosphorylates lipid products of PI3-kinase.

In the same report, homozygous frameshift mutation of the *ATRX* gene resulting in truncation at K1392 was reported to be another possible contributor in the transformational process of DT40 cells [Molnár et al., 2014]. Since ATRX helicase was shown to facilitate various transactions at tandem repeat-rich genomic loci [Law et al., 2010; Clynes and Gibbons, 2013], it is tempting to associate ATRX defects with enhanced deletions at repeats in DT40 cells. However, we noticed that the K1392-encoding 'exon' is only called in the EMBL, but not in NCBI, databases and that BLAST search with this 'exon' sequence in the NCBI EST database retrieved no chicken ESTs. Therefore, further evaluation would be required to determine whether the observed mutations really affect the function of ATRX in DT40 cells.

Mechanisms Maintaining Genomic Integrity of DT40 Cells

In order to understand the molecular mechanisms how vertebrate cells maintain genomic integrity, numerous DT40 mutant cells have been generated and analyzed. DNA repair pathways so far extensively studied include 2 major DNA double-strand repair pathways, i.e. HR and non-homologous end joining (NHEJ), the translesion synthesis (TLS) repair pathway and the Fanconi anemia (FA) pathway [Sale et al., 2006; Sonoda et al., 2006; Takata et al., 2009]. Essentiality of the Rad51 recombinase for vertebrate cellular viability [Sonoda et al., 1998] and differential preference of HR versus NHEJ in phases of the cell cycle [Takata et al., 1998] are the well-recognized observations that came from early DT40 studies. TLS facilitates the restoration of stalled replication forks at damaged templates by extending DNA synthesis past damage by specialized DNA polymerases [Sale et al., 2006]. The FA pathway is mainly composed of 2 major protein complexes and believed to coordinate the selection and/or order of recruitment of DNA repair factors to resolve interstrand DNA crosslinks [Takata et al., 2009]. We demonstrated that DT40 cells singly mutated in FA genes are partly defective in HR and TLS and that double mutants defective in both FA and HR/TLS are epistatic in interstrand crosslink repair [Takata et al., 2009]. Although other DNA repair pathways such as mismatch repair, nucleotide excision repair and base excision repair are much less explored, competency of DT40 cells in these pathways was verified [Wakasugi et al., 2007; Asagoshi et al., 2010; Campo et al., 2013]. Underscoring the accumulating evidence from these reverse genetic studies, no apparent loss-of-function mutations were observed in the whole-genome sequencing analysis [Molnár et al., 2014]. Even if thousands of nonsynonymous SNVs and hundreds of coding InDels were identified to be unique in DT40 cells compared with cells from different breeds, many of them likely had pre-existed in the original animal. Therefore, comparison of unique mutations found in DT40 cells with those in Hyline SC progenitors might allow better defining DT40-specifc mutagenic events. Further evaluation of such mutations could shed lights on yet unclear mechanism(s) explaining the high targeting efficiency in DT40 cells.

RNA-seq: Primary Cells, Cell Lines and Heat Stress

(Prepared by C.J. Schmidt, E.M. Pritchett, L. Sun, R.V.N. Davis, A. Hubbard, K.E. Kniel, S.M. Markland, Q. Wang, C. Ashwell, M. Persia, M.F. Rothschild, and S.J. Lamont)

Transcriptome analysis by RNA-seq has emerged as a high-throughput, cost-effective means to evaluate the expression pattern of genes in organisms. Unlike other methods, such as microarrays or quantitative PCR, RNAseq is a target-free method that permits analysis of essentially any RNA that can be amplified from a cell or tissue. At its most basic, RNA-seq can determine individual gene expression levels by counting the number of times a particular transcript was found in the sequence data. Transcript levels can be compared across multiple samples to identify differentially expressed genes and infer differences in biological states between the samples. We have used this approach to examine gene expression patterns in chicken and human cells, with particular interest in determining response to heat stress.

Three separate in vitro cell culture experiments were conducted using the White Leghorn chicken hepatocarcinoma LMH cell line, the primary chicken liver (e15) cells and the human colorectal cancer epithelial Caco-2 cell line. Cells were freshly plated at subconfluent density and grown for 18 h at 37°C. Six biological replicates were prepared for the LMH cells, while 3 biological replicates were prepared for the primary chicken hepatocytes and Caco-2 cells. After 18 h, half of the plates were transferred to 43°C for 2.5 h after which RNA was isolated, Illumina RNA-seq libraries prepared [Coble et al., 2014; Sun et al., 2015] and sequenced to a depth of 10 million or more reads by the University of Delaware DNA sequencing facility. In addition, RNA-seq libraries were prepared from 6 freshly isolated 7-day-old chicken livers. Sequence data were analyzed using the Tuxedo software suite (Gal4 reference build and annotation) and differentially enriched genes identified as indicated below. Signaling and metabolic pathways affected by enriched genes were identified using GallusReactome+ (http://gallus.reactome.org/) and eGIFT [Tudor et al., 2010].

RNA-seq data were combined using chicken:human orthologs to merge the data [Burt et al., 2009] yielding a total of 5,333 pairs. Initially, hierarchical clustering was used to compare overall gene expression patterns across the cells and tissues (fig. 13). Although prepared at 2 different developmental time points (e15 vs. d7), the gene expression patterns of the primary liver culture and freshly prepared liver samples cluster together. The chicken



Fig. 13. Hierarchical clustering of RNA-seq data from primary cultured liver cells (TCLiver), fresh liver, chicken White-Leghorn hepatocarcioma cell line (LMH) and human colorectal cancer cell line (Caco-2).

LMH and human Caco-2 cells segregate into a second cluster. Overall, the clustering indicates that the primary cultured liver cells are more similar to fresh liver samples than the established cell lines. The clustering of the cell lines likely reflects the transformed nature and long-term culturing of the LMH and Caco-2 lines. Comparing iTerms [Tudor et al., 2010] for genes enriched in either the cell line samples or the liver samples (fig. 14) indicates enrichment for genes involved in cell proliferation in the cell lines. In contrast, the primary liver and cultured liver samples are enriched for genes involved in functions associated with primary hepatic metabolic processes such as coagulation, cholesterol metabolism and bile production. The latter observation suggests also that the shortterm in vitro cultured liver cells may model responses of the intact liver.

Heat Stress

One of the objectives of our work is identifying pathways responsive to heat stress, with a particular interest in identifying evolutionarily conserved responses. To this end, genes were identified that were responsive to heat stress in the cultured liver, LMH and Caco-2 cells. The genes were then mapped to *Gallus*Reactome⁺ [Sun, manuscript in preparation] to identify signaling and metabolic pathways that were modulated (p value <0.05) by heat stress in all 3 cell types. A total of 280 genes mapped to *Gallus*Reactome⁺ and affected 22 pathways (table 17). These pathways impact many cellular activities, including



Fig. 14. Stacked histogram of iTerms describing biological processes affected by genes enriched in the primary cultured liver samples plus fresh liver (red) compared with genes enriched in the LMH plus Caco-2 cell lines (blue).

the immune system, basic metabolism, response to growth factors and other stimuli, membrane transport of small molecules along with the integrity of the extracellular matrix. Presumably, these represent a core set of pathways that are fundamental to the overall heat stress response.

To identify a conserved set of genes that were responsive to heat stress in all 3 cultured cell types, pairwise comparisons were conducted of the 3 experiments to identify genes that were regulated in the same direction (up or down) whose difference was statistically significant (p < 0.05) (table 18). The transcription levels of 17 genes were differentially regulated by heat stress across all 3 cell types; 16 were upregulated by heat stress while 1 was downregulated. Eleven of the upregulated genes function directly as molecular chaperones or as co-chaperones (HSPB8, SERPINH1, BAG3, HSPH1, HSP90AA1, HSPA5, DNAJA1, CHORDC1, HSPA4, HSPA4L, and possibly TPPP). Several of the gene products, such as HSP90AA1, HSPA4 and HSPB8, serve as chaperones for multiple client proteins with broad effects on cellular pathways. Some of these gene products function in specific responses. For example, HSPA5 functions during endoplasmic reticulum stress, while SERPINH1 affects collagen folding and assembly. CHORDC1 binds the nucleotide-binding domain of HSP90 when the ADP:ATP ratio is high suggesting CHORDC1 may modulate HSP90 activity as a function of energy balance [Gano and Simon, 2010].

The hypothesis that TPPP may function as a chaperone or co-chaperone arises from several observations. **Table 17.** Pathways affected by heat stress in in-vitro-cultured pri-mary chicken liver, LMH and Caco-2 cells

*Gallus*Reactome⁺ pathways

Adaptive immune system **Biological** oxidations Collagen formation Degradation of the extracellular matrix Generic transcription pathway Innate immune system Ion channel transport Metabolism of amino acids and derivatives Metabolism of carbohydrates Metabolism of lipids and lipoproteins Peptide hormone metabolism Platelet activation, signaling and aggregation Post-translational protein modification Regulation of insulin-like growth factor (IGF) transport and uptake by IGF-binding proteins (IGFBPs) Regulation of mRNA stability by proteins that bind AU-rich elements Signaling by ERBB2 Signaling by ERBB4 Signaling by GPCR Signaling by NOTCH Signaling by NGF SLC-mediated transmembrane transport Transmission across chemical synapses

One observation is from this work showing that TPPP transcription is induced in chicken and human cells by heat stress. Furthermore, the *TPPP* gene product is known to modulate the tubulin network, promoting tubulin polymerization, microtubule acetylation and bundling, along with alpha-synuclein autophagy [Tirian et al., 2003; Tokesi et al., 2010; Ejlerskov et al., 2013]. Taken together, this evidence suggests that TPPP may function as a chaperone of proteins that interact with the tubulin matrix. While predominately studied in neuronal cells, TPPP may have similar functions in other cells.

Other products encoded by these heat-responsive genes may not function as chaperones. OTUD1 is a deubiquitinase that regulates the level of type 1 interferon in human cells and functions as a suppressor of the innate immune response [Kayagaki et al., 2007]. The deubiquitinase activity may allow OTUD1 to serve as a broad regulator of the ubiquitin protein degradation pathway during the unfolded protein response of heat stress. TRA2 is a nuclear protein, originally identified as controlling sexspecific splicing in *Drosophila* [Baker et al., 1989; Goralski et al., 1989; Amrein et al., 1990] and controls alternative splicing in vertebrates [Zhang et al., 2007]. Possibly, heat stress induction of TRA2 leads to an evolutionarily conserved pattern of alternative splicing [Strasburg and Chiang, 2009].

Three of the conserved heat-responsive genes function either extracellularly or by interaction with the extracellular matrix. OTOA was originally identified as a protein that links the apical surface of inner ear epithelial cells to the extracellular gel [Zwaenepoel et al., 2002]. IMPG2 encodes a hyaluronic binding proteoglycan [Acharya et al., 2000]. The final heat-responsive upregulated gene, SNTB2, encodes a member of the syntrophin family that links the cytoskeleton with the extracellular matrix [Ervasti and Campbell, 1993]. The sole gene downregulated by heat stress across all 3 cell types was VRK1, vaccinia related kinase 1. This kinase functions during the cell cycle and is required for transiting the G1 phase [Valbuena et al., 2008]. Downregulation by heat would likely reduce the rate of cell proliferation, possibly permitting more time to repair stress-associated cell damage.

The great depth of reads provided by RNA-seq can challenge assertions that a gene is only expressed in a specific cell or tissue. For example the OTOA gene was originally identified as expressed only in inner ear, while the IMPG2 gene has been identified as specific to retinal and pineal tissue [Acharya et al., 2000]. However, we detect low levels of both OTOA and IMPG2 transcripts in all 3 cell types. Inspection of the RNA-seq coverage for both genes revealed multiple reads mapping to the corresponding transcripts with the preponderance coming from exons (fig. 15). Also, some of the reads span an intron. Taken together, these observations suggest that the detected reads arise from mature transcripts rather than precursor RNAs or genomic contamination. The conclusion is that the reads from these genes in our experiments arose from true OTOA or IMPG2 transcripts. However, there is no current evidence that the OTOA or IMPG2 transcripts give rise to protein. This demonstrates an important current challenge: linking data from high-throughput sequencing and proteomics analysis to verify translation of such mRNAs.

Future Directions

A major application of RNA-seq transcriptome is determining gene expression profiles. Currently, many such studies focus on identifying genes based upon the annotated genomic sequence for the target organism. However, RNA-seq data can provide a rich source of data for transcripts that are currently not recognized in a genome's annotation file [Smith et al., this report]. Combining data from short-read (such as data generated by Illu-

Symbol	Description	p value ^a			Mean RPMK values					
		TC_Liver	r Caco-2	LMH	TC_Liver		Caco-2		LMH	
					control	heat	control	heat	control	heat
HSPA5	heat shock 70-kDa protein 5	0.0002	0.0088	< 0.0001	418	2,411	265	562	316	1,710
HSPB8	heat shock 22-kDa protein 8	0.0004	0.0145	0.0001	1.17	12.4	19.1	52.7	41.2	1,484
HSP90AA1	heat shock protein 90-kDa alpha 1	< 0.0001	0.0423	< 0.0001	52.4	549	239	943	205	1,369
BAG3	BCL2-associated athanogene 3	0.0002	0.0307	< 0.0001	47.6	665	14.8	198	33.8	806
HSPH1	heat shock 105-kDa/110-kDa protein 1	< 0.0001	0.0180	< 0.0001	12.6	183	43	468	47.9	373
TRA2A	transformer 2 alpha	0.0002	0.0135	< 0.0001	36.5	68.9	33	63.8	79.9	194
HSPA4L	heat shock 70-kDa protein 4	< 0.0001	0.0417	0.0003	30.5	91.3	2.84	33.9	78.7	175
DNAJA1	DnaJ (Hsp40)	< 0.0001	0.0099	< 0.0001	28.3	114	161	509	54.1	171
CHORDC1	cysteine and histidine-rich domain containing 1	0.0009	0.0366	< 0.0001	7.76	29.6	62	115	39.4	124
OTUD1	OTU domain containing 1	0.0055	0.0146	0.0021	15.1	164	8.03	21.3	14	104
HSPA4	heat shock 70-kDa protein 4	< 0.0001	0.0342	< 0.0001	18.5	66.7	65.4	117	29.1	88.8
TPPP	tubulin polymerization promoting protein	0.0064	0.0451	0.0076	0.42	0.89	0.09	0.43	11.6	26.6
SNTB2	syntrophin, beta 2	0.0018	0.0072	0.0014	2.29	4.93	4.98	9.81	6.97	15.4
VRK1	vaccinia-related kinase 1	< 0.0001	0.0106	0.0010	5.54	2.03	22.2	6.79	24.8	7.23
SERPINH1	serpin peptidase inhibitor, clade H-1	0.0005	0.0346	< 0.0001	31.1	87.8	231	799	0.29	2.41
OTOA	otoancorin	0.0018	0.0289	< 0.0001	0.14	0.96	0.02	0.12	0.53	2.19
IMPG2	interphotoreceptor matrix proteoglycan 2	0.0002	0.0315	0.0024	0.11	1.09	0.06	0.12	0.39	1.14

^a p value for difference between heat-stressed and control samples derived from primary cultured liver (TC_Liver), LMH cells and Caco-2 cells.

mina) and long-read (PacBio) RNA-seq technologies can provide an even better understanding of transcript structure, significantly improving genome annotation. Alternative splicing, allele-specific expression, microRNAs, and lncRNAs can all be identified with RNA-seq data, providing a rich catalog of the diversity of transcripts found in samples. Given the (relative) ease with which this data can be collected, an important current need is improving and extending bioinformatics tools. While many such tools exist, the community needs to work together to develop intuitive web-based bioinformatics pipelines and platforms that are appropriate to users with varying levels of computer sophistication. Ultimately, combining an incredibly rich set of sequence data with user-friendly bioinformatics tools will provide complete gene expression profiles of organisms in many different biological states.

Host-Viral Genome Interactions in Marek's Disease

(Prepared by M.C. McPherson, C.M. Robinson, and M.E. Delany)

Marek's Disease and Marek's Disease Virus

Marek's disease (MD) encompasses a variety of symptoms in its host, including paralysis, blindness, immunosuppression, and lymphoma formation, depending on viral serotype, bird genotype and immune status of the affected chicken. In the 1960s, the symptoms of MD were attributed to a herpesvirus, the Marek's disease virus (MDV). The virus was named after József Marek who first identified a widespread polyneuritis common amongst birds in 1907 [Gimeno et al., 1999]. In 1982, electronmicroscopy data identified MDV as an alphaherpesvirus, and sequencing efforts in 2000 further confirmed its alphaherpesvirus designation [Lee et al., 2000; Tulman et al., 2000]. We currently understand that virulent MDV undergoes 4 overlapping infection stages that contribute to viral persistence in a host bird; early cytolytic, latent, late cytolytic, and cell transformation [Addinger and Calnek, 1973; Osterrieder et al., 2006; Gimeno et al., 2011]. During the early cytolytic infection, the virus is actively spreading among host immune cells, and the MDV genome is replicated and packaged into infectious virion particles, to be shed into the environment with the host feather dander. The viral load in the immune tissues peaks during this initial stage. Following the lytic phase

Fig. 15. Track diagrams from UCSC browser plotting chicken LMH RNA-seq reads from *OTOA* (**a**, **b**) and *IMPG2* (**c**, **d**). **a** and **c** correspond to the entire gene region, while **b** and **d** have been zoomed in to show coverage of individual exons along with reads that span the intron indicating splicing events.

(For figure see next page.)



Third Report on Chicken Genes and Chromosomes 2015

Cytogenet Genome Res 2015;145:78–179 DOI: 10.1159/000430927 C. Steinlein - 60989
of infection, latent virus (non-replicating) is established in activated T lymphocytes around 7 days post-infection (dpi) [Baigent and Davison, 2004; Trapp et al., 2006; Arumugaswami et al., 2009]. These T lymphocytes are capable of being transformed, typically around 21 dpi, resulting in lymphoma formation [Witter, 1997; Trapp et al., 2006].

MVD Vaccines and the Poultry Industry

The uncovering of the oncogenic herpesvirus etiology was followed by development of effective vaccines in the early 1970s. The MD vaccines were the first vaccines capable of preventing a cancer in both veterinary and human medicine, making this aspect of MDV research particularly exciting. The Rispens vaccine, developed via serotype 1 Rispens virus attenuation, has proven to be the most effective against very virulent (vv) and very virulent plus (vv+) MDV. Inoculation of birds with herpesvirus of turkey (HVT), a serotype 3 virus, and SB-1, a serotype 2 virus, also prevent MD, but are generally less effective against vv and vv+ strains. Vaccination is currently administered either in ovo or at hatch. Vaccines are surmised to prime the chicken immune system, although they do not prevent MDV infection [Islam et al., 2002]. This process is not well understood and supplementary mechanisms in preventing and/or reducing MD-lymphoma development are doubtlessly involved [Witter, 1984; Osterrieder et al., 2006]. It is a well-studied feature that bird genotype for the major histocompatibility complex confers resistance (e.g. B21) or susceptibility (e.g. B19) to MDV. The poultry industry applies variable MD immunization strategies, administering either monovalent (e.g. Rispens) or polyvalent (e.g. SB-1/HVT) vaccines. Nonetheless, concerns are growing as increasingly virulent MDV strains emerge [Nair, 2005], heightening the need for an enhanced mechanistic understanding of all serotypes of MDV and further vaccine development.

Cytogenetic Investigations of Host-Virus Interactions

Here, we will describe MDV and host genome interactions and review current methods that incorporate genetic tools to more precisely investigate genome-level host-virus interactions. These studies and others have helped elucidate the role of chromosomal integration of MDV in disease pathogenesis.

Initial Studies Detect MDV Integration in Host Telomeres

It was initially understood that all MD-transformed cells maintained a latent infection with the virus; how-

ever, the status of viral DNA in the transformed cells was unknown. Delecluse and Hammerschmidt [1993] were the first to determine that MDV integrated into host chromosomal DNA in transformed cell lines developed from infected birds. Integration into host DNA was supported by detection of doublet MDV FISH signals from sister chromatids, indicating viral DNA was replicated along with the host genome. Integration loci mapping through FISH was limited to the largest macrochromosomes due to a lack of identifying features for the microchromosomes. Integration sites were typically distally-located at the telomeres on the macrochromosomes. Integration profiles were recurrent from cell-tocell within a line, yet differed from the profiles found in other cell lines. Free linear virion, representing processed MDV genomic DNA during the lytic stage, was also detected at low levels in some of the transformed cell lines.

MDV FISH - Current Methods

The availability of chicken BAC clones, many of which have been assigned to a particular macro- or microchromosome [Lee et al., 2003], as well as MDV cosmid and BAC clones, which contain the viral genome or a segment of it [Lee et al., 2000; Tulman et al., 2000; Petherbridge et al., 2003, 2004; Baigent et al., 2006; Niikura et al., 2006; Silva et al., 2010], unlocked new possibilities in cytogenetic research. The former allowed the identification of specific chromosomes [Romanov et al., 2005; Delany et al., 2007], which could not be distinguished by their morphological features alone (i.e. GGA6-28), while the latter was utilized to investigate interactions between the host and viral DNA. Importantly, a number of MDV clones are currently available, including those of non-oncogenic MDV strains (i.e. HVT, SB-1, Meq-deleted). Multi-color FISH techniques, wherein a labeled MDV probe (of one color) was hybridized to interphase nuclei and metaphase chromosomes along with differentially labeled chromosome- and/or telomere-specific probe(s) (of a 2nd and 3rd color), were used for the chromosome mapping of viral DNA integrations and analysis of viral DNA status over time and in different tissues [Robinson et al., 2010].

Cytogenomic Analysis

Robinson et al. [2014] reported 4 viral genome status phenotypes in dividing lymphocytes interrogated by FISH with an MDV-BAC probe. The 4 phenotypes are (1) MDVnull, characterized by the complete absence of fluorescence signals in the nucleus or around the chromosomes, (2) MDV chromosome-associated, represented by distinct flu-

orescence signals around and over the host chromosomes, (3) MDV chromosome-associated/integrated, showing the associated signals as well as punctate fluorescence signals integrated into the sister chromatids at one or more telomeres, and (4) MDV chromosome-integrated only, distinguished by the presence of the distinct, punctate signals at the telomeres with no other signals. Figure 16 shows representative examples of the 4 phenotypes. The interpretations of the phenotypes are that MDV-null cells are free of MDV DNA, indicating the absence of the virus in these cells. The MDV chromosome-associated phenotype suggests cells in the cytolytic stage (free virus evident), while the chromosome-associated/integrated phenotype indicates both the presence of free episomal virus and viral DNA that has integrated into the host genome. The integrated-only phenotype is interpreted to reflect a latently-infected cell, cleared of replicating virus. The interpretations are based both on the appearances and known timing of the disease stages. The appearance of the associated-only MDV phenotype in host immune tissues is clearly linked with the timing and expected viral DNA 'behavior' of the lytic replication stage (1–7 dpi). However, the biological implications of the associated/integrated MDV phenotype are less clear, as it is not known whether viral replication is still occurring after copies of the MDV genome become integrated into the host DNA and precisely how a latent infection is established. The appearance of the MDV chromosome-integrated only phenotype shows significant correlation with the timing of viral latency (7 dpi) and is the predominate phenotype ultimately found in MD-induced lymphomas [Robinson et al., 2010, 2014].

Host-Virus Genome Interactions: Investigations of MDV and the Chicken Genome

MDV-infected chickens provide a unique research model for investigation of herpesvirus infection, virusinduced lymphoma formation and viral latency and cellular persistence, as well as interface between these events. Cytogenetic investigations provide a precise assay of the physical behavior of the virus at the level of the genome.

Fig. 16. Representative images of the MDV and chicken host genome interaction phenotypes in mitotically dividing cells of the spleen of virulent GA-challenged birds. **a** MDV null (no virus detected); **b** MDV chromosome-associated (interpreted as lytic stage of virus replication); **c** MDV chromosome-associated and telomere-integrated (transitional state with virus replicating and integrated into the chicken genome); **d** MDV telomere-integrated only (interpreted as latent stage). Chicken chromosomes are counterstained with DAPI (blue), and the FITC (green) signals represent MDV DNA.



Cytogenet Genome Res 2015;145:78–179 DOI: 10.1159/000430927 151

Third Report on Chicken Genes and Chromosomes 2015

Host-MDV genome interactions appear to have a significant role in the virus' achievement of latency, cellular transformation and reemergence to lytic infection. Understanding the role of these interactions may also help identify new vaccine candidates and characterize emergent, highly virulent MDV strains.

Chromosomal Association during Timing of Cytolytic Replication

Viral genomic DNA reaches considerable levels in host tissues as early as 4 dpi based on quantitative data [Baigent and Davison, 1999]. However, Robinson et al. [2014] detected MDV chromosome-associated FISH signals as early as 1 dpi within the immune organs (bursa, thymus and spleen) of GA-infected birds, thus providing earlier evidence for the lytic replication in host organs even before 4 dpi. Immune tissues, which represented a population of primarily T and B lymphocytes (levels varying by tissue type), showed a peak level of MDV chromosome-associated phenotype cells around 4 dpi, which significantly and continuously declined at 14 and 21 dpi. This is consistent with data indicating a peak of lytic replication between 4 and 7 dpi [Baigent and Davison, 2004].

Telomere Integration by MDV: Mechanism and Mapping

Before the experimental detection of MDV integration into chicken chromosomes, segments of the MDV genome with sequence homology to chicken telomeric repeats were reported in the literature [Kishi et al., 1991]. Shortly thereafter, Delecluse and Hammerschmidt [1993] provided clear evidence of MDV integration into host telomeres. Later studies offered further evidence of a physical interaction between viral and host DNA by detecting viral genes with homology to chicken, including viral interleukin-8 and telomerase RNA [Parcells et al., 2001; Fragnet et al., 2003], and providing direct proof of MDV's capacity to acquire host DNA into its own genome [Niikura et al., 2006]. In 2010, Robinson et al. [2010] determined that oncogenic strains were integrated specifically at the telomeres within chicken MD-induced lymphomas. In corroboration with Delecluse and Hammerschmidt's [1993] findings, doublet FISH signals were observed, indicative of integrated MDV in sister chromatids, and the vast majority of mitoses displayed the MDV chromosome-integrated only phenotype. Notable fluorescence signal strength variation was detected within and across cells and lymphomas, suggesting that a variable number of tandemly repeated MDV genomes integrated into host DNA differing considerably from one integration event to the next. With regard to random, preferred or targeted integration into specific chromosomes, temporal analysis showed strong evidence for random integration initially (although with specific targeting to the terminal telomeric DNA) [Robinson et al., 2014], but there appear to be some 'preferred' sites for viral integration as per the MD tumor data [Robinson et al., 2010], e.g. GGA4, 6, 9, 12, and 20. This fine-tuning may be related to the cell-selection process in transformation and/or tumorigenesis. The mapping analysis in the same cytogenetic study examined the hypothesis as to preferential integration into chromosomes with mega-telomeres, which was not supported. It was later established that herpesvirus telomeric repeats facilitate viral genome integration into host telomeres [Kaufer et al., 2011]. These authors found DNA-junction fragments containing both hosttelomeric and MDV DNA from infected lymphoblastoid cells. These unique fragments were absent when the birds were challenged with the mutant- or deleted-telomeric repeat MDV strains, suggesting such mutant viral genomes exhibit a failure to integrate at the host telomeres. Combined, these data suggest that the mechanism of integration by MDV may involve the host telomere extension pathway or homologous recombination events.

MDV Integration and Latency

As a herpesvirus, MDV enters the stage of latency becoming quiescent and thus evading further host immune attack. The latent stage of infection is characterized by reduced gene expression, decline in viral load (as compared to earlier timepoint in infection) and persistence in the host lymphocytes [Stevens, 1989]. Delecluse and Hammerschmidt [1993] were the first to establish that MDV integration is associated with latently-infected chicken cells, although the nature of the relationship between these 2 events remained unclear. Latently-infected cell lines also maintain very little to no extrachromosomal circular or linear viral DNA. Robinson et al. [2014] confirmed that the telomere-integrated only form of MDV first emerges around 7 dpi (at the timing of latency) and increases in frequency at 14 and 21 dpi in bursa, thymus and spleen. However, it is unclear as to whether the act of integration into host telomeric DNA is necessary to achieve the latent stage or, conversely, if viral integration requires latency-related events to occur. Interestingly, lymphoblastoid cells infected with mutant MDV strains, which cannot integrate at host telomeres, do not show significant changes in latency-associated gene expression (e.g. Meq and vTR) as compared to cells infected with oncogenic wild-type MDV [Kaufer et al., 2011].

MDV Integration and Oncogenesis

Multiple studies have indicated a biological relationship between MDV integration into host telomeres and cellular transformation events leading to lymphoma formation in infected, MD-susceptible birds. MDV shares this trait with a number of other oncogenic viruses, including human hepatitis B and papilloma virus-18, which integrate into the genomic DNA of the host cells they later transform [Popescu et al., 1990]. Nearly all cells within primary MD-lymphomas or transformed cell populations have MDV integrated into one or more telomeric sites [Delecluse and Hammerschmidt, 1993; Robinson et al., 2010, 2014]. Furthermore, very low levels of lytically replicating or free viral DNA are detected from these cell populations [Delecluse and Hammerschmidt, 1993]. Examination of the MDV integration profiles or the distinct telomeric sites of integrated MDV in a nucleus, revealed evidence of both monoclonal and polyclonal tumors [Robinson et al., 2010]. Profiling of MD-induced tumors indicated more heterogeneity in viral-integration profiles within and between early-stage (21 dpi) tumors, whereas late-stage (61 and 73 dpi) tumors demonstrated more homogeneity [Robinson et al., 2014]. The proposed model is that early-stage tumors represent a collection of recently transformed T lymphocytes, many of which do not persist to generate an enhanced cell lineage. The late-stage tumors may mostly represent one transformation event that was selected further by the processes of tumorigenesis to become the predominant cell lineage in the initial tumor, and leading to metastases, resulting in monoclonal tumors in other tissues. In vivo experiments demonstrated that birds infected with non-integrating MDV strains have significantly decreased lymphoma development, providing further evidence of the link between MDV integration into host telomeres and achievement of cellular transformation [Kaufer et al., 2011]. The remaining question is; how does MDV integration contribute to oncogenesis? One hypothetical answer is that integration helps to establish latency, which in turn is concomitant with viral oncogene expression.

Telomeric Integration by Non-Oncogenic MDV Strains

Recently, we characterized in vivo interactions of MDV vaccine-viruses, specifically Rispens, SB-1 and HVT, with the host genome between 1 and 21 dpi [McPherson et al., 2014, manuscript in preparation]. The MDV chromosome-associated/integrated cell phenotype was detected across all vaccinated birds, indicating that MD vaccine-viruses are capable of integrating into chick-

en telomeres. In contrast with the oncogenic MDV-challenged birds, the vaccinated birds indicated a sustained and higher cell population with free, replicating virus (chromosome-associated virus phenotype) from 4 to 21 dpi. The MDV telomere-integrated only phenotype was not established within spleen cell populations of vaccinated birds during early infection. Interestingly, the nononcogenic Meq-deleted MDV strain (Δ Meq) [Silva et al., 2010] also maintains a consistently high level of lytic chromosome-associated signals and does not establish the MDV chromosome-integrated only phenotype in host cells from 1 to 21 dpi [Robinson et al., 2014]. The MDV chromosome-associated/integrated phenotype was detected by FISH in the spleen, bursa and thymus of Δ Meq-infected birds, indicating that the strain is capable of telomeric integration, yet does not transform host lymphocytes in vivo and may not have a capacity for latent infection. These results provided further evidence that Meq expression and telomeric integration by MDV play crucial roles in the establishment of a latent infection and transformation of host cells. Collectively, the data on non-oncogenic MDV strains clarifies that telomeric integration is not sufficient for cellular transformation, although it does appear to be necessary. Reactivation of Lytic Replication in Latently-Infected Host Cells

MD-induced lymphomas principally consist of latently-infected and transformed host cells [Delecluse et al., 1993; Robinson et al., 2010, 2014]. However, an increased number of cells with lytic DNA replication (free linear virion) are detected from MD primary lymphomas in culture conditions and, particularly, after exposure to bromodeoxyuridine [Delecluse et al., 1993; Robinson, 2013]. UA04, a unique latently-infected T lymphocyte line [Dienglewicz and Parcells, 1999], was used to study the behavior of reactivated MDV to investigate the hypothesis that integrated MDV was capable of emerging from its chromosome-embedded locus. Analysis of the cell line showed 2 MDV integration loci. Upon incubation with bromodeoxyuridine, which induces the latent virus to become lytic [Dunn and Nazerian, 1977], multiple, speckled, smaller FISH signals surrounding and associated with the integration loci ('starbursts of virus signal') appeared. The FISH signals in close proximity to MDV integration sites are interpreted as having 'escaped' from their integration loci and are free replicating virus (fig. 17). It has also been found that the presence of telomeric repeats in the MDV genome is crucial to viral reactivation [Kaufer et al., 2011]. These combined data suggest that



Fig. 17. The status of MDV in UA04 metaphase chromosomes and interphase nuclei in control and bromodeoxyuridine (BrdU)-exposed cells. UA04 was found to have a lineage that was trisomic for GGA1. **a** Latently infected control cells (no BrdU treatment), exhibiting MDV-integrated loci at the telomeres of 2 homologs of GGA1p (2 punctate FISH signals). **b** UA04 cells exhibiting 'associ-

ated MDV' signals surrounding and dispersed from (beyond) integrated-MDV loci after BrdU treatment for 48 h. These associated (non-integrated) FISH signals are hypothesized to represent escaped and reactivated replicating MDV genome. Chicken chromosomes are counterstained with DAPI (blue), and the FITC (green) signals represent MDV DNA.

MDV may not only use integration as a means for avoiding immune detection (latency), but also as a mode of persistence in the host with the means to 'escape' integration within the host genome and return to the replication phase of its life cycle. The ability to return to a state of productive replication after integration implies that the MDV genome is integrating intact (without rearrangement) into the host DNA. The mechanism by which the integrated MDV genome is excised or generates free virus involves the viral telomeric repeats; however, the exact host and/or viral machinery involved offers opportunity for future study.

Transcriptome Variation in Response to Marek's Disease Virus Acute Infection

(Prepared by L. Preeyanon, C.T. Brown, and H.H. Cheng)

Marek's disease (MD) is an economically significant chicken disease that affects the poultry industry worldwide with an estimated annual cost of USD 2 billion [Morrow and Fehler, 2004]. The disease is caused by the highly oncogenic Marek's disease virus (MDV), an alphaherpesvirus that induces T-cell lymphomas in susceptible birds. Vaccination is the primary control measure, which is effective in reducing incidence of tumor formation. However, since MD vaccines are not sterilizing, they do not prevent infection or horizontal spread of the virus. As a consequence, MDV field strains that overcome vaccinal protection have arisen repeatedly over time [Atkins et al., 2013]. Therefore, sustainable alternative control measures, such as improving genetic resistance, are needed.

Many studies have reported strong associations between MHC alleles and resistance or susceptibility to MD. For example, chickens with the MHC allele B^{21} are resistant in contrast to chickens with the B^{19} allele, which are susceptible. ADOL lines 6 and 7, both share the same MHC B^2 allele, yet exhibit different phenotypic responses; e.g. challenge with the JM/102W strain typically results in 0 and 100% MD incidence for lines 6 and 7, respectively. Thus, the major unanswered questions are what genetic factors, especially those that are non-MHC, contribute to susceptibility and resistance to the disease, and what are the main contributing mechanisms?

Significant efforts have been made to study variations in global gene expression between MD-resistant and -susceptible birds using microarray and RNA-seq methods in order to identify non-MHC genes that contribute to resistance to MD [Bumstead, 1998; Vallejo et al., 1998; Yonash et al., 1999; Morgan et al., 2001; Sarson et al., 2008; Smith et al., 2011]. However, none of the studies have investigated differential expression of alternative isoforms, which are known to play a significant role in many biological events including immune responses



Fig. 18. Differentially expressed genes in response to MDV infection. The number of unique or common genes showing up- or downregulation in line 6 (MD resistant) or line 7 (MD susceptible) chicks at 4 days post-infection using splenic RNA is given in each Venn diagram.

Table 19. Splenic genes regulated in opposite directions between lines 6 (MD resistant) and 7 (MD susceptible) in response to MDVinfection at 4 days post-infection

Gene		Change in	expression ^a
Symbol	name	line 6	line 7
LL	lung lectin	-3.36	+8.71
GIF	gastric intrinsic factor	-2.15	+3.11
SFTPA1	surfactant protein A1	-4.84	+3.73
SCAF8	SR-related CTD-associated factor 8	-8.70	+8.24
C14ORF1	chromosome 14 open reading frame 1	-3.11	+2.71
PPARG	peroxisome proliferator-activated receptor frame 1	-6.99	+2.06
NDUFA4	NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 4	+1.66	-1.03
RAD17	RAD17 homolog (S. pombe)	+1.45	-1.80
RPL39	ribosomal protein L39	+3.34	-1.63
ATP8A2	ATPase, aminophospholipid transporter, class I, type 8A, member 2	+7.66	-7.80
NDUFB3	NADH dehydrogenase (ubiquinone) 1 beta	+1.03	-1.34
PSMG3	proteosome assembly chaperone 3	+1.52	-1.26
MED9	dediator complex subunit 9	+8.23	-3.15
PNISR	PNN-interacting serine/arginine-rich protein	+5.58	-1.81
S1PR1	sphingosine-1-phosphatase receptor 1	+1.31	-6.96
CD7	CD7 molecule	+2.88	-1.38
LOC100858785	unknown	+1.26	-1.75
THOC7	THO complex 7 homolog (Drosophila)	+7.01	-6.90
DNAJA2	DnaJ (Hsp40) homolog, subfamily A, member 1	+2.34	-6.02

^a The – and + symbols represent the log₂ fold change down- and up-regulated genes in response to MDV infection, respectively.

[Lynch, 2004]. In addition, studies have shown that isoform expression levels can provide better signatures for some diseases [Zhang et al., 2013]. Changes of isoform expression levels are governed partly by 2 types of *cis*regulatory elements: exonic splicing enhancer (ESE) and exonic splicing silencer (ESS), both located within an exon sequence. A number of sequence motifs of ESE and ESS have been identified in human and other organisms and can be predicted in silico. Mutations that disrupt or create those motifs can alter splicing patterns leading to

aberrant alternative splicing. A number of disease-associated single nucleotide polymorphisms (SNPs) in coding regions that affect ESEs and ESSs have been well characterized [Blencowe, 2000; Wang and Cooper, 2007]. Therefore, variations in isoform expression could lead to identification of SNPs that underlie genetic resistance to MD. Here, we report differentially expressed genes and isoforms that may contribute to resistance to MD as well as SNPs that can potentially affect isoform expression levels.

Gene		Change in expression ^a		
Symbol	name	line 6	line 7	
IL2RG	interleukin 2 receptor, γ		0.55	
IL6	interleukin 6 (interferon, β2)		5.11	
IL6ST	interleukin signal transducer (gp130, oncostatin M receptor)		1.36	
IL8L1	interleukin 18-like 1	1.90		
IL18	interleukin 18 (interferon-γ inducing factor)	1.92	4.07	
IL15	interleukin 15		1.06	
IL18R1	interleukin 18 receptor 1	1.94	1.64	
IFNG	interferon-γ	5.14	4.90	
IFNB	interferon- $\hat{\beta}$	4.83	5.64	
IFNA3	interferon- α 3	4.09	5.48	
IFNGR1	interferon-γ receptor 1		2.04	
IFNGR2	interferon-y receptor 2		0.50	
IFNAR1	interferon- $\alpha_{\beta}\beta$ receptor 1		1.46	
IFNAR2	interferon- $\alpha_{\beta}\beta$ receptor 2		0.58	

Table 20. Cytokine-related gene expression in response to MDV infection in lines 6 (MD resistant) and 7 (MD susceptible) chicks

Differential Gene Expression Indicates Active Immune Responses to Ongoing Lytic Infection in the Susceptible Line

To study gene and isoform expression, we incorporated Ensembl gene models (release 73) with de novo and a reference-guided transcriptome assembly to build custom chicken gene models. The models, therefore, include both Ensembl-annotated transcripts and putative genes and isoforms. The advantage of using custom gene models is that it allows an investigation of unannotated genes and isoforms, which is necessary for in-depth study of gene expression. Based on these models, many genes were differentially expressed between control and infected chickens in both lines at 4 days post-infection. While the number of unique downregulated genes in both lines was approximately equal, the number of unique upregulated genes in the susceptible line was much greater compared to the resistant line (fig. 18). Interestingly, some genes that were differentially expressed in both lines were regulated in the opposite direction (table 19). Among genes downregulated in the resistant line but upregulated in the susceptible line were LL (lung lectin) and SFTPA1, which encode a calcium-dependent C-type lectin and a lung surfactant protein, respectively, both important in innate immunity [Kingma and Whitsett, 2006; Hogenkamp et al., 2008], and PPARG, a suppressor of the NFkB-mediated proinflammatory response. On the other hand, nearly all genes upregulated in the resistant line but downregulated

in the susceptible line are involved in cell survival such as mRNA splicing, cell growth and protein synthesis, except CD7 whose function is involved in T cell-B cell interaction. This difference suggests that even at this stage of infection, in the resistant line, the lytic phase could be repressed. Therefore, only genes involved in cell division are upregulated possibly to repair the initial damage due to infection in the resistant line. In comparison, the lytic phase in the susceptible line may still continue and as a result, genes involved in immune responses are still upregulated. In addition, type I interferons (IFN-y and IFN- β) as well as IFN- α 3 were found to be highly upregulated in infected chickens in both lines (table 20). However, genes encoding their corresponding receptors were not differentially expressed in the resistant line, but upregulated in the susceptible line. This could also reflect the ongoing immune response in the susceptible line.

Functional Analysis of Differentially Expressed Genes Indicates Inactive Adaptive Immune Responses in the Resistant Line

To determine pathways that were perturbed during the infection, data were analyzed by GOSeq, which accounts for gene length bias unique to RNA-seq data [Young et al., 2010]. Significantly perturbed pathways (FDR < 0.1) from both lines that are involved in immune response include the TLR signaling pathway, cytokine-cytokine receptor interaction, intestinal immune network for IgA produc-



Fig. 19. KEGG pathways enriched in response to MDV infection. Splenic RNAs from both line 6 (MD resistant, top group) and line 7 (MD susceptible, bottom group) chicks were analyzed for differential gene expression in response to MDV infection with the resulting pathways (y-axis) and corresponding p values (x-axis) shown.

tion, and cell adhesion molecules (fig. 19). Some other pathways important in response to viral infection and only significantly enriched in the susceptible line include phagosome, apoptosis, RIG-I-like receptor signaling pathway, NOD-like receptor signaling pathway, and lysosome. Although MHC class I (BF1) was differentially expressed in both lines, other genes involved in expressing newly synthesized MHC class I were only upregulated in line 7 (MD susceptible) suggesting that new MHC I molecules were actively produced. Furthermore, gene ontology analysis of biological processes (GO:BP) shows that categories involved in both adaptive and innate immune responses were enriched in the susceptible line (data not shown). On the other hand, only categories involved in innate immune responses were enriched in line 6 (MD resistant). In addition, enrichment of the apoptosis pathway in the susceptible line suggests that the programmed cell death could be induced by the cytotoxic T lymphocyte response to eliminate ongoing viral infection.

At this stage of infection, our results suggest that lytic infection of MDV stimulates both innate and adaptive immune responses, which leads to activation of T cells in the susceptible line. Only activated T cells are believed to be infected by MDV; therefore, the lytic phase could facilitate the spread of the virus by enhancing expansion of activated T cells. Due to the cell-associated nature of MDV, the viruses transfer to T cells via cell-to-cell contact between B cells and T cells during antigen presentation, or B cell activation by T helper cells. Therefore, it is beneficial for the host to restrain such contact. However, it is not clear how chickens in the resistant line control the lytic infection of MDV. Two mechanisms have been speculated to contribute to MD resistance. First, innate immune responses could be highly effective and could activate strong adaptive immune responses that rapidly control viral replication and force the viruses to enter into the latent phase. Second, the innate immune response itself could be highly effective in limiting viral replication [Smith et al., 2011].

157

Third Report on Chicken Genes and Chromosomes 2015

Туре	Ensembl ID	Symbol Line 6		Line 7		
			control	infected	control	infected
SE	ENSGALG00000011127	BCL11B	0.07	0.30	0.06	0.04
SE	ENSGALG0000013137	INO80C	0.15	0.35	0.95	0.86
A5SS	ENSGALG0000009824	C7H20RF77	0.49	0.26	0.68	0.62
A5SS	ENSGALG0000013821	GEMIN6	0.84	0.61	0.81	0.85
A5SS	ENSGALG0000002144	THRAP3	0.31	0.51	0.28	0.18
A3SS	ENSGALG00000027665	SYNGR1	0.46	0.23	0.68	0.60
A3SS	ENSGALG0000005685	KSR1	0.77	0.44	0.72	0.65
A3SS	ENSG00000163875 ^a	MEAF6	0.28	0.57	0.40	0.29
A3SS	ENGALG0000020987	ZDHHC7	0.42	0.23	0.57	0.55

Table 21. Group I: Genes showing differential exon usage (ψ) values only in line 6 (MD resistant) between uninfected and MDV-infected birds

SE = Skipped exon; A5SS = alternative 5' splice site; A3SS = alternative 3' splice site. ^a Human homolog.

Genes with Differential Exon Usage in Response to MDV Infection Can Be Divided into 4 Groups Based on Their Patterns of Expression

The immune system is isoform-rich and many genes express different isoforms with distinctive functions in response to stimuli such as stress, chemicals and infection [Stamm, 2002; Hagiwara, 2005]. Changes in expression of splice forms of immune-related genes have been reported to be associated with increased susceptibility to and poor prognosis of diseases [Lynch, 2004]. Studying differential isoform expression could therefore shed light into inherent differences between lines that confer resistance or susceptibility to MD.

In the past, microarray technology has been used to study gene and isoform expression in many studies, but its sensitivity for detecting structurally similar isoforms is low, and known or predicted annotations are required to design probes [Kane et al., 2000]. Although the RNA-seq method can provide a reliable estimate of exon expression compared to microarrays [Pan et al., 2008] and is not constrained to the same limitations, studying isoform expressions using RNA-seq is still not straightforward because the read lengths are not long enough to span across all exons in an isoform. In most cases, only exons in close proximity are covered by the same read, which makes it difficult to accurately predict a full structure of the isoform. In addition, some genes are fused due to overlapping untranslated regions, which can also result in erroneous predicted isoform structures.

Due to those issues, it is not feasible to accurately estimate expression of isoforms, especially when gene annotation is constructed from de novo assembly [Trapnell et al., 2013]. To avoid these issues, we chose to study exon expression instead of isoform expression. Using MISO [Katz et al., 2010] with the exon-centric method, only reads spanning across exons and used in splicing event are examined. The expression of exon inclusion is calculated as 'percent spliced in' (Psi or ψ), which can be used to infer the portion of transcripts that include the exon in each sample [Katz et al., 2010]. In this study, we investigated the 3 most common alternative splicing events in vertebrates, which are skipped exons, an alternative 3' and 5' splice site. Lists of differential exon usage genes from the resistant line that show differences in $\psi > 0.20$ when compared to the susceptible line in infected chickens are shown in tables 21-24. Genes can be categorized roughly into 4 groups based on the pattern of ψ across control and infected birds in both lines.

Group I (table 21) includes genes with ψ values that were up- or downregulated in infected chickens in line 6 (MD resistant) only. This group includes *BCL11B* (B-cell CLL/lymphoma 11B zinc finger proteins), a B-cell lymphoma associated C2H2-type zinc finger protein encoding a tumor-suppressor for T-cell lymphoma in humans. According to homologous alignments, a splice form with the skipped exon is similar to mouse BCL11B isoform b. The skipped exon was expressed 30% in MDV-infected line 6 chickens; whereas it was rarely expressed (4–7%) in the control line 6 birds and both groups in line 7 (MD susceptible). The skipped exon was not found to encode any known protein domain; however, it is in the middle of 2 adjacent C2H2-type finger protein domains.

Туре	Ensembl ID	Symbol	Line 6		Line 7	
			control	infected	control	infected
SE	ENSGALG0000005522	DYNLL2	0.01	0.02	0.20	0.25
SE	ENSGALG0000004971	URM1	0.07	0.03	0.18	0.23
SE	ENSGALG0000015709	TACC3	0.87	0.93	0.77	0.72
SE	ENSGALG00000014642	LOC374195	0.60	0.57	0.70	0.80
SE	ENSGALG0000011682	CNOT4	0.57	0.62	0.40	0.41
SE	ENSGALG0000007511	ITGB2	0.17	0.22	0.02	0.01
SE	ENSGALG0000006522	HCK	0.47	0.59	0.99	0.97
SE	ENSGALG0000000904	C11H16ORF57	0.92	0.98	0.84	0.78
A5SS	ENSGALG00000010836	AHR	0.97	0.99	0.63	0.59
A5SS	ENSGALG00000011488	CMTM7	0.55	0.67	0.37	0.41
A3SS	ENSGALG0000008939	FUBP1	0.42	0.26	0.59	0.54
A3SS	ENSGALG0000008507	THOC2	0.52	0.53	0.69	0.78
A3SS	ENSGALG0000002859	RAC3	0.69	0.84	0.67	0.61
A3SS	ENSGALG0000012050	TNRC6B	0.57	0.39	0.94	0.93
A3SS	ENSGALG0000010410	PFN2	0.71	0.78	0.53	0.50
A3SS	ENSGALG0000027908	LOC422528	0.28	0.39	0.13	0.09
A3SS	ENSGALG00000011476	SEPT11	0.78	0.86	0.60	0.58

Table 22. Group II: Genes showing similar differential exon usage (ψ) values within line between uninfected and MDV-infected birds but different ψ values between lines 6 (MD resistant) and 7 (MD susceptible)

SE = Skipped exon; A5SS = alternative 5' splice site; A3SS = alternative 3' splice site.

Table 23. Group III: Genes showing differential exon usage (ψ) values only in line 7 (MD susceptible) between uninfected and MDV-infected birds

Туре	Ensembl ID	Symbol	Line 6	Line 6		Line 7	
			control	infected	control	infected	
SE	ENSGALG0000003861	HERC4	0.31	0.37	0.45	0.06	
SE	ENSGALG0000009029	TSPAN12	0.08	0.15	0.20	0.47	
SE	ENSGALG0000009520	MARCH1	0.42	0.54	0.66	0.34	
SE	ENSGALG0000008320	EDEM1	0.95	0.99	0.90	0.72	
SE	ENSGALG0000000533	SRSF3	0.36	0.38	0.30	0.16	
SE	ENSGALG0000023199	HNRNPDL	0.39	0.40	0.30	0.18	
SE	ENSGALG0000006157	DDX26B	0.67	0.60	0.57	0.84	
SE	ENSGALG0000001745	PSTPIP2	0.07	0.05	0.11	0.26	
SE	ENSG00000175029 ^a	CTBP2	0.38	0.38	0.23	0.12	
A5SS	ENSGALG0000008038	SF3B1	0.41	0.57	0.55	0.31	
A5SS	ENSGALG0000002487	SFSWAP	0.58	0.73	0.55	0.41	
A3SS	ENSGALG0000005162	RNPC3	0.55	0.36	0.42	0.67	
A3SS	ENSGALG0000000720	LOC419563	0.88	0.94	0.85	0.70	
A3SS	ENSGALG0000009421	SRSF5	0.55	0.72	0.54	0.39	
A3SS	ENSGALG00000014915	THOC1	0.33	0.48	0.33	0.23	
A3SS	ENSGALG0000000189	YTHDC2	0.44	0.59	0.42	0.32	

SE = Skipped exon; A5SS = alternative 5' splice site; A3SS = alternative 3' splice site. ^a Human homolog.

Туре	Ensembl ID	Symbol	Line 6	Line 6		Line 7	
			control	infected	control	infected	
SE	ENSGALG00000001107	GOSR2	0.73	0.92	0.38	0.59	
SE	ENSG00000124193ª	SRSF6	0.43	0.71	0.54	0.34	
A3SS	ENSGALG0000026498	unknown	0.12	0.70	0.10	0.34	

Table 24. Group IV: Genes showing differential exon usage (ψ) values between the 2 chicken lines and between uninfected and MDV-infected birds

^a Human homolog.

In group II (table 22), ψ values were relatively stable in control and infected chickens within line, but not between lines. Genes that could play an important role in immune responses are RAC3, ITGB2 and HCK. RAC3 (Ras-related C3 botulinum toxin substrate 3) encodes small GTPases, belonging to the Ras family, that regulate a wide variety of cellular events including cell growth, cytoskeletal reorganization, and the activation of protein kinases. HCK transmits signals from cell surface receptors such as FCGR1A, FCGR2A, IL2, IL6, IL18, and integrins (ITGB1, ITGB2). *ITGB2* (*CD18*) encodes subunit β_2 integrin of LFA-1 and CR3 receptors. LFA-1 plays an important role in adhesion of lymphocytes with other cells. CR3 binds to a vast array of ligands and molecules including complement C3bi, microbial proteins, ICAM-1 and -2, ECM proteins, and coagulation proteins. It plays a significant role in neutrophil and monocyte activation including phagocytosis, adhesion and migration [Ehlers, 2000]. DYNLL2, SEPT11 and PFN2 are also involved in cell rearrangement and cytokinesis. In particular, DYNLL2 are dynein proteins that have been demonstrated to regulate T cell activation by driving T-cell receptor microclusters toward the center of an immune synapse [Hashimoto-Tane et al., 2011].

Group III (table 23) includes genes that exhibit differential isoform expression only in response to infection in the susceptible line. A number of genes in this group encode proteins that are parts of the spliceosome: SRSF3, HNRNPDL, SFSWAP, THOC1, RNPC3, and SRSF5.

The last group (group IV, table 24) only has 3 genes: GOSR2, SRSF6 and ENSGALG00000026498. The ψ value differences of these genes were greater than 0.20 the cutoff between control and infected chickens in the resistant and susceptible line and were significantly different between infected chickens in the resistant and susceptible lines. GOSR2 encodes a trafficking membrane

protein important for transporting proteins from cis- to trans-golgi network and SRSF6 (serine/arginine-rich splicing factor 6) encodes a protein involved in mRNA splicing.

Roles of LFA-1 and Actin Cytoskeleton in T Cell Activation

By grouping genes based on patterns of ψ values, we found that many genes in group II (ITGB2, PFN2, DYNLL2, SEPT11, and RAC3) are involved in cytokinesis or cell synapse, which are important for T cell activation. As described above, *ITGB2* encodes the β -subunit of integrins including LFA-1, which is exclusively expressed in lymphocytes and plays a major role in lymphoproliferation, antigen presentation, T cell activation, and cytotoxicity. Integrins are a special kind of receptors that transmit signals bidirectionally across the cell membrane. They are heterodimeric composed of α (large) and β (small) subunits [Wang H et al., 2010]. The β (CD18) subunit encoded by ITGB2 is expressed on lymphocytes and antigen-presenting cells (APCs) as a component of LFA-1 and CR3 receptors. LFA-1 binds to its ligand ICAM-1 to help form a synapse that brings APCs and T cells together to initiate antigen presentation leading to T cell activation [Dustin and Cooper, 2000].

Absence of LFA-1 leads to impaired functions of lymphocytes in proliferation and tumor rejection [Schmits et al., 1996; Scharffetter-Kochanek et al., 1998]. Mutations in the ITGB2 gene have been associated with type 1 leukocyte adhesion deficiency (LAD-1), a human autosomal-recessive inherited disease. The disease is characterized by impairment of lymphocytes in adherent-dependent functions, lack of accumulation to the site of infection and recurrent bacterial and fungal infection [Springer et al., 1987]. In addition, the response of lymphocytes to mitogens is decreased in patients with LAD. The decrease in responsiveness to mitogens has been shown to correlate with resistance to MD by Lee and Bacon [1983], who illustrated that resistant birds (MD-resistant lines 6 and N) were less responsive to phytohemagglutinin than MDsusceptible birds (line 7 and P).

The actin cytoskeleton is very important in T cell activation because it enhances the activity of LFA-1 by increasing its avidity and recruiting signaling molecules necessary for downstream signaling [Dustin and Cooper, 2000; van Kooyk and Figdor, 2000]. Cytoskeleton proteins binding to the cytoplasmic domain of LFA-1 are thought to play an important role in driving LFA-1 to aggregate on the cell surface, resulting in increased avidity. Aggregation of LFA-1 has been demonstrated to be essential for lymphocytes to bind to the ligand [van Kooyk et al., 1994]. Interestingly, RAC3 and PFN2, which are involved in the actin cytoskeleton pathway, also expressed different ratios of alternative splice forms between lines. Some of these gene products are also found in other pathways that are involved in immune responses. It could be speculated that pre-mRNA splicing of these genes is coregulated by splicing regulators or some genetic factors.

Prediction of Functional Domains of Splice Forms of Genes in the Actin Cytoskeleton Pathway

To predict the function of the alternative splice forms of genes in the actin cytoskeleton pathway, transcript sequences were translated to protein sequences by ESTscan [Iseli et al., 1999]. Protein sequences were then searched for annotated protein domains using InterPro Scan [Quevillon et al., 2005]. Besides ITGB2, other genes have alternative exons located in coding regions that could potentially affect functional protein domains in some ways. The exon with an alternative 3' splice site of RAC3 encodes part of a protein domain identified as a small GTPase of the Ras subfamily (ProSiteProfiles:PS5142 and SMART:SM00173). Rac3 is highly homologous to Rac1 and has been reported to possess the ability to promote membrane ruffling, transformation, activation of c-Jun transcriptional activity, and co-activation of NFkB [Werbajh et al., 2000]. Activated Rac also regulates production of superoxide in neutrophils and macrophages.

The alternative exon of *PFN2* seems to disrupt the coding sequence that encodes the profilin domain (Pfam: PF00235). The profilin domain is essential for almost all organisms, and its functions include regulating actin polymerization, controlling complex networks of molecular interaction and transmitting signals from small-GTPase pathways. It also binds to Rac effector molecules and a number of other ligands [Witke, 2004]. Even

though the exact mechanism is not known, lack of responsiveness of T cells to stimuli appears to benefit resistant birds because in these birds, MDV cannot induce T cells to proliferate and cause them to undergo neoplastic transformation. It has also been suggested that the mechanism that controls both lymphocyte proliferation induced by MDV and lymphocyte proliferation induced by the immune response is the same [Pazderka et al., 1975]. Therefore, it may be useful to consider a link between the deficiency of lymphocytes in the resistant line and the alternative splice form of *ITGB2* that is only expressed in the resistant line. Although the exon included in the alternative splice form is noncoding, it could serve important functions in translation or posttranscriptional regulation.

Prediction of cis-Regulatory Elements in Alternative Splicing Exons of Genes in Group II

Among all groups, alternative splicing of genes in the group II is most likely to be regulated by genetic factors because the ratios of isoform expression in this group were relatively stable within line, but were significantly different between lines. Investigation of nucleotide differences within exons of both lines could reveal a possible role of SNPs in regulating alternative splicing in this group. We obtained a sequence of alternative exons from the resistant line and used Human Splicing Finder to determine whether SNPs from the susceptible line could alter predicted ESEs or ESSs. Results from some genes involved in cytokinesis are discussed below.

For ITGB2, a SNP at position 26 of the cDNA from the resistant line located at position 7,183,696 on chromosome 7 is in a predicted binding site for SC35, which is an exon enhancer. Although the exon is not expressed in the susceptible line, we found that there are no polymorphisms between lines based on genome resequencing. Therefore, this SNP may not account for exclusion of the exon in the susceptible line (fig. 20). Exon sequences of PFN2 from the lines 6 and 7 differ at position 23,221,934 on chromosome 9. A small insertion of 2 AA nucleotides is predicted to create a new binding site for Tra2-β splicing regulator (table 25), which serves as a stabilizer of an enhancer complex [Lopez, 1998]. From exon expression data, ψ of an exon with alternative splicing increases from 0.20-0.30 to about 0.50 in the susceptible line. Tra2 could possibly increase inclusion of the exon with alternative 3' splice site via ESE-dependent 3' splice site activation.

Replacement of an A with a G nucleotide in the skipped exon of *DYNLL2* from line 6 is predicted to slightly alter the binding site of several ESEs as well as to create a new



Fig. 20. An SNP in an *ITGB2* exon can explain alternate splicing. A single-nucleotide substitution (T>C) at position 7,183,696 of an *ITGB2* exon is predicted to disrupt a putative ESE motif. The numbers not on the axes in the figure on the left side reflect the actual counts for each exon splice junction read for each sample, and the ψ values given on the right side are the percent of variants containing the potentially skipped exon.

binding site for 9G8 (table 25). This exon is upregulated in the susceptible line compared to the resistant line, therefore, the presence of the new binding site for 9G8 exon enhancer helps support the expression results. In addition, the G nucleotide in this position matches the reference nucleotide; therefore, we could expect this exon to be expressed in other datasets. According to EST tags on the UCSC genome browser, the exon has been found and sequenced from chicken eyes (EST sequence: DR424100).

Conclusions

Custom gene models built by combining gene models from de novo assembly, reference-based assembly, and Ensembl have allowed us to identify genes and isoforms that might play an important role in resistance to MD. Results from gene expression analysis indicated that adaptive immune responses were more highly activated during lytic infection in the susceptible line than in the resistant line. Because only activated T cells are thought to be infected by MDV, we speculate that enhancement of adaptive immune responses could help spread the viruses by recruiting and activating more T cells. In contrast, the delay or reduction of adaptive immune responses could benefit the host by limiting infection of activated T cells.

To elucidate the molecular mechanism of MD resistance, we investigated differential isoform expression between lines and identified a number of genes that could be responsible for differences in immune responses. Notably, this includes several genes involved in actin cyto-

Gene	cDNA position	Linked SR protein	Туре	Reference motif	Mutant motif	Variation
DYNLL2	2 2 4 4 6 6	SF2/ASF (IgM-BRCA1) SF2/ASF, SF2/ASF (IgM-BRCA1) SF2/ASF (IgM-BRCA1) SF2/ASF, SF2/ASF (IgM-BRCA1) 9G8 hnRNP A1	ESE ^a ESE ^a ESE ^a ESE ^b ESE ^d	CTCCGGG (86.38) CTCCGGG (76.91) CCGGGGT (73.00) CCGGGGT (73.00)	CTCCGAG (72.69) CTCCGAG (72.69) CCGAGGT (86.23) CCGAGGT (82.94) GAGGTG (60.67) GAGGTG (74.05)	new site new site
PFN2	2068 2069 2070 2066 2067	Tra2-β Tra2-β Tra2-β	ESE ^a ESE ^a ESS ^c ESS ^c	AAAAT (81.02)	AAAAa AAAaaa (94.14) AAAaaT (81.02) ACAAAAaa (38.13) CAAAAaaT (28.85)	+16.19% new site new site new site new site

Table 25. Selected group II genes (see table 22) showing differential exon usage (ψ) between lines 6 and 7 with potential sequence variants in motifs that may account for the alternative splicing

^a ESE Finder matrices for SRp40, SC35, SF2/ASF and SRp55 proteins.

^b ESE motifs from Human Splicing Finder.

^c Predicted PESS octamers from Zhang and Chasin [2004].

^d hnRNP motif.

skeleton structure and cytokinesis, which are important for the functions of lymphocytes and immune cells but have not been of great interest in the field of MD research. Even though we mainly discuss the possible role of *ITGB2* in MD resistance, other genes cannot be precluded and should be candidates for further investigation and experimental studies. Moreover, the full mechanism of MD resistance is highly complex and more data from different stages of infection as well as greater sequencing depth will be required to identify all genes and isoforms involved. To enhance the study of unannotated gene and isoform expression, our approach of constructing gene models from RNA-seq should be iteratively used to extend the Ensembl data to construct more complete gene models.

Materials and Methods

Detailed computational analysis and methods are available at https://github.com/likit/mdv_rnaseq_paper.

The National Avian Research Facility

(Prepared by A. Hart, R. Kuo, L. Eöry, P. Kaiser, and D.W. Burt)

The National Avian Research Facility (NARF) was created as a partnership between The Roslin Institute and The Pirbright Institute to provide a wide range of resources for the avian research community including access to

Third Report on Chicken Genes and Chromosomes 2015

avian lines, biological tools and genomic databases. We have received capital support from The Roslin Foundation, BBSRC, University of Edinburgh and The Roslin Institute as well as funding from the Wellcome Trust.

NARF is currently comprised of avian rearing facilities and a large group of collaborators contributing to the services that we aim to provide for the community. There are 2 avian units, The Greenwood Building, a conventional avian facility, and The Bumstead Building, a National Avian Specific Pathogen Free (SPF) Research Facility. The purpose of the Bumstead Building is to maintain unique inbred genetic lines of poultry currently housed at The Pirbright Institute Compton Laboratories. Both facilities will support research programmes currently running at both institutes, and NARF will also offer UK and international investigators access to these unique bioresources.

The Greenwood Building is named after Dr Alan Greenwood. Alan Greenwood moved to Edinburgh in the mid 1920's from Australia to study with Dr Francis Albert Crew at the Institute of Animal Genetics, The University of Edinburgh. In the late 1920's, Alan Greenwood started an inbreeding programme with Brown Leghorn chickens to study a variety of traits, egg laying, plumage, vigour, etc. Through this programme he generated 9 lines denoted A through to I. During 1947 Dr Greenwood founded and was Director of the Edinburgh Poultry Research Centre (PRC); he continued in this role until his retirement in the 1960's. His inbred chicken

lines were intercrossed at this time to generate what is known as the J-line where this line continues to be studied at NARF.

The Bumstead Building is named after Dr Nat Bumstead, former Head of Avian Genetics Group, BBSRC Institute for Animal Health. Nat Bumstead was a leading avian geneticist who published extensively on the genetics of immunity and disease resistance and susceptibility using chicken as the model organism. He was instrumental in establishing worldwide research on the genome of the chicken. The inbred chicken lines that Dr Bumstead studied throughout his research career continue to be studied today and will move to NARF from late 2014. These birds will be kept in SPF conditions, and NARF will supply these to researchers to gain greater insights into gene function in relation to disease susceptibility and resistance.

Considerable investment by BBSRC and other governmental departments over several years has supported several of these chicken lines, with some tracing their origins to the 1930's. All of the lines, including those that are currently housed at The Roslin Institute and The Pirbright Institute offer a unique resource backed with a wealth of data. From late 2014, NARF will become fully operational taking over the role as curator for these unique chicken lines.

NARF has also been working on creating an avian biobank. The advantages of developing an avian biobank are many:

- Cryopreserve rare or endangered breeds of chicken to maintain genetic diversity for future generations;
- In the event of disease outbreak, flocks could be rederived from frozen material;
- The cost of maintaining flocks increases year-on-year, and therefore, if we were able to cryopreserve lines using primordial germ cells and/or semen, then this could free up space in avian facilities for those lines that are being actively researched.

Robust procedures that would allow us to freeze, maintain and rederive lines in the future would provide confidence that unique genetic resources will not be lost if the live flock is no longer 'on the shelf'.

In addition to providing various animal resources, NARF collaborators are developing biological tools such as antibodies, primers and reagents. Information regarding available chicken lines and biological tools are on the NARF website (www.narf.ac.uk). The NARF website also provides links to relevant resources as well as our own contributions to the annotation of avian genomes. At present we are developing several informaticsbased resources including a variation database, alternative splice variant annotation, and avian comparative genomics annotations.

The variation database will contain single nucleotide polymorphisms (SNPs) and short insertions/deletions (InDels) detected in the inbred lines at NARF. Where appropriate, specific SNPs and InDels will be linked to relevant annotations and biological tools to provide a more information-rich experience.

The alternative splice variant annotation will contain full-length transcript sequences obtained using long-read sequencing technology. This information will greatly expand our current knowledge of isoforms occurring in chicken as well as expanding the list of putative noncoding RNA. Using this annotation, researchers will be able to run RNA-seq analyses on the isoform level.

To facilitate genetics and genomics research, NARF has been making considerable efforts to improve and share the genome annotations currently available for poultry genomes. Having good quality, functional annotations are crucial for the understanding of the link between genetic variations and phenotypic differences between individuals. Although annotations from largescale functional assays are at present only available for mammals (see e.g. the human ENCODE project [EN-CODE Project Consortium, 2012]), annotations can still be improved by incorporating information on selective forces from comparative genome analyses of recently sequenced bird genomes [Zhang G et al., 2014b]. For example, regions in the chicken genome that are known to be under purifying selection can serve as an extra layer of annotation as these regions can be predictive for function [Lindblad-Toh et al., 2011].

While the primary focus at NARF is on poultry genomics, integrating and sharing genome sequences and annotations for bird species in general can also benefit the diverse bird research communities. Therefore, we provided access to our integrated genome resource through a web portal called Avianbase (http://avianbase.narf. ac.uk) [Eöry et al., 2015], an Ensembl-based [Flicek et al., 2014] genome browser instance, which provides visual displays of the genome sequences, gene models, transcripts, and translations along with the BLAST search facility for 44 newly sequenced bird genomes [Zhang G et al., 2014b].

The goal of NARF is to become a hub for the avian research community. We are working to provide new resources to help researchers find the information and tools they need to progress.

References

- Abasht B, Lamont SJ: Genome-Wide Association Study of Fatness in Chickens. Animal Industry Report: AS 653, ASL R2218 2007 (2007). http://lib.dr.iastate.edu/ans_air/vol653/ iss1/44
- Abbott UK, Asmundson VS: Scaleless, an inherited ectodermal defect in the domestic fowl. J Hered 48:63–70 (1957).
- Abe H, Nagao K, Inoue-Murayama M: Short copy number variations potentially associated with tonic immobility responses in newly hatched chicks. PLoS One 8:e80205 (2013).
- Abernathy J, Li X, Jia X, Chou W, Lamont SJ, et al: Copy number variation in Fayoumi and leghorn chickens analyzed using array comparative genomic hybridization. Anim Genet 45:400–411 (2014).
- Acharya S, Foletta VC, Lee JW, Rayborn ME, Rodriguez IR, et al: SPACRCAN, a novel human interphotoreceptor matrix hyaluronan-binding proteoglycan synthesized by photoreceptors and pinealocytes. J Biol Chem 275:6945– 6955 (2000).
- Addinger HK, Calnek BW: Pathogenesis of Marek's disease: early distribution of virus and viral antigens in infected chickens. J Natl Cancer Inst 50:1287–1298 (1973).
- Adli M, Bernstein BE: Whole-genome chromatin profiling from limited numbers of cells using nano-ChIP-seq. Nat Protoc 6:1656–1668 (2011).
- Aerts J, Megens HJ, Veenendaal T, Ovcharenko I, Crooijmans R, et al: Extent of linkage disequilibrium in chicken. Cytogenet Genome Res 117:338–345 (2007).
- Albers CA, Lunter G, MacArthur DG, McVean G, Ouwehand WH, Durbin R: Dindel: accurate indel calls from short-read data. Genome Res 21:961–973 (2011).
- Ali M, Hocking PM, McKibbin M, Finnegan S, Shires M, et al: *Mpdz* null allele in an avian model of retinal degeneration and mutations in human leber congenital amaurosis and retinitis pigmentosa. Invest Ophthalmol Vis Sci 52:7432–7440 (2011).
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ: Basic local alignment search tool. J Mol Biol 215:403–410 (1990).
- Amrein H, Maniatis T, Nöthiger R: Alternatively spliced transcripts of the sex-determining gene *tra-2* of *Drosophila* encode functional proteins of different size. EMBO J 9:3619– 3629 (1990).
- Andreescu C, Avendano S, Brown SR, Hassen A, Lamont SJ, Dekkers JCM: Linkage disequilibrium in related breeding lines of chickens. Genetics 177:2161–2169 (2007).
- Arakawa H, Buerstedde JM: Activation-induced cytidine deaminase-mediated hypermutation in the DT40 cell line. Philos Trans R Soc Lond B Biol Sci 364:639–644 (2009).
- Arnold AP: The end of gonad-centric sex determination in mammals. Trends Genet 28:55– 61 (2012).

- Arnold AP, Chen X, Itoh Y: What a difference an X or Y makes: sex chromosomes, gene dose, and epigenetics in sexual differentiation. Handb Exp Pharmacol 214:67–88 (2012).
- Arnold AP, Chen X, Link JC, Itoh Y, Reue K: Cellautonomous sex determination outside of the gonad. Dev Dyn 242:371–379 (2013).
- Arumugaswami V, Kumar PM, Konjufca V, Dienglewicz RL, Reddy SM, Parcells MS: Latency of Marek's disease virus (MDV) in a reticuloendotheliosis virus-transformed T-cell line. II: Expression of the latent MDV genome. Avian Dis 53:156–165 (2009).
- Asagoshi K, Tano K, Chastain PD 2nd, Adachi N, Sonoda E, et al: FEN1 functions in long patch base excision repair under conditions of oxidative stress in vertebrate cells. Mol Cancer Res 8:204–215 (2010).
- Atkins KE, Read AF, Savill NJ, Renz KG, Islam AF, et al: Vaccination and reduced cohort duration can drive virulence evolution: Marek's disease virus and industrialized agriculture. Evolution 67:851–860 (2013).
- Aubert G, Hills M, Lansdorp PM: Telomere length measurement – caveats and a critical assessment of the available technologies and tools. Mutat Res 730:59–67 (2012).
- Auguste A, Chassot AA, Grégoire EP, Renault L, Pannetier M, et al: Loss of *R-Spondin1* and *Foxl2* amplifies female-to-male sex reversal in XX mice. Sex Dev 5:304–317 (2011).
- Avendaño S, Watson K, Kranis A: Genomics in poultry breeding: from Utopia to deliverables, in: Proceedings of the 9th World Congress on Genetics Applied to Livestock Production, August 1–6, 2010, Leipzig, Germany (2010).
- Ayers KL, Davidson NM, Demiyah D, Roeszler KN, Grützner F, et al: RNA sequencing reveals sexually dimorphic gene expression before gonadal differentiation in chicken and allows comprehensive annotation of the Wchromosome. Genome Biol 14:R26 (2013a).
- Ayers KL, Sinclair AH, Smith CA: The molecular genetics of ovarian differentiation in the avian model. Sex Dev 7:80–94 (2013b).
- Ayers KL, Smith CA, Lambeth LS: The molecular genetics of avian sex determination and its manipulation. Genesis 51:325–336 (2013c).
- Baba TW, Giroir BP, Humphries EH: Cell lines derived from avian lymphomas exhibit two distinct phenotypes. Virology 144:139–151 (1985).
- Baigent SJ, Davison TF: Development and composition of lymphoid lesions in the spleens of Marek's disease virus-infected chickens: association with virus spread and the pathogenesis of Marek's disease. Avian Pathol 28:287– 300 (1999).
- Baigent SJ, Davison F: Marek's disease virus: biology and life cycle, in Davison F, Nair V (eds): Marek's Disease, an Evolving Problem, pp 62–77 (Elsevier Academic Press, London 2004).

- Baigent SJ, Petherbridge LJ, Smith LP, Zhao Y, Chesters PM, Nair VK: Herpesvirus of turkey reconstituted from bacterial artificial chromosome clones induces protection against Marek's disease. J Gen Virol 87:769–776 (2006).
- Baker AJ, Pereira SL: Ratites and tinamous (Paleognathae), in Hedges SB, Kumar S (eds): The Timetree of Life, pp 412–414 (Oxford University Press, New York 2009).
- Baker BS, Burtis K, Goralski T, Mattox W, Nagoshi R: Molecular genetic aspects of sex determination in *Drosophila melanogaster*. Genome 31:638–645 (1989).
- Barbin A, Montpellier C, Kokalj-Vocak N, Gibaud A, Niveleau A, et al: New sites of methylcytosine-rich DNA detected on metaphase chromosomes. Hum Genet 94:684–692 (1994).
- Barker FK, Barrowclough GF, Groth JG: A phylogenetic hypothesis for passerine birds: taxonomic and biogeographic implications of an analysis of nuclear DNA sequence data. Proc Biol Sci 269:295–308 (2002).
- Barrangou R, Fremaux C, Deveau H, Richards M, Boyaval P, et al: CRISPR provides acquired resistance against viruses in prokaryotes. Science 315:1709–1712 (2007).
- Bartlett JR: Linkage analysis of endogenous viral element 1, blue eggshell, and pea comb loci in chickens. J Hered 87:67–70 (1996).
- Bateson W: Experiments with poultry. Rep Evol Comm Roy Soc 187–124 (1902).
- Bateson W, Punnet R: Experimental studies in the physiology of heredity. Poultry Repts Evol Comm Roy Soc 418–35 (1908).
- Bauer MM, Reed KM: Extended sequence of the turkey MHC B-locus and sequence variation in the highly polymorphic B-G loci. Immunogenetics 63:209–221 (2011).
- Baumjohann D, Ansel KM: MicroRNA regulation of the germinal center response. Curr Opin Immunol 28:6–11 (2014).
- Begun DJ, Aquadro CF: Levels of naturally occurring DNA polymorphism correlate with recombination rates in *D. melanogaster*. Nature 356:519–520 (1992).
- Bell O, Tiwari VK, Thoma NH, Schübeler D: Determinants and dynamics of genome accessibility. Nat Rev Genet 12:554–564 (2011).
- Bellott DW, Skaletsky H, Pyntikova T, Mardis ER, Graves T, et al: Convergent evolution of chicken Z and human X chromosomes by expansion and gene acquisition. Nature 466: 612–616 (2010).
- Bensaada M, Kiefer H, Tachdjian G, Lapierre JM, Cacheux V, et al: Altered patterns of DNA methylation on chromosomes from leukemia cell lines: identification of 5-methylcytosines by indirect immunodetection. Cancer Genet Cytogenet 103:101–109 (1998).
- Berger SL, Kouzarides T, Shiekhattar R, Shilatifard A: An operational definition of epigenetics. Genes Dev 23:781–783 (2009).

- Beridze T: Satellite DNA (Springer, Berlin 1986).
- Berlin S, Ellegren H: Chicken W: a genetically uniform chromosome in a highly variable genome. Proc Natl Acad Sci USA 101:15967– 15969 (2004).
- Bernardino J, Lamoliatte E, Lombard M, Niveleau A, Malfoy N, et al: DNA methylation of the X chromosomes of the human female: an in situ semi-quantitative analysis. Chromosoma 104:528–535 (1996).
- Bernardino J, Lombard M, Niveleau A, Dutrillaux B: Common methylation characteristics of sex chromosomes in somatic and germ cells from mouse, lemur and human. Chromosome Res 8:513–525 (2000).
- Bernstein BE, Meissner A, Lander ES: The mammalian epigenome. Cell 128:669–681 (2007).
- Bernstein BE, Stamatoyannopoulos JA, Costello JF, Ren B, Milosavljevic A, et al: The NIH Roadmap Epigenomics Mapping Consortium. Nat Biotechnol 28:1045–1048 (2010).
- Berthouly C, Leroy G, Van TN, Thanh HH, Bed'Hom B, et al: Genetic analysis of local Vietnamese chickens provides evidence of gene flow from wild to domestic populations. BMC Genet 10:1 (2009).
- Bett RC, Bhuiyan AK, Khan MS, Silva GL, Thuy LT, et al: Phenotypic variation of native chicken populations in the South and South East Asia. Int J Poultry Sci 13:449–460 (2014).
- Beyer C, Schett G, Distler O, Distler JHW: Animal models of systemic sclerosis: prospects and limitations. Arthritis Rheum 62:2831–2844 (2010).
- Bird A: DNA methylation patterns and epigenetic memory. Genes Dev 16:6–21 (2002).
- Bitgood JJ, Somes RG: Linkage relationships and gene mapping, in Crawford RD (ed): Poultry Breeding and Genetics, pp 469–496 (Elsevier Publications, Amsterdam 1990).
- Bitgood JJ, Shoffner RN, Otis JS, Briles WE: Mapping of the genes for pea comb, blue egg, barring, silver, and blood groups A, E, H, and P in the domestic fowl. Poult Sci 59:1686–1693 (1980).
- Blagoveshchenskiĭ IIu, Sazanova AL, Stekol'nikova VA, Fomichev KA, Barkova OIu, et al: Investigation of pseudoautosomal and bordering regions in avian Z and W chromosomes with the use of large insert genomic BAC clones (in Russian). Genetika 47:312– 319 (2011).
- Blencowe BJ: Exonic splicing enhancers: mechanism of action, diversity and role in human genetic diseases. Trends Biochem Sci 25:106– 110 (2000).
- Blyth JSS, Sang JH: Survey of line crosses in a brown leghorn flock. 1. Egg production. Genet Res 1:408–421 (1960).
- Bock C: Analysing and interpreting DNA methylation data. Nat Rev Genet 13:705–719 (2012).
- Bonasio R, Tu S, Reinberg D: Molecular signals of epigenetic states. Science 330:612–616 (2010).

- Booth MJ, Branco MR, Ficz G, Oxley D, Krueger F, et al: Quantitative sequencing of 5-methylcytosine and 5-hydroxymethylcytosine at single-base resolution. Science 336:934–937 (2012).
- Boschiero C, Gheyas AA, Ralph HK, Eory L, Paton B, et al: Detection and characterization of small insertions and deletions in modern layer chicken genomes. BMC Genomics (in press).
- Bovine Genome Sequencing and Analysis Consortium, Elsik CG, Tellam RL, Worley KC, Gibbs RA, et al: The genome sequence of taurine cattle: a window to ruminant biology and evolution. Science 324:522–528 (2009).
- Boyle AP, Davis S, Shulha HP, Meltzer P, Margulies EH, et al: High-resolution mapping and characterization of open chromatin across the genome. Cell 132:311–322 (2008).
- Brandström M, Ellegren H: The genomic landscape of short insertion and deletion polymorphisms in the chicken (*Gallus gallus*) genome: a high frequency of deletions in tandem duplicates. Genetics 176:1691–1701 (2007).
- Breg WR, Schreck RR, Miller OJ: Familial trisomy 15: identification of a deleted No. 15 confirmed by anti-5-methylcytosine antibody binding. Am J Hum Genet 26:17 (1974).
- Brutlag DL: Molecular arrangement and evolution of heterochromatic DNA. Annu Rev Genet 14:121–144 (1980).
- Buenrostro JD, Giresi PG, Zaba LC, Chang HY, Greenleaf WJ: Transposition of native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding proteins and nucleosome position. Nat Methods 10:1213–1218 (2013).
- Buerstedde JM, Takeda S: Reviews and Protocols in DT40 Research. Subcellular Biochemistry, vol 40 (Springer, Netherlands 2006).
- Bumstead N: Genomic mapping of resistance to Marek's disease. Avian Path 27:S78–S81 (1998).
- Burge SW, Daub J, Eberhardt R, Tate J, Barquist L, et al: Rfam 11.0: 10 years of RNA families. Nucleic Acids Res 41:D226–D232 (2013).
- Burt DW: Origin and evolution of avian microchromosomes. Cytogenet Genome Res 96: 97-112 (2002).
- Burt DW: Chicken genome: current status and future opportunities. Genome Res 15:1692– 1698 (2005).
- Burt DW, Bruley C, Dunn IC, Jones CT, Ramage A, et al: The dynamics of chromosome evolution in birds and mammals. Nature 402:411–413 (1999).
- Burt DW, Carre W, Fell M, Law AS, Antin PB, et al: The chicken gene nomenclature committee report. BMC Genomics 10 Suppl 2:S5 (2009).
- Cai Q, Qian X, Lang Y, Luo Y, Xu J, et al: Genome sequence of ground tit *Pseudopodoces humilis* and its adaptation to high altitude. Genome Biol 14:R29 (2013).

- Campo VA, Patenaude AM, Kaden S, Horb L, Firka D, et al: MSH6- or PMS2-deficiency causes re-replication in DT40 B cells, but it has little effect on immunoglobulin gene conversion or on repair of AID-generated uracils. Nucleic Acids Res 41:3032–3046 (2013).
- Cech TR, Steitz JA: The noncoding RNA revolution – trashing old rules to forge new ones. Cell 157:77–94 (2014).
- Chan PP, Lowe TM: GtRNAdb: a database of transfer RNA genes detected in genomic sequence. Nucleic Acids Res 37:D93–D97 (2009).
- Chang CF, Schock EN, O'Hare EA, Dodgson J, Cheng HH, et al: The cellular and molecular etiology of the craniofacial defects in the avian ciliopathic mutant *talpid2*. Development 141: 3003–3012 (2014).
- Chang CM, Coville JL, Coquerelle G, Gourichon D, Oulmouden A, Tixier-Boichard M: Complete association between a retroviral insertion in the tyrosinase gene and the recessive white mutation in chickens. BMC Genomics 7:19 (2006).
- Chang H, Delany ME: Karyotype stability of the DT40 chicken B cell line: macrochromosome variation and cytogenetic mosaicism. Chromosome Res 12:299–307 (2004).
- Chapus C, Edwards SV: Genome evolution in Reptilia: in silico chicken mapping of 12,000 BACend sequences from two reptiles and a basal bird. BMC Genomics 10 Suppl 2:S8 (2009).
- Charlesworth B, Sniegowski P, Stephan W: The evolutionary dynamics of repetitive DNA in eukaryotes. Nature 371:215–220 (1994).
- Chassot AA, Ranc F, Grégoire EP, Roepers-Gajadien HL, Taketo MM, et al: Activation of β-catenin signaling by Rspo1 controls differentiation of the mammalian ovary. Hum Mol Genet 17:1264–1277 (2008).
- Chassot AA, Gillot I, Chaboissier MC: R-spondin1, WNT4, and the CTNNB1 signaling pathway: strict control over ovarian differentiation. Reproduction 148:R97–R110 (2014).
- Chaves LD, Krueth SB, Reed KM: Characterization of the turkey MHC chromosome through genetic and physical mapping. Cytogenet Genome Res 117:213–220 (2007).
- Chen CY, Misztal I, Aguilar I, Tsuruta S, Meuwissen THE, et al: Genome-wide marker-assisted selection combining all pedigree phenotypic information with genotypic data in one step: an example using broiler chickens. J Anim Sci 89:23–28 (2011).
- Chen N, Bellott DW, Page DC, Clark AG: Identification of avian W-linked contigs by short-read sequencing. BMC Genomics 13:183 (2012).
- Chiappe LM, Dyke GJ: The Mesozoic radiation of birds. Annu Rev Ecol Syst 33:91–124 (2002).
- Chodroff RA, Goodstadt L, Sirey TM, Oliver PL, Davies KE, et al: Long noncoding RNA genes: conservation of sequence and brain expression among diverse amniotes. Genome Biol 11:R72 (2010).
- Choi Y, Sims GE, Murphy S, Miller JR, Chan AP: Predicting the functional effect of amino acid substitutions and indels. PLoS One 7:e46688 (2012).

- Christian M, Cermak T, Doyle EL, Schmidt C, Zhang F, et al: Targeting DNA double-strand breaks with TAL effector nucleases. Genetics 186:757–761 (2010).
- Christidis L: Aves, in John B, Kayano H, Levan A (eds): Animal Cytogenetics, vol. 4: Chordata 3 B (Gebrüder Borntraeger, Berlin 1990).
- Chue J, Smith CA: Sex determination and sexual differentiation in the avian model. FEBS J 278: 1027–1034 (2011).
- Clayton DF, Balakrishnan CN, London SE: Integrating genomes, brain and behavior in the study of songbirds. Curr Biol 19:R865–R873 (2009).
- Clurman BE, Hayward WS: Multiple proto-oncogene activations in avian leukosis virus-induced lymphomas: evidence for stage-specific events. Mol Cell Biol 9:2657–2664 (1989).
- Clynes D, Gibbons RJ: ATRX and the replication of structured DNA. Curr Opin Genet Dev 23: 289–294 (2013).
- Coble DJ, Fleming D, Persia ME, Ashwell CM, Rothschild MF, et al: RNA-seq analysis of broiler liver transcriptome reveals novel responses to high ambient temperature. BMC Genomics 15:1084 (2014).
- Cole RK: Hereditary hypothyroidism in the domestic fowl. Genetics 53:1021–1033 (1966).
- Cooley C, Baird KM, Faure V, Wenner T, Stewart JL, et al: Trf1 is not required for proliferation or functional telomere maintenance in chicken DT40 cells. Mol Biol Cell 20:2563–2571 (2009).
- Cooper A, Penny D: Mass survival of birds across the Cretaceous-Tertiary boundary: molecular evidence. Science 275:1109–1113 (1997).
- Cooper A, Lalueza-Fox C, Anderson S, Rambaut A, Austin J, Ward R: Complete mitochondrial genome sequences of two extinct moas clarify ratite evolution. Nature 409:704–707 (2001).
- Cooper GM, Shendure J: Needles in stacks of needles: finding disease-causal variants in a wealth of genomic data. Nat Rev Genet 12: 628–640 (2011).
- Cracraft J: Continental drift, paleoclimatology, and the evolution and biogeography of birds. J Zool 169:455–545 (1973).
- Cracraft J: Avian evolution, Gondwana biogeography and the Cretaceous-Tertiary mass extinction event. Proc Biol Sci 268:459–469 (2001).
- Cracraft J, Barker FK, Braun M, Harshman J, Dyke GJ, et al: Phylogenetic relationships among modern birds (Neornithes), in Cracraft J, Donoghue MJ (eds): Assembling the Tree of Life, pp 468–478 (Oxford University Press, Oxford 2004).
- Craven AJ, Ormandy CJ, Robertson FG, Wilkins RJ, Kelly PA, et al: Prolactin signaling influences the timing mechanism of the hair follicle: analysis of hair growth cycles in prolactin receptor knockout mice. Endocrinology 142: 2533–2539 (2001).
- Crawford RD: Poultry biology, in Crawford RD (ed): Poultry Breeding and Genetics, pp 1–108 (Elsevier, Amsterdam 1990).

- Crooijmans RPMA, Vrebalov J, Dijkhof RJM, van der Poel JJ, Groenen MAM: Two-dimensional screening of the Wageningen chicken BAC library. Mamm Genome 11:360–363 (2000).
- Crooijmans RPMA, Fife MS, Fitzgerald TW, Strickland S, Cheng HH, et al: Large scale variation in DNA copy number in chicken breeds. BMC Genomics 14:398 (2013).
- Cui J, Zhao W, Huang Z, Jarvis ED, Gilbert MT, et al: Low frequency of paleoviral infiltration across the avian phylogeny. Genome Biol 15: 539 (2014).
- Cui K, Zhao K: Genome-wide approaches to determining nucleosome occupancy in metazoans using MNase-Seq. Methods Mol Biol 833: 413–419 (2012).
- Cunningham F, Amode MR, Barrell D, Beal K, Billis K, et al: Ensembl 2015. Nucleic Acids Res 43:D662–D669 (2015).
- Dalloul RA, Long JA, Zimin AV, Aslam L, Beal K, et al: Multi-platform next-generation sequencing of the domestic turkey (*Meleagris gallopavo*): genome assembly and analysis. PLoS Biol 8:e1000475 (2010).
- Dana N, van der Waaij LH, Dessie T, van Arendonk JAM: Production objectives and trait preferences of village poultry producers of Ethiopia: implications for designing breeding schemes utilizing indigenous chicken genetic resources. Trop Anim Health Prod 42:1519– 1529 (2010).
- Dana N, Megens HJ, Crooijmans RPMA, Hanotte O, Mwacharo J, et al: East Asian contributions to Dutch traditional and western commercial chickens inferred from mtDNA analysis. Anim Genet 42:125–133 (2011).
- Daniels LM, Delany ME: Molecular and cytogenetic organization of the 5S ribosomal DNA array in chicken (*Gallus gallus*). Chromosome Res 11:305–317 (2003).
- Davey MG, Paton IR, Yin Y, Schmidt M, Bangs FK, et al: The chicken *talpid*³ gene encodes a novel protein essential for hedgehog signaling. Genes Dev 20:1365–1377 (2006).
- Dekker J, Rippe K, Dekker M, Kleckner N: Capturing chromosome conformation. Science 295:1306–1311 (2002).
- de Koning DJ, Nadaf J, Liu X, Li X, Le Bihan-Duval E, Dunn I: Integrating QTL mapping, gene-expression and next-generation sequencing to unravel a QTL affecting meat quality in chicken. Adv Anim Biosci 3:184– 206 (2012).
- Delacour J: The Pheasants of the World, ed 2 (Spur Publications and World Pheasant Association, 1977).
- Delany ME: Genetic variants for chick biology research: from breeds to mutants. Mech Dev 121:1169–1177 (2004).
- Delany ME, Gessaro TM: Genetic stocks for immunological research, in Davison F, Kaspers B, Schat KA (eds): Avian Immunology, pp 443–458 (Academic Press, Elsevier, London 2008).

- Delany ME, Gessaro TM, Rodrigue KL, Daniels LM: Chromosomal mapping of chicken mega-telomere arrays to GGA9, 16, 28 and W using a cytogenomic approach. Cytogenet Genome Res 117:54–63 (2007).
- Delecluse HJ, Hammerschmidt W: Status of Marek's disease virus in established lymphoma cell lines: herpesvirus integration is common. J Virol 67:82–92 (1993).
- Delecluse HJ, Schüller S, Hammerschmidt W: Latent Marek's disease virus can be activated from its chromosomally integrated state in herpesvirus-transformed lymphoma cells. EMBO J 12:3277–3286 (1993).
- de Medeiros RB: Sequential chromatin immunoprecipitation assay and analysis. Methods Mol Biol 791:225–237 (2011).
- Derrien T, Johnson R, Bussotti G, Tanzer A, Djebali S, et al: The GENCODE v7 catalog of human long noncoding RNAs: analysis of their gene structure, evolution, and expression. Genome Res 22:1775–1789 (2012).
- Dessie T, Taye T, Dana N, Ayalew W, Hanotte O: Current state of knowledge on phenotypic characteristics of indigenous chickens in the tropics. World Poult Sci J 67:507–516 (2011).
- Desta TT: Phenomic and Genomic Landscape of Ethiopian Village Chickens. PhD thesis, The University of Nottingham (2015).
- Desta TT, Wakeyo O: Uses and flock management practices of scavenging chickens in Wolaita Zone of southern Ethiopia. Trop Anim Health Prod 44:537–544 (2012).
- Desta TT, Dessie T, Bettridge J, Lynch SE, Melese K, et al: Signature of artificial selection and ecological landscape on morphological structures of Ethiopian village chickens. Anim Genet Res 52:17–29 (2013).
- Dev VG, Warburton D, Miller OJ, Miller DA, Erlanger BF, Beiser SM: Consistent pattern of binding of anti-adenosine antibodies to human metaphase chromosomes. Exp Cell Res 74:288–295 (1972).
- de Wit E, de Laat W: A decade of 3C technologies: insights into nuclear organization. Genes Dev 26:11–24 (2012).
- de Yébenes VG, Bartolome-Izquierdo N, Ramiro AR: Regulation of B-cell development and function by microRNAs. Immunol Rev 253: 25–39 (2013).
- Dienglewicz RL, Parcells MS: Establishment of a lymphoblastoid cell line using a mutant MDV containing a green fluorescent protein expression cassette. Acta Virol 43:106–112 (1999).
- Dorsett Y, McBride KM, Jankovic M, Gazumyan A, Thai TH, et al: MicroRNA-155 suppresses activation-induced cytidine deaminase-mediated *Myc-Igh* translocation. Immunity 28: 630–638 (2008).
- Dorshorst BJ, Ashwell CM: Genetic mapping of the sex-linked barring gene in the chicken. Poult Sci 88:1811–1817 (2009).
- Dorshorst B, Okimoto R, Ashwell C: Genomic regions associated with dermal hyperpigmentation, polydactyly and other morphological traits in the Silkie chicken. J Hered 101:339– 350 (2010).

- Dorshorst B, Molin AM, Rubin CJ, Johansson AM, Strömstedt L, et al: A complex genomic rearrangement involving the *Endothelin 3* locus causes dermal hyperpigmentation in the chicken. PLoS Genet 7:e1002412 (2011).
- Dostie J, Richmond TA, Arnaout RA, Selzer RR, Lee WL, et al: Chromosome Conformation Capture Carbon Copy (5C): a massively parallel solution for mapping interactions between genomic elements. Genome Res 16: 1299–1309 (2006).
- Douaud M, Feve K, Pituello F, Gourichon D, Boitard S, et al: Epilepsy caused by an abnormal alternative splicing with dosage effect of the SV2A gene in a chicken model. PLoS One 6:e26932 (2011).
- Downing T, Lynn DJ, Connell S, Lloyd AT, Bhuiyan AK, et al: Evidence of balanced diversity at the chicken interleukin 4 receptor alpha chain locus. BMC Evol Biol 9:136–149 (2009a).
- Downing T, Lynn DJ, Connell S, Lloyd AT, Bhuiyan AK, et al: Contrasting evolution of diversity at two disease-associated chicken genes. Immunogenetics 61:303–314 (2009b).
- Doyle JM, Katzner TE, Bloom PH, Ji Y, Wijayawardena BK, DeWoody JA: The genome sequence of a widespread apex predator, the golden eagle (*Aquila chrysaetos*). PLoS One 9:e95599 (2014).
- Dunn IC, Paton IR, Clelland AK, Sebastian S, Johnson EJ, et al: The chicken polydactyly (Po) locus causes allelic imbalance and ectopic expression of *Shh* during limb development. Dev Dyn 240:1163–1172 (2011).
- Dunn K, Nazerian K: Induction of Marek's disease virus antigens by IdUrd in a chicken lymphoblastoid cell line. J Gen Virol 34:413–419 (1977).
- Dustin ML, Cooper JA: The immunological synapse and the actin cytoskeleton: molecular hardware for T cell signaling. Nat Immunol 1: 23–29 (2000).
- Ehlers MR: CR3: a general purpose adhesion-recognition receptor essential for innate immunity. Microbes Infect 2:289–294 (2000).
- Ejlerskov P, Rasmussen I, Nielsen TT, Bergström AL, Tohyama Y, et al: Tubulin polymerization-promoting protein (TPPP/p25α) promotes unconventional secretion of α-synuclein through exophagy by impairing autophagosome-lysosome fusion. J Biol Chem 288:17313–17335 (2013).
- Elferink MG, Vallée AAA, Jungerius A, Crooijmans RPMA, Groenen MAM: Partial duplication of the *PRLR* and *SPEF2* genes at the late feathering locus in chicken. BMC Genomics 9:391–399 (2008).
- Elferink MG, van As P, Veenendaal T, Crooijmans RPMA, Groenen MAM: Regional differences in recombination hotspots between two chicken populations. BMC Genet 11:11 (2010).
- Elferink MG, Megens HJ, Vereijken A, Hu XX, Crooijmans RPMA, Groenen MAM: Signatures of selection in the genomes of commercial and non-commercial chicken breeds. PLoS One 7:e32720 (2012).

- Elkin RG, Bauer R, Schneider WJ: The restricted ovulator chicken strain: an oviparous vertebrate model of reproductive dysfunction caused by a gene defect affecting an oocytespecific receptor. Anim Reprod Sci 136:1–13 (2012).
- Ellegren H: Emergence of male-biased genes on the chicken Z-chromosome: sex-chromosome contrasts between male and female heterogametic systems. Genome Res 21:2082– 2086 (2011).
- Ellegren H: The evolutionary genomics of birds. Annu Rev Ecol Evol Syst 44:239–259 (2013).
- Ellegren H: Genome sequencing and population genomics in non-model organisms. Trends Ecol Evol 29:51–63 (2014).
- Ellegren H, Hultin-Rosenberg L, Brunström B, Dencker L, Kultima K, Scholz B: Faced with inequality: chicken do not have a general dosage compensation of sex-linked genes. BMC Biol 5:40 (2007).
- Ellegren H, Smeds L, Burri R, Olason PI, Backström N, et al: The genomic landscape of species divergence in *Ficedula* flycatchers. Nature 491:756–760 (2012).
- Ellis HL, Shioda K, Rosenthal NF, Coser KR, Shioda T: Masculine epigenetic sex marks of the *CYP19A1/aromatase* promoter in genetically male chicken embryonic gonads are resistant to estrogen-induced phenotypic sex conversion. Biol Reprod 87:23 (2012).
- ENCODE Project Consortium: An integrated encyclopedia of DNA elements in the human genome. Nature 489:57–74 (2012).
- Eöry L, Thomas M, Gilbert P, Li C, Li B, et al: Avianbase: a community resource for bird genomics. Genome Biol 16:21 (2015).
- Erdmann VA, Szymanski M, Hochberg A, Groot N, Barciszewski J: Non-coding, mRNA-like RNAs database Y2K. Nucleic Acids Res 28: 197–200 (2000).
- Eriksson J, Larson G, Gunnarsson U, Bed'hom B, Tixier-Boichard M, et al: Identification of the *yellow skin* gene reveals a hybrid origin of the domestic chicken. PLoS Genet 4:e1000010 (2008).
- Erlanger BF, Beiser SM: Antibodies specific for ribonucleosides and ribonucleotides and their reaction with DNA. Proc Natl Acad Sci USA 52:68–74 (1964).
- Erlanger BF, Senitzer D, Miller OJ, Beiser SM: Nucleic acid-reactive antibodies specific for nucleosides and nucleotides. Karolinska Symposia on Research Methods in Reproductive Endocrinology, 5th Symposium: Gene Transcription in Reproductive Tissue (1972).
- Ervasti JM, Campbell KP: A role for the dystrophin-glycoprotein complex as a transmembrane linker between laminin and actin. J Cell Biol 122:809–823 (1993).
- Fan WL, Ng CS, Chen CF, Lu MY, Chen YH, et al: Genome-wide patterns of genetic variation in two domestic chickens. Genome Biol Evol 5:1376–1392 (2013).
- Feduccia A: Explosive evolution in tertiary birds and mammals. Science 267:637–638 (1995).

- Feduccia A: 'Big bang' for tertiary birds? Trends Ecol Evol 18:172–176 (2003).
- Feng C, Gao Y, Dorshorst B, Song C, Gu X, et al: A *cis*-regulatory mutation of *PDSS2* causes silky-feather in chickens. PLoS Genet 10:e1004576 (2014).
- Fernando TR, Rodriguez-Malave NI, Rao DS: MicroRNAs in B cell development and malignancy. J Hematol Oncol 5:7 (2012).
- Fillon V, Morisson M, Zoorob R, Auffray C, Douaire M, et al: Identification of 16 chicken microchromosomes by molecular markers using two-colour fluorescence in situ hybridization (FISH). Chromosome Res 6:307–313 (1998).
- Flicek P, Amode MR, Barrell D, Beal K, Brent S, et al: Ensembl 2011. Nucleic Acids Res 39:D800–D806 (2011).
- Flicek P, Amode MR, Barrell D, Beal K, Billis K, et al: Ensembl 2014. Nucleic Acids Res 42:D749–D755 (2014).
- Flink LG, Allen R, Barnett R, Malmström H, Peters J, et al: Establishing the validity of domestication genes using DNA from ancient chickens. Proc Natl Acad Sci USA 111:6184–6189 (2014).
- Flusberg BA, Webster DR, Lee JH, Travers KJ, Olivares EC, et al: Direct detection of DNA methylation during single-molecule, real-time sequencing. Nat Methods 7:461–465 (2010).
- Fragnet L, Blasco MA, Klapper W, Rasschaert D: The RNA subunit of telomerase is encoded by Marek's disease virus. J Virol 77:5985–5996 (2003).
- Frankl-Vilches C, Kuhl H, Werber M, Klages S, Kerick M, et al: Using the canary genome to decipher the evolution of hormone-sensitive gene regulation in seasonal singing birds. Genome Biol 16:19 (2015).
- Frésard L, Leroux S, Servin B, Gourichon D, Dehais P, et al: Transcriptome-wide investigation of genomic imprinting in chicken. Nucleic Acids Res 42:3768–3782 (2014).
- Fridolfsson AK, Cheng H, Copeland NG, Jenkins NA, Liu HC, et al: Evolution of the avian sex chromosomes from an ancestral pair of autosomes. Proc Natl Acad Sci USA 95:8147–8152 (1998).
- Fulton JE: Genomic selection for poultry breeding. Anim Front 2:30–36 (2012).
- Fumihito A, Miyake T, Sumi S, Takada M, Ohno S, Kondo N: One subspecies of the red junglefowl (*Gallus gallus gallus*) suffices as the matriarchic ancestor of all domestic breeds. Proc Natl Acad Sci USA 91:12505–12509 (1994).
- Fumihito A, Miyake T, Takada M, Shingu R, Endo T, et al: Monophyletic origin and unique dispersal patterns of domestic fowls. Proc Natl Acad Sci USA 93:6792–6795 (1996).
- Furlan-Magaril M, Rincon-Arano H, Recillas-Targa F: Sequential chromatin immunoprecipitation protocol: ChIP-reChIP. Methods Mol Biol 543:253–266 (2009).
- Gaffney DJ, McVicker G, Pai AA, Fondufe-Mittendorf YN, Lewellen N, et al: Controls of nucleosome positioning in the human genome. PLoS Genet 8:e1003036 (2012).

- Ganapathy G, Howard JT, Ward JM, Li J, Li B, et al: High-coverage sequencing and annotated assemblies of the budgerigar genome. Gigascience 3:11 (2014).
- Gano JJ, Simon JA: A proteomic investigation of ligand-dependent HSP90 complexes reveals CHORDC1 as a novel ADP-dependent HSP90-interacting protein. Mol Cell Proteomics 9:255–270 (2010).
- Garcia-Moreno J, Sorenson MD, Mindell DP: Congruent avian phylogenies inferred from mitochondrial and nuclear DNA sequences. J Mol Evol 57:27–37 (2003).
- Gardner PP, Fasold M, Burge SW, Ninova M, Hertel J, et al: Conservation and losses of noncoding RNAs in avian genomes. PLoS One 10:e0121797 (2015).
- Garro AG, Erlanger BF, Beiser SM: Specificity in the reaction between anti-pyrimidine nucleoside antibodies and DNA, in Plescia OJ, Braun W (eds): Nucleic Acids in Immunology, pp 47–55 (Springer, New York 1968).
- Genome 10K Community of Scientists: Genome 10K: a proposal to obtain whole-genome sequence for 10,000 vertebrate species. J Hered 100:659–674 (2009).
- Ghebrewold RAR, Heidaritabar M, Vereijken A, Ducro BJ, Bastiaansen JWM: A genomewide association study for egg shell strength in the genome of brown-egg layers. 2014 Joint Annual Meeting, Kansas City, MO, July 20–24, 2014. Available at https://asas. confex.com/asas/jam2014/webprogram/ Paper5582.html
- Gheyas AA, Burt DW: Microarray resources for genetic and genomic studies in chicken: a review. Genesis 51:337–356 (2013).
- Gheyas AA, Boschiero C, Eory L, Ralph H, Kuo R, Woolliams JA, Burt DW: Functional classification of 15 million SNPs detected from diverse chicken populations. DNA Res 2015, DOI: 10.1093/dnares/dsv005.
- Gholami M, Erbe M, Gärke C, Preisinger R, Weigend A, et al: Population genomic analyses based on 1 million SNPs in commercial egg layers. PLoS One 9:e94509 (2014).
- Gimeno IM, Witter RL, Reed WM: Four distinct neurologic syndromes in Marek's disease: effect of viral strain and pathotype. Avian Dis 43:721–737 (1999).
- Gimeno IM, Witter RL, Cortes AL, Reed WM: Replication ability of three highly protective Marek's disease vaccines: implications in lymphoid organ atrophy and protection. Avian Pathol 40:573–579 (2011).
- Giresi PG, Kim J, McDaniell RM, Iyer VR, Lieb JD: FAIRE (Formaldehyde-Assisted Isolation of Regulatory Elements) isolates active regulatory elements from human chromatin. Genome Res 17:877–885 (2007).
- Giuffra E, Törnsten A, Marklund S, Bongcam-Rudloff E, Chardon P, et al: A large duplication associated with dominant white color in pigs originated by homologous recombination between LINE elements flanking KIT. Mamm Genome 13:569–577 (2002).

- Glazov EA, Cottee PA, Barris WC, Moore RJ, Dalrymple BP, Tizard ML: A microRNA catalog of the developing chicken embryo identified by a deep sequencing approach. Genome Res 18:957–964 (2008).
- Godoy TF, Moreira GCM, Boschiero C, Gheyas AA, Gasparin G, et al: SNP and INDEL detection in a QTL region on chicken chromosome 2 associated with muscle deposition. Anim Genet 46:158–163 (2015).
- Goralski TJ, Edström JE, Baker BS: The sex determination locus *transformer-2* of *Drosophila* encodes a polypeptide with similarity to RNA binding proteins. Cell 56:1011–1018 (1989).
- Govoroun MS, Pannetier M, Pailhoux E, Cocquet J, Brillard JP, et al: Isolation of chicken homolog of the *FOXL2* gene and comparison of its expression patterns with those of aromatase during ovarian development. Dev Dyn 231: 859–870 (2004).
- Graves JAM: Sex and death in birds: a model of dosage compensation that predicts lethality of sex chromosome aneuploids. Cytogenet Genome Res 101:278–282 (2003).
- Graves JAM: Avian sex, sex chromosomes, and dosage compensation in the age of genomics. Chromosome Res 22:45–57 (2014).
- Greer EL, Shi Y: Histone methylation: a dynamic mark in health, disease and inheritance. Nat Rev Genet 13:343–357 (2012).
- Griffin D, Burt DW: All chromosomes great and small: 10 years on. Chromosome Res 22:1–6 (2014).
- Griffin DK, Robertson LB, Tempest HG, Skinner BM: The evolution of the avian genome as revealed by comparative molecular cytogenetics. Cytogenet Genome Res 117:64–77 (2007).
- Griffin DK, Robertson LB, Tempest HG, Vignal A, Fillon V, et al: Whole genome comparative studies between chicken and turkey and their implications for avian genome evolution. BMC Genomics 9:168 (2008).
- Groenen MAM, Wahlberg P, Foglio M, Cheng HH, Megens HJ, et al: A high-density SNP-based linkage map of the chicken genome reveals sequence features correlated with recombination rate. Genome Res 19:510–519 (2009).
- Groenen MAM, Megens HJ, Zare Y, Warren WC, Hillier LW, et al: The development and characterization of a 60K SNP chip for chicken. BMC Genomics 12:274 (2011).
- Groenen MA, Archibald AL, Uenishi H, Tuggle CK, Takeuchi Y, et al: Analyses of pig genomes provide insight into porcine demography and evolution. Nature 491:393–398 (2012).
- Groth JG, Barrowclough GF: Basal divergences in birds and the phylogenetic utility of the nuclear RAG-1 gene. Mol Phylogenet Evol 12:115– 123 (1999).
- Grützner F, Zend-Ajusch E, Stout K, Munsche S, Niveleau A, et al: Chicken microchromosomes are hypermethylated and can be identified by specific painting probes. Cytogenet Cell Genet 93:265–269 (2001).

- Guioli S, Lovell-Badge R: PITX2 controls asymmetric gonadal development in both sexes of the chick and can rescue the degeneration of the right ovary. Development 134:4199–4208 (2007).
- Guioli S, Nandi S, Zhao D, Burgess-Shannon J, Lovell-Badge R, Clinton M: Gonadal asymmetry and sex determination in birds. Sex Dev 8:227-242 (2014).
- Gunnarsson U, Kerje S, Bed'hom B, Sahlqvist AS, Ekwall O, et al: The dark brown plumage color in chickens is caused by an 8.3-kb deletion upstream of *SOX10*. Pigment Cell Melanoma Res 24:268–274 (2011).
- Hackett SJ, Kimball RT, Reddy S, Bowie RCK, Braun EL, et al: A phylogenomic study of birds reveals their evolutionary history. Science 320:1763–1768 (2008).
- Haddrath O, Baker AJ: Complete mitochondrial DNA genome sequences of extinct birds: ratite phylogenetics and the vicariance biogeography hypothesis. Proc Biol Sci 268:939–945 (2001).
- Hagiwara M: Alternative splicing: a new drug target of the post-genome era. Biochim Biophys Acta 1754:324–331 (2005).
- Halley YA, Dowd SE, Decker JE, Seabury PM, Bhattarai E, et al: A draft de novo genome assembly for the northern bobwhite (*Colinus virginianus*) reveals evidence for a rapid decline in effective population size beginning in the Late Pleistocene. PLoS One 9:e90240 (2014).
- Hamburger V, Hamilton HL: A series of normal stages in the development of the chick embryo. Dev Dyn 195:231–272 (1992).
- Han R, Yang P, Tian Y, Wang D, Zhang Z, et al: Identification and functional characterization of copy number variations in diverse chicken breeds. BMC Genomics 15:934 (2014).
- Härlid A, Arnason U: Analyses of mitochondrial DNA nest ratite birds within the Neognathae: supporting a neotenous origin of ratite morphological characters. Proc R Soc Lond B Biol 266:305–309 (1999).
- Hashimoto-Tane A, Yokosuka T, Sakata-Sogawa K, Sakuma M, Ishihara C, et al: Dynein-driven transport of T cell receptor microclusters regulates immune synapse formation and T cell activation. Immunity 34:919–931 (2011).
- Hayward WS, Neel BG, Astrin SM: Activation of a cellular *onc* gene by promoter insertion in ALV-induced lymphoid leukosis. Nature 290: 475–480 (1981).
- He Y, Carrillo JA, Luo J, Ding Y, Tian F, et al: Genome-wide mapping of DNase I hypersensitive sites and association analysis with gene expression in MSB1 cells. Front Genet 5:308 (2014).
- Hedges SB: Amniote phylogeny and the position of turtles. BMC Biol 10:64 (2012).
- Hedges SB, Kumar S: Genomic clocks and evolutionary timescales. Trends Genet 19:200–206 (2003).
- Hedges SB, Kumar S: Precision of molecular time estimates. Trends Genet 20:242–247 (2004).

169

tatsbibliothek Würzburg .78.1 - 7/15/2015 10:03:51

- Hedges SB, Sibley CG: Molecules vs. morphology in avian evolution: the case of the 'pelecaniform' birds. Proc Natl Acad Sci USA 91:9861– 9865 (1994).
- Hedges SB, Simmons MD, VanDikj MA, Caspers G, DeJong WW, Sibley CG: Phylogenetic relationships of the hoatzin, an enigmatic South American bird. Proc Natl Acad Sci USA 92: 11662–11665 (1995).
- Hedges SB, Parker PH, Sibley CG, Kumar S: Continental breakup and the ordinal diversification of birds and mammals. Nature 381:226– 229 (1996).
- Hellström AR, Sundström E, Gunnarsson U, Bed'Hom B, et al: Sex-linked barring in chickens is controlled by the *CDKN2A/B* tumour suppressor locus. Pigment Cell Melanoma Res 23:521–530 (2010).
- Hillel J, Granevitze Z, Twito T, Ben-Avraham D, Blum S, et al: Molecular markers for the assessment of chicken biodiversity. Worlds Poult Sci J 63:33–45 (2007).
- Hocking PM, Guggenheim JA: The chick as an animal model of eye disease. Drug Disc Today Dis Models 10:225–230 (2013).
- Hofacker IL, Fontana W, Stadler PF, Bonhoeffer S, Tacker M, Schuster P: Fast folding and comparison of RNA secondary structures. Monatsh Chemie 125:167–188 (1994).
- Hogenkamp A, Isohadouten N, Reemers SS, Romijn RA, Hemrika W, et al: Chicken lung lectin is a functional C-type lectin and inhibits haemagglutination by influenza A virus. Vet Microbiol 130:37–46 (2008).
- Hori T, Asakawa S, Itoh Y, Shimizu N, Mizuno S: *Wpkci*, encoding an altered form of *PKCI*, is conserved widely on the avian W chromosome and expressed in early female embryos: implication of its role in female sex determination. Mol Biol Cell 11:3645–3660 (2000).
- Hsu PD, Lander ES, Zhang F: Development and applications of CRISPR-Cas9 for genome engineering. Cell 157:1262–1278 (2014).
- Hu Y, Xu H, Li Z, Zheng X, Jia X, et al: Comparison of the genome-wide DNA methylation profiles between fast-growing and slowgrowing broilers. PLoS One 8:e56411 (2013).
- Huang Y, Zhao Y, Haley CS, Hu S, Hao J, et al: A genetic and cytogenetic map for the duck (*Anas platyrhynchos*). Genetics 173:287-296 (2006).
- Huang Y, Li Y, Burt DW, Chen H, Zhang Y, et al: The duck genome and transcriptome provide insight into an avian influenza virus reservoir species. Nat Genet 45:776–783 (2013).
- Hughes AL, Hughes MK: Small genomes for better flyers. Nature 377:391 (1995).
- Hughes JM, Baker AJ: Phylogenetic relationships of the enigmatic hoatzin (*Opisthocomus hoazin*) resolved using mitochondrial and nuclear gene sequences. Mol Biol Evol 16:1300– 1307 (1999).
- Hyun BR, McElwee JL, Soloway PD: Single molecule and single cell epigenomics. Methods 72:41-50 (2015).

- Imsland F, Feng C, Boije H, Bed'hom B, Fillon V, et al: The *Rose-comb* mutation in chickens constitutes a structural rearrangement causing both altered comb morphology and defective sperm motility. PLoS Genet 8:e1002775 (2012).
- International Chicken Genome Sequencing Consortium: Sequence and comparative analysis of the chicken genome provide unique perspectives on vertebrate evolution. Nature 432: 695–716 (2004).
- International Chicken Polymorphism Map Consortium: A genetic variation map for chicken with 2.8 million single-nucleotide polymorphisms. Nature 432:717–722 (2004).
- International HapMap Consortium: A haplotype map of the human genome. Nature 437:1299– 1320 (2005).
- Iseli C, Jongeneel CV, Bucher P: ESTScan: a program for detecting, evaluating, and reconstructing potential coding regions in EST sequences. Proc Int Conf Intell Syst Mol Biol 99: 138–148 (1999).
- Ishimaru Y, Komatsu T, Kasahara M, Katoh-Fukui Y, Ogawa H, et al: Mechanism of asymmetric ovarian development in chick embryos. Development 135:677–685 (2008).
- Ishishita S, Tsuruta Y, Uno Y, Nakamura A, Nishida C, et al: Chromosome size-correlated and chromosome size-uncorrelated homogenization of centromeric repetitive sequences in New World quails. Chromosome Res 22: 15–34 (2014).
- Islam A, Wong CW, Walkden-Brown SW, Colditz IG, Arzey KE, Groves PJ: Immunosuppressive effects of Marek's disease virus (MDV) and herpesvirus of turkeys (HVT) in broiler chickens and the protective effect of HVT vaccination against MDV challenge. Avian Pathol 31:449–461 (2002).
- Itoh Y, Melamed E, Yang X, Kampf K, Wang S, et al: Dosage compensation is less effective in birds than in mammals. J Biol 6:2 (2007).
- IUCN: The IUCN Redlist of Threatened Species (International Union for the Conservation of Nature, Cambridge 2014).
- Janes DE, Ezaz T, Graves JA, Edwards SV: Characterization, chromosomal location, and genomic neighborhood of a ratite ortholog of a gene with gonadal expression in mammals. Integr Comp Biol 48:505–511 (2008).
- Janes DE, Ezaz T, Marshall Graves JA, Edwards SV: Recombination and nucleotide diversity in the sex chromosomal pseudoautosomal region of the emu, *Dromaius novaehollandiae*. J Hered 100:125–136 (2009).
- Janes DE, Valenzuela N, Ezaz T, Amemiya C, Edwards SV: Sex chromosome evolution in amniotes: applications for bacterial artificial chromosome libraries. J Biomed Biotechnol 2011:132975 (2011).
- Jang HJ, Seo HW, Lee BR, Yoo M, Womack JE, Han JY: Gene expression and DNA methylation status of chicken primordial germ cells. Mol Biotechnol 54:177–186 (2013).
- Janz S: *Myc* translocations in B cell and plasma cell neoplasms. DNA Repair 5:1213–1224 (2006).

- Jarvis ED, Mirarab S, Aberer AJ, Li B, Houde P, et al: Whole-genome analyses resolve early branches in the tree of life of modern birds. Science 346:1320–1331 (2014).
- Jeffares DC, Poole AM, Penny D: Relics from the RNA world. J Mol Evol 46:18–36 (1998).
- Jia X, Chen S, Zhou H, Li D, Liu W, Yang N: Copy number variation identified in the chicken using a 60K SNP BeadChip. Anim Genet 44: 276–284 (2013).
- Johansson US, Parsons TJ, Irestedt M, Ericson PGP: Clades within the 'higher land birds', evaluated by nuclear DNA sequences. J Zool Syst Evol Res 39:37–51 (2001).
- Johansson US, Irestedt M, Parsons TJ, Ericson PGP: Basal phylogeny of the Tyrannoidea based on comparisons of cytochrome *b* and exons of nuclear *c-myc* and RAG-1 genes. Auk 119:984–995 (2002).
- Kane MD, Jatkoe TA, Stumpf CR, Lu J, Thomas JD, Madore SJ: Assessment of the sensitivity and specificity of oligonucleotide (50mer) microarrays. Nucleic Acids Res 28:4552–4557 (2000).
- Kanginakudru S, Metta M, Jakati RD, Nagaraju J: Genetic evidence from Indian red jungle fowl corroborates multiple domestication of modern day chicken. BMC Evol Biol 8:174 (2008).
- Kaplan N, Moore IK, Fondufe-Mittendorf Y, Gossett AJ, Tillo D, et al: The DNA-encoded nucleosome organization of a eukaryotic genome. Nature 458:362–366 (2009).
- Katz Y, Wang ET, Airoldi EM, Burge CB: Analysis and design of RNA sequencing experiments for identifying isoform regulation. Nat Methods 7:1009–1015 (2010).
- Kaufer BB, Jarosinski KW, Osterrieder N: Herpesvirus telomeric repeats facilitate genomic integration into host telomeres and mobilization of viral DNA during reactivation. J Exp Med 208:605–615 (2011).
- Kawahara-Miki R, Sano S, Nunome M, Shimmura T, Kuwayama T, et al: Next-generation sequencing reveals genomic features in the Japanese quail. Genomics 101:345–353 (2013).
- Kawakami T, Smeds L, Backström N, Husby A, Qvarnström A, et al: A high-density linkage map enables a second-generation collared flycatcher genome assembly and reveals the patterns of avian recombination rate variation and chromosomal evolution. Mol Ecol 23: 4035–4058 (2014).
- Kayagaki N, Phung Q, Chan S, Chaudhari R, Quan C, et al: DUBA: a deubiquitinase that regulates type I interferon production. Science 318:1628–1632 (2007).
- Kellner WA, Sullivan RT, Carlson BH, NISC Comparative Sequencing Program, Thomas JW: Uprobe: a genome-wide universal probe resource for comparative physical mapping in vertebrates. Genome Res 15:166–173 (2005).
- Kelly TK, Liu Y, Lay FD, Liang G, Berman BP, Jones PA: Genome-wide mapping of nucleosome positioning and DNA methylation within individual DNA molecules. Genome Res 22:2497–2506 (2012).

- King M: Species Evolution: the Role of Chromosome Change (Cambridge University Press, Cambridge 1995).
- Kingma PS, Whitsett JA: In defense of the lung: surfactant protein A and surfactant protein D. Curr Opin Pharmacol 6:277–283 (2006).
- Kishi M, Bradley G, Jessip J, Tanaka A, Nonoyama M: Inverted repeat regions of Marek's disease virus DNA possess a structure similar to that of the *a* sequence of herpes simplex virus DNA and contain host cell telomere sequences. J Virol 65:2791–2797 (1991).
- Koboldt DC, Steinberg KM, Larson DE, Wilson RK, Mardis ER: The next-generation sequencing revolution and its impact on genomics. Cell 155:27–38 (2013).
- Kokalj-Vokac N, Zagorac A, Pristovnik M, Burgeois CA, Dutrillaux B: DNA methylation of the extraembryonic tissues: an in situ study on human metaphase chromosomes. Chromosome Res 6:161–166 (1998).
- Kos R, Reedy MV, Johnson RL, Erickson CA: The winged-helix transcription factor FoxD3 is important for establishing the neural crest lineage and repressing melanogenesis in avian embryos. Development 128:1467–1479 (2001).
- Kos R, Tucker RP, Hall R, Duong TD, Erickson CA: Methods for introducing morpholinos into the chicken embryo. Dev Dyn 226:470– 477 (2003).
- Koskela K, Kohonen P, Nieminen P, Buerstedde JM, Lassila O: Insight into lymphoid development by gene expression profiling of avian B cells. Immunogenetics 55:412–422 (2003).
- Kothapalli NR, Collura KM, Norton DD, Fugmann SD: Separation of mutational and transcriptional enhancers in Ig genes. J Immunol 187:3247–3255 (2011).
- Kouzarides T: Chromatin modifications and their function. Cell 128:693–705 (2007).
- Kozomara A, Griffiths-Jones S: miRBase: annotating high confidence microRNAs using deep sequencing data. Nucleic Acids Res 42:D68–D73 (2014).
- Kranis A, Gheyas AA, Boschiero C, Turner F, Yu L, et al: Development of a high density 600K SNP genotyping array for chicken. BMC Genomics 14:59 (2013).
- Kumar S, Hedges SB: A molecular timescale for vertebrate evolution. Nature 392:917–920 (1998).
- Kurochkin EN, Dyke GJ, Karhu AA: A new presbyornithid bird (Aves, Anseriformes) from the late Cretaceous of Southern Mongolia. Am Mus Novit 1–11 (2002).
- Laird PW: Principles and challenges of genomewide DNA methylation analysis. Nat Rev Genet 11:191–203 (2010).
- Lambeth LS, Cummins DM, Doran TJ, Sinclair AH, Smith CA: Overexpression of aromatase alone is sufficient for ovarian development in genetically male chicken embryos. PLoS One 8:e68362 (2013).

- Lambeth LS, Raymond CS, Roeszler KN, Kuroiwa A, Nakata T, et al: Over-expression of DMRT1 induces the male pathway in embryonic chicken gonads. Dev Biol 389:160–172 (2014).
- Lan H, Chen H, Chen LC, Wang BB, Sun L, et al: The first report of a Pelecaniformes defensin cluster: characterization of β -defensin genes in the crested ibis based on BAC libraries. Sci Rep 4:6923 (2014).
- Langhorst LJ, Fechheimer NS: Shankless, a new mutation on chromosome 2 in the chicken. J Hered 76:182–186 (1985).
- Larkin DM, Pape G, Donthu R, Auvil L, Welge M, Lewin HA: Breakpoint regions and homologous synteny blocks in chromosomes have different evolutionary histories. Genome Res 19:770–777 (2009).
- Law MJ, Lower KM, Voon HP, Hughes JR, Garrick D, et al: ATR-X syndrome protein targets tandem repeats and influences allele-specific expression in a size-dependent manner. Cell 143:367–378 (2010).
- Lee K, Feinstein J, Cracraft JC: The phylogeny of ratite birds: resolving conflicts between molecular and morphological data sets, in Mindell DP (ed): Avian Molecular Evolution and Systematics, pp 173–211 (Academic Press, San Diego 1997).
- Lee LF, Bacon LD: Ontogeny and line differences in the mitogenic response of chicken lymphocytes. Poult Sci 62:579–584 (1983).
- Lee LF, Wu P, Sui D, Ren D, Kamil J, et al: The complete unique long sequence and the overall genomic organization of the GA strain of Marek's disease virus. Proc Natl Acad Sci USA 97:6091–6096 (2000).
- Lee MK, Ren CW, Yan B, Cox B, Zhang HB, et al: Construction and characterization of three BAC libraries for analysis of the chicken genome. Anim Genet 34:151–152 (2003).
- Lee SI, Lee WK, Shin JH, Han BK, Moon S, et al: Sexually dimorphic gene expression in the chick brain before gonadal differentiation. Poult Sci 88:1003–1015 (2009).
- Leroy G, Kayang BB, Youssao IA, Yapi-Gnaoré CV, Osei-Amponsah R, et al: Gene diversity, agroecological structure and introgression patterns among village chicken populations across North, West and Central Africa. BMC Genet 13:34 (2012).
- Lewin HA, Larkin DM, Pontius J, O'Brien SJ: Every genome sequence needs a good map. Genome Res 19:1925–1928 (2009).
- Li C, Zhang Y, Li J, Kong L, Hu H, et al: Two Antarctic penguin genomes reveal insights into their evolutionary history and molecular changes related to the Antarctic environment. Gigascience 3:27 (2014).
- Li H, Durbin R: Fast and accurate short read alignment with Burrows-Wheeler transform. Bioinformatics 25:1754–1760 (2009).
- Li H, Durbin R: Inference of human population history from individual whole-genome sequences. Nature 475:493–496 (2011).

- Li H, Ruan J, Durbin R: Mapping short DNA sequencing reads and calling variants using mapping quality scores. Genome Res 18: 1851–1858 (2008).
- Li Q, Li N, Hu X, Li J, Du Z, et al: Genome-wide mapping of DNA methylation in chicken. PLoS One 6:e19428 (2011).
- Li S, Li B, Cheng C, Xiong Z, Liu Q, et al: Genomic signatures of near-extinction and rebirth of the crested ibis and other endangered bird species. Genome Biol 15:557 (2014).
- Lieberman-Aiden E, van Berkum NL, Williams L, Imakaev M, Ragoczy T, et al: Comprehensive mapping of long-range interactions reveals folding principles of the human genome. Science 326:289–293 (2009).
- Lindblad-Toh K, Garber M, Zuk O, Lin MF, Parker BJ, et al: A high-resolution map of human evolutionary constraint using 29 mammals. Nature 478:476–482 (2011).
- Lithgow PE, O'Connor R, Smith D, Fonseka G, Al Mutery A, et al: Novel tools for characterising inter and intra chromosomal rearrangements in avian microchromosomes. Chromosome Res 22:85–97 (2014).
- Liu W, Liu Z, Hu X, Zhang Y, Yuan J, et al: Construction and characterization of a novel 13.34-fold chicken bacterial artificial chromosome library. Anim Biotechnol 14:145– 153 (2003).
- Liu YP, Wu GS, Yao YG, Miao YW, Luikart G, et al: Multiple maternal origins of chickens: out of the Asian jungles. Mol Phylogenet Evol 38: 12–19 (2006).
- Lohe A, Roberts P: Evolution of satellite DNA sequences in *Drosophila*, in Verma RS (ed): Heterochromatin. Molecular and Structural Aspects, pp 148–186 (Cambridge University Press, Cambridge 1988).
- Long EO, Dawid IB: Repeated genes in eukaryotes. Ann Rev Biochem 49:727–764 (1980).
- Lopez AJ: Alternative splicing of pre-mRNA: developmental consequences and mechanisms of regulation. Annu Rev Genet 32:279–305 (1998).
- Lovell PV, Wirthlin M, Wilhelm L, Minx P, Lazar NH, et al: Conserved syntenic clusters of protein coding genes are missing in birds. Genome Biol 15:565–592 (2014).
- Lowe TM, Eddy SR: tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. Nucleic Acids Res 25:955– 964 (1997).
- Luo CL, Qu H, Ma J, Wang J, Li CY, et al: Genome-wide association study of antibody response to Newcastle disease virus in chicken. BMC Genet 14:42 (2013).
- Luo CL, Qu H, Ma J, Wang J, Hu XX, et al: A genome-wide association study identifies major loci affecting the immune response against infectious bronchitis virus in chicken. Infect Genet Evol 21:351–358 (2014).
- Luo J, Mitra A, Tian F, Chang S, Zhang H, et al: Histone methylation analysis and pathway predictions in chickens after MDV infection. PLoS One 7:e41849 (2012).

- Luo J, Yu Y, Mitra A, Chang S, Zhang H, et al: genome-wide copy number variant analysis in inbred chickens lines with different susceptibility to Marek's disease. G3 (Bethesda) 3: 217–223 (2013).
- Luo M, Yu Y, Kim H, Kudrna D, Itoh Y, et al: Utilization of a zebra finch BAC library to determine the structure of an avian androgen receptor genomic region. Genomics 87:181– 190 (2006).
- Lyimo CM, Weigend S, Msoffe PL, Eding H, Simianer H, Weigend S: Global diversity and genetic contributions of chicken populations from African, Asian and European regions. Anim Genet 45:836–848 (2014).
- Lynch KW: Consequences of regulated premRNA splicing in the immune system. Nat Rev Immunol 4:931–940 (2004).
- Maas SA, Suzuki T, Fallon JF: Identification of spontaneous mutations within the long-range limb-specific *Sonic hedgehog* enhancer (ZRS) that alter *Sonic hedgehog* expression in the chicken limb mutants *oligozeugodactyly* and Silkie breed. Dev Dyn 240:1212–1222 (2011).
- Maatouk DM, DiNapoli L, Alvers A, Parker KL, Taketo MM, Capel B: Stabilization of β-catenin in XY gonads causes male-to-female sex-reversal. Hum Mol Genet 17:2949– 2955 (2008).
- Madsen BE, Villesen P, Wiuf C: Short tandem repeats in human exons: a target for disease mutations. BMC Genomics 9:410 (2008).
- Maher B: Personal genomes: the case of the missing heritability. Nature 456:18–21 (2008).
- Mamanova L, Coffey AJ, Scott CE, Kozarewa I, Turner EH, et al: Target-enrichment strategies for next-generation sequencing. Nat Methods 7:111–118 (2010).
- Mank JE, Ellegren H: All dosage compensation is local: gene-by-gene regulation of sex-biased expression on the chicken Z chromosome. Heredity 102:312–320 (2009).
- Marz M, Kirsten T, Stadler PF: Evolution of spliceosomal snRNA genes in metazoan animals. J Mol Evol 67:594–607 (2008).
- Masabanda J, Burt DW, O'Brien PCM, Vignal A, Fillon V, et al: Molecular cytogenetic definition of the chicken genome: the first complete avian karyotype. Genetics 166:1367–1373 (2004).
- Matson CK, Zarkower D: Sex and the singular DM domain: insights into sexual regulation, evolution and plasticity. Nat Rev Genet 13: 163–174 (2012).
- Matson CK, Murphy MW, Sarver AL, Griswold MD, Bardwell VJ, Zarkower D: DMRT1 prevents female reprogramming in the postnatal mammalian testis. Nature 476:101–104 (2011).
- Matsumine H, Herbst MA, Ou SH, Wilson JD, McPhaul MJ: Aromatase mRNA in the extragonadal tissues of chickens with the hennyfeathering trait is derived from a distinctive promoter structure that contains a segment of a retroviral long terminal repeat. Functional organization of the Sebright, Leghorn, and Campine aromatase genes. J Biol Chem 266: 19900–19907 (1991).

- Matsumoto H, Maruse H, Inaba Y, Yoshizawa K, Sasazaki S, et al: The ubiquitin ligase gene *(WWP1)* is responsible for the chicken muscular dystrophy. FEBS Lett 582:2212–2218 (2008).
- Matsumoto Y, Buemio A, Chu R, Vafaee M, Crews D: Epigenetic control of gonadal aromatase (*cyp19a1*) in temperature-dependent sex determination of red-eared slider turtles. PLoS One 8:e63599 (2013).
- Mayer W, Niveleau A, Walter J, Fundele R, Haaf T: Demethylation of the zygotic paternal genome. Nature 403:501–502 (2000).
- McGowan P, Madge S: Pheasants, Partridges & Grouse: Including Buttonquails, Sandgrouse and Allies (A&C Black Publishers, London 2010).
- McPherson MC, Robinson CM, Gehlen LP, Delany ME: Comparative cytogenomics of poultry: mapping of single gene and repeat loci in the Japanese quail (*Coturnix japonica*). Chromosome Res 22:71–83 (2014).
- McQueen HA, Clinton M: Avian sex chromosomes: dosage compensation matters. Chromosome Res 17:687–697 (2009).
- McQueen HA, Fantes J, Cross SH, Clark VH, Archibald AL, Bird AP: CpG islands of chicken are concentrated on microchromosomes. Nat Genet 12:321–324 (1996).
- McQueen HA, Siriaco G, Bird AP: Chicken microchromosomes are hyperacetylated, early replicating, and gene rich. Genome Res 8: 621–630 (1998).
- Megens HJ, Groenen MAM: Domesticated species form a treasure-trove for molecular characterization of Mendelian traits by exploiting the specific genetic structure of these species in across-breed genome wide association studies. Heredity 109:1–3 (2012).
- Megens HJ, Crooijmans RP, Bastiaansen JW, Kerstens HH, Coster A, et al: Comparison of linkage disequilibrium and haplotype diversity on macro- and microchromosomes in chicken. BMC Genet 10:86 (2009).
- Mekchay S, Supakankul P, Assawamakin A, Wilantho A, Chareanchim W, Tongsima S: Population structure of four Thai indigenous chicken breeds. BMC Genet 15:40 (2014).
- Melamed E, Elashoff D, Arnold A: Evaluating dosage compensation on the chicken Z chromosome: should effective dosage compensation eliminate sexual bias? Heredity 103:357– 359 (2009).
- Mellor P, Furber LA, Nyarko JN, Anderson DH: Multiple roles for the p85α isoform in the regulation and function of PI3K signalling and receptor trafficking. Biochem J 441:23–37 (2012).
- Mendel G: Versuche über Pflanzenhybriden. Verhandlungen des naturforschenden Vereines in Brünn 10 (1866). Available from http:// www.gutenberg.org/ebooks/40854
- Meredith RW, Zhang G, Gilbert MT, Jarvis ED, Springer MS: Evidence for a single loss of mineralized teeth in the common avian ancestor. Science 346:1254390 (2014).

- Messer PW, Arndt PF: The majority of recent short DNA insertions in the human genome are tandem duplications. Mol Biol Evol 24: 1190–1197 (2007).
- Metzker ML: Sequencing technologies the next generation. Nat Rev Genet 11:31-46 (2010).
- Meuwissen THE, Hayes BJ, Goddard ME: Prediction of total genetic value using genome-wide dense marker maps. Genetics 157:1819–1829 (2001).
- Miao YW, Peng MS, Wu GS, Ouyang YN, Yang ZY, et al: Chicken domestication: an updated perspective based on mitochondrial genomes. Heredity 110:277–282 (2012).
- Miklos GLC: Localized highly repetitive DNA sequence in vertebrate and invertebrate genomes, in MacIntyre RJ (ed): Molecular Evolutionary Genetics, pp 241–321 (Plenum Press, New York 1985).
- Miller OJ: Heterochromatin visualized by various techniques, in Pfeiffer RA (ed): Modern Aspects of Cytogenetics: Constitutive Heterochromatin in Man, Symposia Medica Hoechst, Vol 6, pp 135–148 (Schattauer, Stuttgart 1973).
- Miller OJ, Schnedl W, Allen J, Erlanger BF: 5-Methylcytosine localised in mammalian constitutive heterochromatin. Nature 251: 636–637 (1974).
- Mindell DP, Sorenson MD, Dimcheff DE, Hasegawa M, Ast JC, Yuri T: Interordinal relationships of birds and other reptiles based on whole mitochondrial genomes. Syst Biol 48:138–152 (1999).
- Miniou P, Jeanpierre M, Blanquet V, Sibella V, Boneau D, et al: Abnormal methylation pattern in constitutive and facultative (X inactive chromosome) heterochromatin of ICF patients. Hum Mol Genet 3:2093–2102 (1994).
- Minkina A, Matson CK, Lindeman RE, Ghyselinck NB, Bardwell VJ, Zarkower D: DMRT1 protects male gonadal cells from retinoid-dependent sexual transdifferentiation. Dev Cell 29:511–520 (2014).
- modENCODE Consortium, Roy S, Ernst J, Kharchenko PV, Kheradpour P, et al: Identification of functional elements and regulatory circuits by *Drosophila* modENCODE. Science 330:1787–1797 (2010).
- Modi WS, Romanov M, Green ED, Ryder O: Molecular cytogenetics of the California condor: evolutionary and conservation implications. Cytogenet Genome Res 127:26–32 (2009).
- Moghadam HK, Pointer MA, Wright AE, Berlin S, Mank JE: W chromosome expression responds to female-specific selection. Proc Natl Acad Sci USA 109:8207–8211 (2012).
- Molnár J, Poti A, Pipek O, Krzystanek M, Kanu N, et al: The genome of the chicken DT40 bursal lymphoma cell line. G3 (Bethesda) 4: 2231–2240 (2014).
- Moniot B, Boizet-Bonhoure B, Poulat F: Male specific expression of lipocalin-type prostaglandin D synthase (cPTGDS) during chicken gonadal differentiation: relationship with cSOX9. Sex Dev 2:96–103 (2008).

- Montpellier C, Burgeois CA, Kokalj-Vokac N, Muleris M, Niveleau A, et al: Detection of methylcytosine-rich heterochromatin on banded chromosomes. Application to cells with various status of DNA methylation. Cancer Genet Cytogenet 78:87–93 (1994).
- Moreira GCM, Godoy TF, Boschiero C, Gheyas AA, Gasparin G, et al: Variant discovery in a QTL region on chromosome 3 associated with fatness in chickens. Anim Genet 46:141– 147 (2015).
- Morejohn GV: Breakdown of isolation mechanisms in two species of captive junglefowl (*Gallus gallus* and *Gallus sonneratii*). Evolution 22:576–582 (1968).
- Morgan RW, Sofer L, Anderson AS, Bernberg EL, Cui J, Burnside J: Induction of host gene expression following infection of chicken embryo fibroblasts with oncogenic Marek's disease virus. J Virol 75:533–539 (2001).
- Morisson M, Pitel F, Fillon V, Pouzadoux A, Bergé R, et al: Integration of chicken cytogenetic and genetic maps: 18 new polymorphic markers isolated from BAC and PAC clones. Anim Genet 29:348–355 (1998).
- Morisson M, Denis M, Milan D, Klopp C, Leroux S, et al: The chicken RH map: current state of progress and microchromosome mapping. Cytogenet Genome Res 117:14–21 (2007).
- Morrow C, Fehler F: Marek's disease: a worldwide problem, in Davison F, Nair V (eds): Marek's Disease, An Evolving Problem, pp 49–61 (Elsevier Academic Press, San Diego 2004).
- Motegi A, Takata M: Multiple genetic manipulations of DT40 cell line. Methods Mol Biol 1114:25–35 (2014).
- Mou C, Pitel F, Gourichon D, Vignoles F, Tzika A, et al: Cryptic patterning of avian skin confers a developmental facility for loss of neck feathering. PLoS Biol 9:e1001028 (2011).
- Muchadeyi FČ, Eding H, Simianer H, Wollny CBA, Groeneveld E, Weigend S: Mitochondrial DNA D-loop sequences suggest a Southeast Asian and Indian origin of Zimbabwean village chickens. Anim Genet 39:615–622 (2008).
- Muir WM, Wong GK, Zhang Y, Wang J, Groenen MA, et al: Genome-wide assessment of worldwide chicken SNP genetic diversity indicates significant absence of rare alleles in commercial breeds. Proc Natl Acad Sci USA 105: 17312–17317 (2008a).
- Muir WM, Wong GK, Zhang Y, Wang J, Groenen MAM, et al: Review of the initial validation and characterization of a 3K chicken SNP array. World's Poult Sci J 64:219–225 (2008b).
- Mullaney JM, Mills RE, Pittard WS, Devine SE: Small insertions and deletions (INDELs) in human genomes. Hum Mol Genet 19:R131– R136 (2010).
- Murnane JP: Telomere dysfunction and chromosome instability. Mutat Res 730:28–36 (2012).
- Murphy PJ, Cipriany BR, Wallin CB, Ju CY, Szeto K, et al: Single-molecule analysis of combinatorial epigenomic states in normal and tumor cells. Proc Natl Acad Sci USA 110:7772–7777 (2013).

- Muscarella DE, Vogt VM, Bloom SE: The ribosomal RNA gene cluster in aneuploid chickens: evidence for increased gene dosage and regulation of gene expression. J Cell Biol 101: 1749–1756 (1985).
- Mwacharo JM, Bjørnstad G, Mobegi V, Nomura K, Hanada H, et al: Mitochondrial DNA reveals multiple introductions of domestic chicken in East Africa. Mol Phylogenet Evol 58:374–382 (2011).
- Mwacharo JM, Bjørnstad G, Han JL, Hanotte O: The history of African village chickens: an archaeological and molecular perspective. Afr Archaeol Rev 30:97–114 (2013).
- Nachman MW: Single nucleotide polymorphisms and recombination rate in humans. Trends Genet 17:481–485 (2001).
- Nadachowska-Brzyska K, Burri R, Olason PI, Kawakami T, Smeds L, Ellegren H: Demographic divergence history of pied flycatcher and collared flycatcher inferred from wholegenome re-sequencing data. PLoS Genet 9:e1003942 (2013).
- Nadachowska-Brzyska K, Li C, Smeds L, Zhang G, Ellegren H: Temporal dynamics of avian populations during Pleistocene revealed by whole-genome sequences. Curr Biol 25:1375–1380 (2015).
- Nair V: Evolution of Marek's disease a paradigm for incessant race between the pathogen and the host. Vet J 170:175–183 (2005).
- Nakata T, Ishiguro M, Aduma N, Izumi H, Kuroiwa A: Chicken hemogen homolog is involved in the chicken-specific sex-determining mechanism. Proc Natl Acad Sci USA 110: 3417–3422 (2013).
- Nam K, Ellegren H: Recombination drives vertebrate genome contraction. PLoS Genet 8:e1002680 (2012).
- Nam K, Mugal C, Nabholz B, Schielzeth H, Wolf JB, et al: Molecular evolution of genes in avian genomes. Genome Biol 11:R68 (2010).
- Nanda I, Schrama D, Feichtinger W, Haaf T, Schartl M, Schmid M: Distribution of telomeric (TTAGGG)(n) sequences in avian chromosomes. Chromosoma 111:215–227 (2002).
- Nätt D, Agnvall B, Jensen P: Large sex differences in chicken behavior and brain gene expression coincide with few differences in promoter DNA-methylation. PLoS One 9:e96376 (2014).
- Nawrocki EP, Eddy SR: Infernal 1.1: 100-fold faster RNA homology searches. Bioinformatics 29:2933–2935 (2013).
- Necsulea A, Soumillon M, Warnefors M, Liechti A, Daish T, et al: The evolution of lncRNA repertoires and expression patterns in tetrapods. Nature 505:635–640 (2014).
- Neiman PE, Ruddell A, Jasoni C, Loring G, Thomas SJ, et al: Analysis of gene expression during *myc* oncogene-induced lymphomagenesis in the bursa of Fabricius. Proc Natl Acad Sci USA 98:6378–6383 (2001).

- Neiman PE, Burnside J, Elsaesser K, Hwang H, Clurman BE, et al: Analysis of gene expression, copy number and palindrome formation with a DT40 enriched cDNA microarray. Subcell Biochem 40:245–256 (2006).
- Neuman JA, Isakov O, Shomron N: Analysis of insertion-deletion from deep-sequencing data: software evaluation for optimal detection. Brief Bioinform 14:46–55 (2013).
- Newman LA, Robinson PR: Cone visual pigments of aquatic mammals. Vis Neurosci 22:873– 879 (2005).
- Ng CS, Wu P, Foley J, Foley A, McDonald ML, et al: The chicken frizzle feather is due to an α-keratin (*KRT75*) mutation that cuses a defective rachis. PLoS Genet 8:e1002748 (2012).
- Ng PC, Henikoff S: SIFT: predicting amino acid changes that affect protein function. Nucleic Acids Res 31:3812–3814 (2003).
- Nicol B, Yao HH: Building an ovary: insights into establishment of somatic cell lineages in the mouse. Sex Dev 8:243–251 (2014).
- Niikura M, Dodgson J, Cheng H: Direct evidence of host genome acquisition by the alphaherpesvirus Marek's disease virus. Arch Virol 151:537–549 (2006).
- Nishibori M, Shimogiri T, Hayashi T, Yasue H: Molecular evidence for hybridization of species in the genus *Gallus* except for *Gallus varius*. Anim Genet 36:367–375 (2005).
- Nishida-Umehara C, Tsuda Y, Ishijima J, Ando J, Fujiwara A, et al: The molecular basis of chromosomal orthologies and sex chromosomal differentiation in palaeognathus birds. Chromosome Res 15:721–734 (2007).
- Noorai RE, Freese NH, Wright LM, Chapman SC, Clark LA: Genome-wide association mapping and identification of candidate genes for the rumpless and ear-tufted traits of the Araucana chicken. PLoS One 7:e40974 (2012).
- O'Connor JK, Chiappe LM: A revision of enantiornithine (aves: Ornithothoraces) skull morphology. J Syst Palaeontol 9:135–157 (2011).
- Ogawa A, Murata K, Mizuno S: The location of Z- and W-linked marker genes and sequence on the homomorphic sex chromosomes of the ostrich and the emu. Proc Natl Acad Sci USA 95:4415–4418 (1998).
- O'Hare TH, Delany ME: Genetic variation exists for telomeric array organization within and among the genomes of normal, immortalized, and transformed chicken systems. Chromosome Res 17:947–964 (2009).
- Okano M, Bell DW, Haber DA, Li E: DNA methyltransferases Dnmt3a and Dnmt3b are essential for de novo methylation and mammalian development. Cell 99:247–257 (1999).
- Oleksyk TK, Pombert JF, Siu D, Mazo-Vargas A, Ramos B, et al: A locally funded Puerto Rican parrot (*Amazona vittata*) genome sequencing project increases avian data and advances young researcher education. Gigascience 1:14 (2012).

- Organ CL, Shedlock AM, Meade A, Pagel M, Edwards SV: Origin of avian genome size and structure in non-avian dinosaurs. Nature 446: 180–184 (2007).
- Osterrieder N, Kamil JP, Schumacher D, Tischer BK, Trapp S: Marek's disease virus: from miasma to model. Nat Rev Microbiol 4:283–294 (2006).
- Ottolenghi C, Pelosi E, Tran J, Colombino M, Douglass E, et al: Loss of *Wnt4* and *Foxl2* leads to female-to-male sex reversal extending to germ cells. Hum Mol Genet 16:2795–2804 (2007).
- Pacheco MA, Battistuzzi FU, Lentino M, Aguilar RF, Kumar S, Escalante AA: Evolution of modern birds revealed by mitogenomics: timing the radiation and origin of major orders. Mol Biol Evol 28:1927–1942 (2011).
- Pan Q, Shai O, Lee LJ, Frey BJ, Blencowe BJ: Deep surveying of alternative splicing complexity in the human transcriptome by high-throughput sequencing. Nat Genet 40:1413–1415 (2008).
- Pannetier M, Fabre S, Batista F, Kocer A, Renault L, et al: FOXL2 activates *P450 aromatase* gene transcription: towards a better characterization of the early steps of mammalian ovarian development. J Mol Endocrinol 36:399–413 (2006).
- Parcells MS, Lin SF, Dienglewicz RL, Majerciak V, Robinson DR, et al: Marek's disease virus (MDV) encodes an interleukin-8 homolog (vIL-8): characterization of the vIL-8 protein and a vIL-8 deletion mutant MDV. J Virol 75: 5159–5173 (2001).
- Park MN, Choi JA, Lee KT, Lee HJ, Choi BH, et al: Genome-wide association study of chicken plumage pigmentation. Asian-Australas J Anim Sci 26:1523–1528 (2013).
- Park TS, Han JY: Genetic modification of chicken germ cells. Ann N Y Acad Sci 1271:104–109 (2012).
- Park TS, Lee HC, Rengaraj D, Han JY: Germ cell, stem cell, and genomic modification in birds. J Stem Cell Res Ther 4:201 (2014a).
- Park TS, Lee HJ, Kim KH, Kim JS, Han JY: Targeted gene knockout in chickens mediated by TALENs. Proc Natl Acad Sci USA 111:12716– 12721 (2014b).
- Parker P, Li B, Li H, Wang J: The genome of Darwin's finch (*Geospiza fortis*). GigaScience, http://dx.doi.org/10.5524/100040 (2012).
- Pastinen T: Genome-wide allele-specific analysis: insights into regulatory variation. Nat Rev Genet 11:533–538 (2010).
- Pastor W, Aravind L, Rao A: TETonic shift: biological roles of TET proteins in DNA demethylation and transcription. Nat Rev Mol Cell Biol 14:341–356 (2013).
- Paton T, Haddrath O, Baker AJ: Complete mitochondrial DNA genome sequences show that modern birds are not descended from transitional shorebirds. Proc Biol Sci 269:839–846 (2002).

- Pazderka F, Longenecker BM, Law GR, Stone HA, Ruth RF: Histocompatibility of chicken populations selected for resistance to Marek's disease. Immunogenetics 2:93–100 (1975).
- Pereira SL, Baker AJ: Waterfowl and gamefowl (Galloanserae), in Hedges SB, Kumar S (eds): The Timetree of Life, pp 415–418 (Oxford University Press, New York 2009).
- Petherbridge L, Howes K, Baigent SJ, Sacco MA, Evans S, et al: Replication-competent bacterial artificial chromosomes of Marek's disease virus: novel tools for generation of molecularly defined herpesvirus vaccines. J Virol 77: 8712–8718 (2003).
- Petherbridge L, Brown AC, Baignet SJ, Howes K, Sacco MA, et al: Oncogenicity of virulent Marek's disease virus cloned as bacterial artificial chromosomes. J Virol 78:13376–13380 (2004).
- Pevzner P, Tesler G: Human and mouse genomic sequences reveal extensive breakpoint reuse in mammalian evolution. Proc Natl Acad Sci USA 100:7672–7677 (2003).
- Pisenti JM, Delany ME, Taylor RL, Abbott UK, Abplanalp H, et al: Avian genetic resources at risk: an assessment and proposal for conservation of genetic stocks in the USA and Canada (Report No. 20, Genetic Resources Conservation Program, Division of Agriculture and Natural Resources, University of California, Davis, CA 1999).
- Poelstra JW, Vijay N, Bossu CM, Lantz H, Ryll B, et al: The genomic landscape underlying phenotypic integrity in the face of gene flow in crows. Science 344:1410–1414 (2014).
- Pokorná M, Altmanová M, Kratochvíl L: Multiple sex chromosomes in the light of female meiotic drive in amniote vertebrates. Chromosome Res 22:35–44 (2014).
- Popescu NC, Zimonjic D, DiPaolo JA: Viral integration, fragile sites, and proto-oncogenes in human neoplasia. Hum Genet 84:383–386 (1990).
- Preisinger R: Genome-wide selection in poultry. Anim Prod Sci 52:121–125 (2012).
- Qanbari S, Hansen M, Weigend S, Preisinger R, Simianer H: Linkage disequilibrium reveals different demographic history in egg laying chickens. BMC Genet 11:103 (2010).
- Qanbari S, Strom TM, Haberer G, Weigend S, Gheyas AA, et al: A high resolution genomewide scan for significant selective sweeps: an application to pooled sequence data in laying chickens. PLoS One 7:e49525 (2012).
- Qin S, Chock PB: Implication of phosphatidylinositol 3-kinase membrane recruitment in hydrogen peroxide-induced activation of PI3K and Akt. Biochemistry 42:2995–3003 (2003).
- Qu Y, Zhao H, Han N, Zhou G, Song G, et al: Ground tit genome reveals avian adaptation to living at high altitudes in the Tibetan plateau. Nat Commun 4:2071 (2013).
- Quevillon E, Silventoinen V, Pillai S, Harte N, Mulder N, et al: InterProScan: protein domains identifier. Nucleic Acids Res 33:W116– W120 (2005).

- Raiber EA, Beraldi D, Ficz G, Burgess HE, Branco MR, et al: Genome-wide distribution of 5-formylcytosine in embryonic stem cells is associated with transcription and depends on thymine DNA glycosylase. Genome Biol 13:R69 (2012).
- Rainey MD, Zachos G, Gillespie DA: Analysing the DNA damage and replication checkpoints in DT40 cells. Subcell Biochem 40:107–117 (2006).
- Ran FA, Hsu PD, Wright J, Agarwala V, Scott DA, Zhang F: Genome engineering using the CRISPR-Cas9 system. Nat Protoc 8:2281– 2308 (2013).
- Rands CM, Darling A, Fujita M, Kong L, Webster MT, et al: Insights into the evolution of Darwin's finches from comparative analysis of the *Geospiza magnirostris* genome sequence. BMC Genomics 14:95 (2013).
- Rands CM, Meader S, Ponting CP, Lunter G: 8.2% of the human genome is constrained: variation in rates of turnover across functional element classes in the human lineage. PLoS Genet 10:e1004525 (2014).
- Rao M, Morisson M, Faraut T, Bardes S, Fève K, et al: A duck RH panel and its potential for assisting NGS genome assembly. BMC Genomics 13:513 (2012).
- Ratnakumar A, Mousset S, Glémin S, Berglund J, Galtier N, et al: Detecting positive selection within genomes: the problem of biased gene conversion. Philos Trans R Soc Lond B Biol Sci 365:2571–2580 (2010).
- Raymond CS, Kettlewell JR, Hirsch B, Bardwell VJ: Expression of *Dmrt1* in the genital ridge of mouse and chicken embryos suggests a role in vertebrate sexual development. Dev Biol 215:208–220 (1999).
- Razanfindraibe H, Mobegi VA, Ommeh S, Rakotondravao, ML, Bjørnstad G, et al: Mitochondrial DNA origin of indigenous Malagasy chicken. Ann N Y Acad Sci 1149:77–79 (2008).
- Redon R, Ishikawa S, Fitch KR, Feuk L, Perry GH, et al: Global variation in copy number in the human genome. Nature 444:444–454 (2006).
- Reed KM, Faile GM, Kreuth SB, Chaves LD, Sullivan LM: Association and in silico assignment of sequences from turkey BACs. Anim Biotechnol 19:80–83 (2008).
- Ren CW, Lee MK, Yan B, Ding K, Cox B, et al: A BAC-based physical map of the chicken genome. Genome Res 13:2754–2758 (2003).
- Reynaud C, Bruno C, Boullanger P, Grange J, Barbesti S, Niveleau A: Monitoring of urinary excretion of modified nucleosides in cancer patients using a set of six monoclonal antibodies. Cancer Lett 61:255–262 (1992).
- Rinn JL, Chang HY: Genome regulation by long noncoding RNAs. Annu Rev Biochem 81: 145–166 (2012).
- Rivera CM, Ren B: Mapping human epigenomes. Cell 155:39–55 (2013).
- Robb EA: Developmental Syndromes in the Chicken: Mapping and Candidate Gene Studies. Dissertation, University of California, Davis, CA (2012).

- Robb EA, Delany ME: The expression of preaxial polydactyly is influenced by modifying genetic elements and is not maintained by chromosomal inversion in an avian biomedical model. Cytogenet Genome Res 136:50-68 (2012a).
- Robb EA, Delany ME: Case study of sequence capture enrichment technology: identification of variation underpinning developmental syndromes in an amniote model. Genes 3: 233-247 (2012b).
- Robb EA, Gitter CL, Cheng HH, Delany ME: Chromosomal mapping and candidate gene discovery of chicken developmental mutants and genome-wide variation analysis of MHC congenics. J Hered 102:141-156 (2011).
- Robb EA, Antin PB, Delany ME: Defining the sequence elements and candidate genes for the coloboma mutation. Plos One 8:e60267 (2013).
- Roberts JA: Assessing the scavenging feed resources base for sustainable smallholder poultry development, in Sonaiya EB (ed): Sustainable Poultry Production in Africa. Proc. International Workshop, 13-16 June 1995, International Livestock Research Institute, Addis Ababa (1997).
- Robertson G, Schein J, Chiu R, Corbett R, Field M, et al: De novo assembly and analysis of RNA-seq data. Nat Methods 7:909-912 (2010).
- Robinson CM: Investigation of the Interactions between Marek's Disease Virus and the Chicken Host Genome throughout Pathogenesis. PhD Dissertation, University of California, Davis, CA (2013).
- Robinson CM, Hunt HD, Cheng HH, Delany ME: Chromosomal integration of an avian oncogenic herpesvirus reveals telomeric preferences and evidence for lymphoma clonality. Herpesviridae 1:5 (2010).
- Robinson CM, Cheng HH, Delany ME: Temporal kinetics of Marek's disease herpesvirus: integration occurs early after infection in both B and T cells. Cytogenet Genome Res 144:142-154 (2014).
- Rodríguez-León J, Rodríguez Esteban C, Martí M, Santiago-Josefat B, Dubova I, et al: Pitx2 regulates gonad morphogenesis. Proc Natl Acad Sci USA 105:11242-11247 (2008).
- Roeszler KN, Itman C, Sinclair AH, Smith CA: The long non-coding RNA, MHM, plays a role in chicken embryonic development, including gonadogenesis. Dev Biol 366:317-326 (2012).
- Romanov MN, Dodgson JB: Cross-species overgo hybridization and comparative physical mapping within avian genomes. Anim Genet 37: 397-399 (2006).
- Romanov MN, Weigend S: Genetic diversity in chicken populations based on microsatellite markers, in Dekkers JCM, Lamont SJ, Rothschild MF: From Jay Lush to Genomics: Visions for Animal Breeding and Genetics' [Conference Proceedings], p 174 (Iowa State University, Ames 1999).

- Romanov MN, Price JA, Dodgson JB: Integration of animal linkage and BAC contig maps using overgo hybridization. Cytogenet Genome Res 102:277-281 (2003).
- Romanov MN, Daniels LM, Dodgson JB, Delany ME: Integration of the cytogenetic and physical maps of chicken chromosome 17. Chromosome Res 13:215-222 (2005).
- Romanov MN, Koriabine M, Nefedov M, de Jong PJ, Ryder OA: Construction of a California condor BAC library and first-generation chicken-condor comparative physical map as an endangered species conservation genomics resource. Genomics 88:711-718 (2006).
- Romanov MN, Sazanov AA, Moiseyeva IG, Smirnov AF: Poultry, in Cockett NE, Kole C (eds): Genome Mapping and Genomics in Animals, Vol. 3: Genome Mapping and Genomics in Domestic Animals, pp 75-141 (Springer-Verlag, New York 2009a).
- Romanov MN, Tuttle EM, Houck ML, Modi WS, Chemnick LG, et al: The value of avian genomics to the conservation of wildlife. BMC Genomics 10 Suppl 2:S10 (2009b).
- Romanov MN, Dodgson JB, Gonser RA, Tuttle EM: Comparative BAC-based mapping in the white-throated sparrow, a novel behavioral genomics model, using interspecies overgo hybridization. BMC Res Notes 4:211 (2011).
- Romanov MN, Farré M, Lithgow PE, Fowler KE, Skinner BM, et al: Reconstruction of gross avian genome structure, organization and evolution suggests that the chicken lineage most closely resembles the dinosaur avian ancestor. BMC Genomics 15:1060 (2014).
- Roux PF, Boutin M, Desert C, Djari A, Esquerre D, et al: Re-sequencing data for refining candidate genes and polymorphisms in QTL regions affecting adiposity in chicken. PLoS One 9:e111299 (2014).
- Rubin CJ, Zody MC, Eriksson J, Meadows JRS, Sherwood E, et al: Whole-genome resequencing reveals loci under selection during chicken domestication. Nature 464:587-591 (2010)
- Ruiz-Herrera A, Farré M, Robinson TJ: Molecular cytogenetic and genomic insights into chromosomal evolution. Heredity 108:28-36 (2012).
- Saelao P, Zhou H: Tissue-specific transcriptome regulation through histone modifications in chickens. Plant and Animal Genome Meeting XXIII, San Diego, CA (2015).
- Sajan SA, Hawkins RD: Methods for identifying higher-order chromatin structure. Annu Rev Genomics Hum Genet 13:59-82 (2012).
- Sale JE, Ross AL, Simpson LJ: Analysis of DNA replication damage bypass and its role in immunoglobulin repertoire development. Subcell Biochem 40:271-294 (2006).
- Sander S, Calado DP, Srinivasan L, Köchert K, Zhang B, et al: Synergy between PI3K signaling and MYC in Burkitt lymphomagenesis. Cancer Cell 22:167-179 (2012).
- Sanjana NE, Cong L, Zhou Y, Cunniff MM, Feng G, Zhang F: A transcription activator-like effector toolbox for genome engineering. Nat Protoc 7:171-192 (2012).

- Sarda S, Hannenhalli S: Next-generation sequencing and epigenomics research: a hammer in search of nails. Genomics Inform 12: 2-11 (2014).
- Sarson A, Parvizi P, Lepp D, Quinton M, Sharif S: Transcriptional analysis of host responses to Marek's disease virus infection in genetically resistant and susceptible chickens. Anim Genet 39:232-240 (2008).
- Sawai H, Kim HL, Kuno K, Suzuki S, Gotoh H, et al: The origin and genetic variation of domestic chickens with special reference to junglefowls Gallus g. gallus and G. varius. PLoS One 5:e10639 (2010).
- Sawicki DL, Erlanger BF, Beiser SM: Immunochemical detection of minor bases in nucleic acids. Science 174:70-72 (1971).
- Scharffetter-Kochanek K, Lu H, Norman K, Van Nood N, Munoz F, et al: Spontaneous skin ulceration and defective T cell function in CD18 null mice. J Exp Med 188:119-131 (1998).
- Schattner P, Brooks AN, Lowe TM: The tRNAscan-SE, snoscan and snoGPS web servers for the detection of tRNAs and snoRNAs. Nucleic Acids Res 33:W686-W689 (2005).
- Schmid M, Enderle E, Schindler D, Schempp W: Chromosome banding and DNA replication patterns in bird karyotypes. Cytogenet Cell Genet 52:139-146 (1989).
- Schmid M, Nanda I, Guttenbach M, Steinlein C, Hoehn M, et al: First report on chicken genes and chromosomes 2000. Cytogenet Cell Genet 90:169-218 (2000).
- Schmid M, Nanda I, Hoehn H, Schartl M, Haaf T, et al: Second report on chicken genes and chromosomes 2005. Cytogenet Genome Res 109:415-479 (2005).
- Schmits R, Kündig T, Baker DM, Shumaker G, Simard J, et al: LFA-1-deficient mice show normal CTL responses to virus but fail to reject immunogenic tumor. J Exp Med 183: 1415-1426 (1996).
- Schnedl W, Dev VG, Tantravahi R, Miller DA, Erlanger BF, Miller OJ: 5-Methylcytosine in heterochromatic regions of chromosomes: chimpanzee and gorilla compared to the human. Chromosoma 52:59-66 (1975).
- Schnedl W, Erlanger BF, Miller OJ: 5-Methylcytosine in heterochromatic regions of chromosomes in Bovidae. Hum Genet 31:21-26 (1976).
- Schreck RR, Erlanger BF, Miller OJ: The use of antinucleoside antibodies to probe the organization of chromosomes denatured by ultraviolet irradiation. Exp Cell Res 88:31-39 (1974).
- Schreck RR, Erlanger BF, Miller OJ: Binding of anti-nucleoside antibodies reveals different classes of DNA in the chromosomes of the kangaroo rat (Dipodomys ordii). Exp Cell Res 108:403-411 (1977).
- Scrivener D: Rare Poultry Breeds (The Crowood Press, Ramsbury 2006).
- Scrivener D: Popular Poultry Breeds (The Crowood Press, Ramsbury 2009).

- Seabury CM, Dowd SE, Seabury PM, Raudsepp T, Brightsmith DJ, et al: A multi-platform draft de novo genome assembly and comparative analysis for the scarlet macaw (*Ara macao*). PLoS One 8:e62415 (2013).
- Sekido R, Lovell-Badge R: Mechanisms of gonadal morphogenesis are not conserved between chick and mouse. Dev Biol 302:132–142 (2007).
- Semple-Rowland SL, Lee NR, Van Hooser JP, Palczewski K, Baehr W: A null mutation in the photoreceptor guanylate cyclase gene causes the retinal degeneration chicken phenotype. Proc Natl Acad Sci USA 95:1271–1276 (1998).
- Shan Z, Nanda I, Wang Y, Schmid M, Vortkamp A, Haaf T: Sex-specific expression of an evolutionarily conserved male regulatory gene, *DMRT1*, in birds. Cytogenet Genome Res 89: 252–257 (2000).
- Shao P, Liao JY, Guan DG, Yang JH, Zheng LL, et al: Drastic expression change of transposonderived piRNA-like RNAs and microRNAs in early stages of chicken embryos implies a role in gastrulation. RNA Biol 9:212–227 (2012).
- Shapiro MD, Kronenberg Z, Li C, Domyan ET, Pan H, et al: Genomic diversity and evolution of the head crest in the rock pigeon. Science 339:1063–1067 (2013).
- Shedlock AM: Phylogenomic investigation of CR1 LINE diversity in reptiles. Syst Biol 55: 902–911 (2006).
- Shen Y, Yue F, McCleary DF, Ye Z, Edsall L, et al: A map of the *cis*-regulatory sequences in the mouse genome. Nature 488:116–120 (2012).
- Shepard JB, Gliga DA, Morrow AP, Hoffman S, Capehart AA: Versican knock-down compromises chondrogenesis in the embryonic chick limb. Anat Rec 291:19–27 (2008).
- Shetty S, Griffin DK, Graves JA: Comparative painting reveals strong chromosome homology over 80 million years of bird evolution. Chromosome Res 7:289–295 (1999).
- Shinomiya A, Kayashima Y, Kinoshita K, Mizutani M, Namikawa T, et al: Gene duplication of endothelin 3 is closely correlated with the hyperpigmentation of the internal organs (fibromelanosis) in silky chickens. Genetics 190: 627–638 (2012).
- Shiomi M, Ito T: The Watanabe heritable hyperlipidemic (WHHL) rabbit, its characteristics and history of development: a tribute to the late Dr. Yoshio Watanabe. Atherosclerosis 207:1–7 (2009).
- Sibley CG, Ahlquist JE: Phylogeny and Classification of Birds (Yale University Press, New Haven 1990).
- Sibley CG, Monroe BL Jr: Distribution and Taxonomy of Birds of the World (Yale University Press, New Haven 1990).
- Silva P, Guan X, Ho-Shing O, Jones J, Xu J, et al: Mitochondrial DNA-based analysis of genetic variation and relatedness among Sri Lankan indigenous chickens and the Ceylon junglefowl (*Gallus lafayetti*). Anim Genet 40:1–9 (2009).

- Silva RF, Dunn JR, Cheng HH, Niikura M: A MEQ-deleted Marek's disease virus cloned as a bacterial artificial chromosome is a highly efficacious vaccine. Avian Dis 54:862–869 (2010).
- Singer MF: Highly repeated sequences in mammalian genomes. Int Rev Cytol 76:67–112 (1982).
- Siwek M, Wragg D, Slawińska A, Malek M, Hanotte O, Mwacharo J: Insights into the genetic history of green-legged partridgelike fowl: mtDNA and genome-wide SNP analysis. Anim Genet 44:522–352 (2013).
- Skinner BM, Griffin DK: Intrachromosomal rearrangements in avian genome evolution: evidence for regions prone to breakpoints. Heredity 108:37–41 (2012).
- Skinner BM, Robertson LB, Tempest HG, Langley EJ, Ioannou D, et al: Comparative genomics in chicken and Pekin duck using FISH mapping and microarray analysis. BMC Genomics 10:357 (2009).
- Skinner BM, Al Mutery A, Smith D, Völker M, Hojjat N, et al: Global patterns of apparent copy number variation in birds revealed by cross-species comparative genomic hybridization. Chromosome Res 22:59–70 (2014).
- Slater GS, Birney E: Automated generation of heuristics for biological sequence comparison. BMC Bioinformatics 6:31 (2005).
- Smeds LA, Kawakami T, Burri R, Bolivar P, Husby A, et al: Genomic identification and characterization of the pseudoautosomal region in highly differentiated avian sex chromosomes. Nat Commun 5:5448 (2014).
- Smeds L, Warmuth V, Bolivar P, Uebbing S, Burri R, et al: Evolutionary analysis of the femalespecific avian W chromosome. Nat Commun 6:7330 (2015).
- Smith CA, Andrews J, Sinclair AH: Gonadal sex differentiation in chicken embryos: expression of estrogen receptor and aromatase genes. J Steroid Biochem Mol Biol 60:295–302 (1997).
- Smith CA, McClive P, Western P, Reed K, Sinclair AH: Conservation of a sex-determining gene. Nature 402:601–602 (1999).
- Smith CA, Clifford V, Western PS, Wilcox SA, Bell KS, Sinclair AH: Cloning and expression of a DAX1 homologue in the chicken embryo. J Mol Endocrinol 24:23–32 (2000).
- Smith CA, Shoemaker CM, Roeszler KN, Queen J, Crews D, Sinclair AH: Cloning and expression of R-Spondin1 in different vertebrates suggests a conserved role in ovarian development. BMC Dev Biol 8:72 (2008).
- Smith CA, Roeszler KN, Ohnesorg T, Cummins DM, Farlie PG, et al: The avian Z-linked gene DMRT1 is required for male sex determination in the chicken. Nature 461:267–271 (2009).
- Smith J, Bruley CK, Paton IR, Dunn I, Jones CT, et al: Differences in gene density on chicken macrochromosomes and microchromosomes. Anim Genet 31:96–103 (2000).

- Smith J, Sadeyen JR, Paton IR, Hocking PM, Salmon N, et al: Systems analysis of immune responses in Marek's disease virus-infected chickens identifies a gene involved in susceptibility and highlights a possible novel pathogenicity mechanism. J Virol 85:11146–11158 (2011).
- Smyth J: Genetics of plumage, skin and eye pigmentation in chickens, in Crawford RD (ed): Poultry Breeding and Genetics, pp 109–167 (Elsevier, Amsterdam 1990).
- Solinhac R, Leroux S, Galkina S, Chazara O, Feve K, et al: Integrative mapping analysis of chicken microchromosome 16 organization. BMC Genomics 11:616 (2010).
- Somes RG: Mutations and major variants of muscles and skeleton in chickens, in Crawford RD (ed): Poultry Breeding and Genetics, pp 209– 238 (Elsevier, Amsterdam 1990a).
- Somes RG: Lethal mutant traits in chickens, in Crawford RD (ed): Poultry Breeding and Genetics, pp 293–316 (Elsevier, Amsterdam 1990b).
- Song CX, Szulwach KE, Dai Q, Fu Y, Mao SQ, et al: Genome-wide profiling of 5-formylcytosine reveals its roles in epigenetic priming. Cell 153:678–691 (2013).
- Sonoda E, Sasaki MS, Buerstedde JM, Bezzubova O, Shinohara A, et al: Rad51-deficient vertebrate cells accumulate chromosomal breaks prior to cell death. EMBO J 17:598–608 (1998).
- Sonoda E, Hochegger H, Saberi A, Taniguchi Y, Takeda S: Differential usage of non-homologous end-joining and homologous recombination in double strand break repair. DNA Repair 5:1021–1029 (2006).
- Sorenson MD, Oneal E, Garcia-Moreno J, Mindell DP: More taxa, more characters: the hoatzin problem is still unresolved. Mol Biol Evol 20:1484–1498 (2003).
- Southern EM: DNA sequences and chromosome structure. J Cell Sci 1 (Suppl):31–41 (1984).
- Spillman WJ: Spurious allelomorphism: results of some recent investigations. Am Nat 42:610– 615 (1908).
- Springer MS, Cleven GC, Madsen O, de Jong WW, Waddell VG, et al: Endemic African mammals shake the phylogenetic tree. Nature 388:61–64 (1997).
- Springer TA, Dustin ML, Kishimoto TK, Marlin SD: The lymphocyte function-associated LFA-1, CD2, and LFA-3 molecules: cell adhesion receptors of the immune system. Annu Rev Immunol 5:223–252 (1987).
- Stamm S: Signals and their transduction pathways regulating alternative splicing: a new dimension of the human genome. Hum Mol Genet 11:2409–2416 (2002).
- Stanhope MJ, Waddell VG, Madsen O, de Jong W, Hedges SB, et al: Molecular evidence for multiple origins of Insectivora and for a new order of endemic African insectivore mammals. Proc Natl Acad Sci USA 95:9967–9972 (1998).
- Stern CD: The chick embryo past, present and future as a model system in developmental biology. Mech Dev 121:1011–1013 (2004).

- Stevens JG: Human herpesviruses: a consideration of the latent state. Microbiol Rev 53: 318-332 (1989).
- Storchová R, Divina P: Nonrandom representation of sex-biased genes on chicken Z chromosome. J Mol Evol 63:676–681 (2006).
- Storey AA, Athens JS, Bryant D, Carson M, Emery K, et al: Investigating the global dispersal of chickens in prehistory using ancient mitochondrial DNA signatures. PLoS One 7:e39171 (2012).
- Strahl BD, Allis CD: The language of covalent histone modifications. Nature 403:41–45 (2000).
- Strasburg GM, Chiang W: Pale, soft, exudative turkey – the role of ryanodine receptor variation in meat quality. Poult Sci 88:1497–1505 (2009).
- Sumner AT: A simple technique for demonstrating centromeric heterochromatin. Exp Cell Res 75:304–306 (1972).
- Sun L, Lamont SJ, Cooksey AM, McCarthy F, Tudor CO, et al: Transcriptome response to heat stress in a chicken hepatocellular carcinoma cell line. Cell Stress Chaperones, in press (2015).
- Sun YF, Zhao GP, Liu RR, Zheng MQ, Hu YD, et al: The identification of 14 new genes for meat quality traits in chicken using a genome-wide association study. BMC Genomics 14:458 (2013).
- Sundström H, Webster MT, Ellegren H: Reduced variation on the chicken Z chromosome. Genetics 167:377–385 (2004).
- Suzuki K, Kobayashi E, Yamashita H, Uenishi H, Churkina I, et al: Structural analysis of MHC alleles in an RSV tumour regression chicken using a BAC library. Anim Genet 43:348–351 (2012).
- Swanberg SE, Delany ME: Dynamics of telomere erosion in transformed and non-transformed avian cells in vitro. Cytogenet Genome Res 102:318–325 (2003).
- Tahiliani M, Koh KP, Shen Y, Pastor WA, Bandukwala H, et al: Conversion of 5-methylcytosine to 5-hydroxymethylcytosine in mammalian DNA by MLL partner TET1. Science 324:930–935 (2009).
- Takao N, Kato H, Mori R, Morrison C, Sonada E, et al: Disruption of ATM in p53-null cells causes multiple functional abnormalities in cellular response to ionizing radiation. Oncogene 18:7002–7009 (1999).
- Takata M, Sasaki MS, Sonoda E, Morrison C, Hashimoto M, et al: Homologous recombination and non-homologous end-joining pathways of DNA double-strand break repair have overlapping roles in the maintenance of chromosomal integrity in vertebrate cells. EMBO J 17:5497–5508 (1998).
- Takata M, Ishiai M, Kitao H: The Fanconi anemia pathway: insights from somatic cell genetics using DT40 cell line. Mutat Res 668:92–102 (2009).

- Tam W, Hughes SH, Hayward WS, Besmer P: Avian *bic*, a gene isolated from a common retroviral site in avian leukosis virus-induced lymphomas that encodes a noncoding RNA, cooperates with *c-myc* in lymphomagenesis and erythroleukemogenesis. J Virol 76:4275– 4286 (2002).
- Tan M, Luo H, Lee S, Jin F, Yang JS, et al: Identification of 67 histone marks and histone lysine crotonylation as a new type of histone modification. Cell 146:1016–1028 (2011).
- Tanikawa J, Ichikawa-Iwata E, Kanei-Ishii C, Nakai A, Matsuzawa S, et al: p53 suppresses the c-Myb-induced activation of heat shock transcription factor 3. J Biol Chem 275:15578– 15585 (2000).
- Teng G, Hakimpour P, Landgraf P, Rice A, Tuschl T, et al: MicroRNA-155 is a negative regulator of activation-induced cytidine deaminase. Immunity 28:621–629 (2008).
- Teranishi M, Shimada Y, Hori T, Nakabayashi O, Kikuchi T, et al: Transcripts of the MHM region on the chicken Z chromosome accumulate as non-coding RNA in the nucleus of female cells adjacent to the *DMRT1* locus. Chromosome Res 9:147–165 (2001).
- The FAANG Consortium: Coordinated international action to accelerate genome-to-phenome with FAANG, the Functional Annotation of Animal Genomes. Genome Biol 16:57 (2015).
- Thomas JW, Cáceres M, Lowman JJ, Morehouse CB, Short ME, et al: The chromosomal polymorphism linked to variation in social behavior in the white-throated sparrow (*Zonotrichia albicollis*) is a complex rearrangement and suppressor of recombination. Genetics 179:1455–1468 (2008).
- Thomson VA, Lebrasseur O, Austin JJ, Hunt TL, Burney DA, et al: Using ancient DNA to study the origins and dispersal of ancestral Polynesian chickens across the Pacific. Proc Natl Acad Sci USA 111:4826–4831 (2014).
- Tian F, Zhan F, VanderKraats ND, Hiken JF, Edwards JR, et al: DNMT gene expression and methylome in Marek's disease resistant and susceptible chickens prior to and following infection by MDV. Epigenetics 8:431–444 (2013).
- Tian M, Wang Y, Gu X, Feng C, Fang S, et al: Copy number variants in locally raised Chinese chicken genomes determined using array comparative genomic hybridization. BMC Genomics 14:262 (2013).
- Tian Z, Tolic N, Zhao R, Moore RJ, Hengel SM, et al: Enhanced top-down characterization of histone post-translational modifications. Genome Biol 13:R86 (2012).
- Tirian L, Hlavanda E, Olah J, Horvath I, Orosz F, et al: TPPP/p25 promotes tubulin assemblies and blocks mitotic spindle formation. Proce Natl Acad Sci USA 100:13976–13981 (2003).
- Tobita-Teramoto T, Jang GY, Kino K, Salter DW, Brumbaugh J, Akiyama T: Autosomal albino chicken mutation (c^a/c^a) deletes hexanucleotide ($-\Delta$ GACTGG817) at a copper-binding site of the tyrosinase gene. Poult Sci 79:46–50 (2000).

- Tokesi N, Lehotzky A, Horvath I, Szabo B, Olah J, et al: TPPP/p25 promotes tubulin acetylation by inhibiting histone deacetylase 6. J Biol Chem 285:17896–17906 (2010).
- Trapnell C, Williams BA, Pertea G, Mortazavi AM, Kwan G, et al: Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and isoform switching during cell differentiation. Nat Biotech 28:511– 515 (2010).
- Trapnell C, Hendrickson DG, Sauvageau M, Goff L, Rinn JL, Pachter L: Differential analysis of gene regulation at transcript resolution with RNA-seq. Nat Biotech 31:46–53 (2013).
- Trapp S, Parcells MS, Kamil JP, Schumacher D, Tischer BK, et al: A virus-encoded telomerase RNA promotes malignant T cell lymphomagenesis. J Exp Med 203:1307–1317 (2006).
- Tsompana M, Buck MJ: Chromatin accessibility: a window into the genome. Epigenetics Chromatin 7:33 (2014).
- Tudor CO, Schmidt CJ, Vijay-Shanker K: eGIFT: mining gene information from the literature. BMC Bioinform 11:418 (2010).
- Tulman ER, Afonso CL, Lu Z, Zsak L, Rock DL, Kutish GF: The genome of a very virulent Marek's disease virus. J Virol 74:7980–7988 (2000).
- Tummala H, Ali M, Getty P, Hocking PM, Burt DW, et al: Mutation in the guanine nucleotide-binding protein beta-3 causes retinal degeneration and embryonic mortality in chickens. Invest Ophthal Vis Sci 47:4714–4718 (2006).
- Twito T, Weigend S, Blum S, Granevitze Z, Feldman MW, et al: Biodiversity of 20 chicken breeds assessed by SNPs located in gene regions. Cytogenet Genome Res 117:319–326 (2007).
- Uetz P, Hošek J: The Reptile Database. http:// www.reptile-database.org, accessed Dec 8, 2014 (2014).
- Vaez M, Follett SA, Bed'hom B, Gourichon D, Tixier-Boichard M, Burke T: A single pointmutation within the melanophilin gene causes the *lavender* plumage colour dilution phenotype in the chicken. BMC Genetics 9:7 (2008).
- Vaillant S, Dorizzi M, Pieau C, Richard-Mercier N: Sex reversal and aromatase in chicken. J Exp Zool 290:727–740 (2001).
- Valbuena A, Lopez-Sanchez I, Lazo PA: Human VRK1 is an early response gene and its loss causes a block in cell cycle progression. PLoS One 3:e1642 (2008).
- Vallejo RL, Bacon LD, Liu HC, Witter RL, Groenen MA, et al: Genetic mapping of quantitative trait loci affecting susceptibility to Marek's disease virus induced tumors in F2 intercross chickens. Genetics 148:349–360 (1998).
- van de Lavoir MC, Diamond JH, Leighton PA, Mather-Love C, Heyer BS, et al: Germline transmission of genetically modified primordial germ cells. Nature 441:766–769 (2006).

- van Kooyk Y, Figdor CG: Avidity regulation of integrins: the driving force in leukocyte adhesion. Curr Opin Cell Biol 12:542–547 (2000).
- van Kooyk Y, Weder P, Heije K, Figdor CG: Extracellular Ca²⁺ modulates leukocyte function-associated antigen-1 cell surface distribution on T lymphocytes and consequently affects cell adhesion. J Cell Biol 124:1061– 1070 (1994).
- van Steensel B, Dekker J: Genomics tools for unraveling chromosome architecture. Nat Biotechnol 28:1089–1095 (2010).
- van Tuinen M: Birds (Aves), in Hedges SB, Kumar S (eds): The Timetree of Life, pp 409–411 (Oxford University Press, New York 2009).
- van Tuinen M, Hedges SB: Calibration of avian molecular clocks. Mol Biol Evol 18:206–213 (2001).
- van Tuinen M, Hedges SB: The effect of external and internal fossil calibrations on the avian evolutionary timescale. J Paleontol 78:45–50 (2004).
- van Tuinen M, Sibley CG, Hedges SB: Phylogeny and biogeography of ratite birds inferred from DNA sequences of the mitochondrial ribosomal genes. Mol Biol Evol 15:370–376 (1998).
- van Tuinen M, Sibley CG, Hedges SB: The early history of modern birds inferred from DNA sequences of nuclear and mitochondrial ribosomal genes. Mol Biol Evol 17:451–457 (2000).
- van Tuinen M, Butvill DB, Kirsch JA, Hedges SB: Convergence and divergence in the evolution of aquatic birds. Proc Biol Sci 268:1345–1350 (2001).
- Varricchio DJ, Jackson F, Borkowski JJ, Horner JR: Nest and egg clutches of the dinosaur *Troodon formosus* and the evolution of avian reproductive traits. Nature 385:247–250 (1997).
- Vasilikaki-Baker H, Nishioka Y: Immunological visualization of 5-methylcytosine-rich regions in Indian muntjac metaphase chromosomes. Exp Cell Res 147:226–230 (1983).
- Venkatesan RN, Price C: Telomerase expression in chickens: constitutive activity in somatic tissues and down-regulation in culture. Proc Natl Acad Sci USA 95:14763–14768 (1998).
- Verma RS: Heterochromatin. Molecular and Structural Aspects (Cambridge University Press, Cambridge 1988).
- Völker M, Backström N, Skinner BM, Langley EJ, Bunzey SK, et al: Copy number variation, chromosome rearrangement, and their association with recombination during avian evolution. Genome Res 20:503–511 (2010).
- Wakasugi M, Matsuura K, Nagasawa A, Fu D, Shimizu H, et al: *DDB1* gene disruption causes a severe growth defect and apoptosis in chicken DT40 cells. Biochem Biophys Res Commun 364:771–777 (2007).
- Wallis JW, Aerts J, Groenen M, Crooijmans R, Layman D, et al: A physical map of the chicken genome. Nature 432:761–764 (2004).

- Wang B, Ekblom R, Bunikis I, Siitari H, Hoglund J: Whole genome sequencing of the black grouse (*Tetrao tetrix*): reference guided assembly suggests faster-Z and MHC evolution. BMC Genomics 15:180 (2014).
- Wang DS, Kobayashi T, Zhou LY, Paul-Prasanth B, Ijiri S, et al: Foxl2 up-regulates aromatase gene transcription in a female-specific manner by binding to the promoter as well as interacting with Ad4 binding protein/steroidogenic factor 1. Mol Endocrinol 21:712–725 (2007).
- Wang GS, Cooper TA: Splicing in disease: disruption of the splicing code and the decoding machinery. Nat Rev Genet 8:749–761 (2007).
- Wang H, Lim D, Rudd CE: Immunopathologies linked to integrin signalling. Semin Immunopathol 32:173–182 (2010).
- Wang J, He XM, Ruan J, Dai MT, Chen J, et al: ChickVD: a sequence variation database for the chicken genome. Nucleic Acids Res 33:D438–D441 (2005).
- Wang K, Li M, Hadley D, Liu R, Glessner J, et al: PennCNV: an integrated hidden Markov model designed for high-resolution copy number variation detection in whole-genome SNP genotyping data. Genome Res 17:1665– 1674 (2007).
- Wang X, Nahashon S, Feaster T, Bohannon-Stewart A, Adefope N: An initial map of chromosomal segmental copy number variations in the chicken. BMC Genomics 11:351 (2010).
- Wang Y, Gu X, Feng C, Song C, Hu X, Li N: A genome-wide survey of copy number variation regions in various chicken breeds by array comparative genomic hybridization method. Anim Genet 43:282–289 (2012).
- Wang Z, Qu L, Yao J, Yang X, Li G, et al: An EAV-HP insertion in 5' flanking region of SLCO1B3 causes blue eggshell in the chicken. PLoS Genet 9:e1003183 (2013).
- Warren WC, Clayton DF, Ellegren H, Arnold AP, Hillier LW, et al: The genome of a songbird. Nature 464:757–762 (2010).
- Weber C, Boussau B, Romiguier J, Jarvis E, Ellegren H: Evidence for GC-biased gene conversion as a driver of between-lineage differences in avian base composition. Genome Biol 15:549 (2014a).
- Weber C, Nabholz B, Romiguier J, Ellegren H: K_r/K_c but not d_N/d_S correlates positively with body mass in birds, raising implications for inferring lineage-specific selection. Genome Biol 15:542 (2014b).
- Wei C, Skopp R, Takata M, Takeda S, Price CM: Effects of double-strand break repair proteins on vertebrate telomere structure. Nucleic Acids Res 30:2862–2870 (2002).
- Wells KL, Hadad Y, Ben-Avraham D, Hillel J, Cahaner A, Headon DJ: Genome-wide SNP scan of pooled DNA reveals nonsense mutation in *FGF20* in the scaleless line of featherless chickens. BMC Genomics 13:257 (2012).
- Werbajh S, Nojek I, Lanz R, Costas MA: RAC-3 is a NF-κB coactivator. FEBS Lett 485:195–199 (2000).

- Whitney O, Pfenning AR, Howard JT, Blatti CA, Liu F, et al: Core and region-enriched networks of behaviorally regulated genes and the singing genome. Science 346:1256780 (2014).
- Wiseman J: A History of the Britisch Pig (Duckworth, London 1986).
- Witke W: The role of profilin complexes in cell motility and other cellular processes. Trends Cell Biol 14:461–469 (2004).
- Witter RL: A new strategy for Marek's disease immunization – bivalent vaccine. Avian Pathol 13:133–135 (1984).
- Witter RL: Increased virulence of Marek's disease virus field isolates. Avian Dis 41:149–163 (1997).
- Wolc A, Arango J, Jankowski T, Settar P, Fulton JE, et al: Genome-wide association study for Marek's disease mortality in layer chickens. Avian Dis 57:395–400 (2013).
- Wolc A, Arango J, Jankowski T, Dunn I, Settar P, et al: Genome-wide association study for egg production and quality in layer chickens. J Anim Breed Genet 131:173–182 (2014).
- Wong GKS, Liu B, Wang J, Zhang Y, Yang X, et al: A genetic variation map for chicken with 2.8 million single-nucleotide polymorphisms. Nature 432:717–722 (2004).
- Woolfe A, Goodson M, Goode DK, Snell P, Mc-Ewen GK, et al: Highly conserved non-coding sequences are associated with vertebrate development. PLoS Biol 3:e7 (2005).
- Wragg D, Mwacharo JM, Alcalde JA, Hocking PM, Hanotte O: Analysis of genome-wide structure, diversity and fine mapping of Mendelian traits in traditional and village chickens. Heredity 109:6–18 (2012).
- Wragg D, Mwacharo JM, Alcalde JA, Wang C, Han JL, et al: Endogenous retrovirus EAV-HP linked to blue egg phenotype in Mapuche fowl. PLoS One 8:e71393 (2013).
- Wright D, Boije H, Meadows JRS, Bed'hom B, Gourichon D, et al: Copy number variation in intron 1 of SOX5 causes the pea-comb phenotype in chickens. PLoS Genet 5:e1000512 (2009).
- Xie L, Luo CL, Zhang CG, Zhang R, Tang J, et al: Genome-wide association study identified a narrow chromosome 1 region associated with chicken growth traits. PLoS One 7:e30910 (2012).
- Xu X, Zhou ZH, Wang XL, Kuang XW, Zhang FC, Du XK: Four-winged dinosaurs from China. Nature 421:335–340 (2003).
- Yamamoto KN, Kobayashi S, Tsuda M, Kurumizaka H, Takata M, et al: Involvement of SLX4 in interstrand cross-link repair is regulated by the Fanconi anemia pathway. Proc Natl Acad Sci USA 108:6492–6496 (2011).
- Yan Y, Yi G, Sun C, Qu L, Yang N: Genome-wide characterization of insertion and deletion variation in chicken using next generation sequencing. PLoS One 9:e104652 (2014).
- Yao Y, Charlesworth J, Nair V, Watson M: MicroRNA expression profiles in avian haemopoietic cells. Front Genet 4:153 (2013).

- Yasuda T, Maeda A, Kurosaki M, Tezuka T, Hironaka K, et al: Cbl suppresses B cell receptormediated phospholipase C (PLC)- γ 2 activation by regulating B cell linker protein-PLC- γ 2 binding. J Exp Med 191:641–650 (2000).
- Ye L, Hillier LW, Minx P, Thane N, Locke DP, et al: A vertebrate case study of the quality of assemblies derived from next-generation sequences. Genome Biol 12:R31 (2011).
- Ye Q, He K, Wu SY, Wan QH: Isolation of a 97-kb minimal essential MHC B locus from a new reverse-4D BAC library of the golden pheasant. PLoS One 7:e32154 (2012).
- Yi G, Qu L, Liu J, Yan Y, Xu G, Yang N: Genomewide patterns of copy number variation in the diversified chicken genomes using next-generation sequencing. BMC Genomics 15:962 (2014).
- Yokomine T, Kuroiwa A, Tanaka K, Tsudzuki M, Matsuda Y, Sasaki H: Sequence polymorphisms, allelic expression status and chromosome locations of the chicken *IGF2* and *MPR1* genes. Cytogenet Cell Genet 93:109–113 (2001).
- Yokomine T, Shirohzu H, Purbowasito W, Toyoda A, Iwama H, et al: Structural and functional analysis of a 0.5-Mb chicken region orthologous to the imprinted mammalian *Ascl2/ Mash2-Igf2-H19* region. Genome Res 15:154– 165 (2005).
- Yonash N, Bacon L, Witter R, Cheng H: High resolution mapping and identification of new quantitative trait loci (QTL) affecting susceptibility to Marek's disease. Anim Genet 30: 126–135 (1999).
- Young MD, Wakefield MJ, Smyth GK, Oshlack A: Gene ontology analysis for RNA-seq: accounting for selection bias. Genome Biol 11:R14 (2010).
- Yu M, Hon GC, Szulwach KE, Song CX, Zhang L, et al: Base-resolution analysis of 5-hydroxymethylcytosine in the mammalian genome. Cell 149:1368–1380 (2012).
- Yu Y, Zhang H, Tian F, Zhang W, Fang H, Song J: An integrated epigenetic and genetic analysis of DNA methyltransferase genes (*DNMTs*) in tumor resistant and susceptible chicken lines. PLoS One 3:e2672 (2008).

- Yuan X, Zhang M, Ruan W, Song C, Ren L, et al: Construction and characterization of a duck bacterial artificial chromosome library. Anim Genet 37:599–600 (2006).
- Yuri T, Mindell DP: Molecular phylogenetic analysis of Fringillidae, 'New World nine-primaried oscines' (Aves: Passeriformes). Mol Phylogenet Evol 23:229–243 (2002).
- Zhan X, Pan S, Wang J, Dixon A, He J, et al: Peregrine and saker falcon genome sequences provide insights into evolution of a predatory lifestyle. Nat Genet 45:563–566 (2013).
- Zhang G, Parker P, Li B, Li H, Wang J: The genome of Darwin's finch (*Geospiza fortis*). GigaScience. http://dx.doi.org/10.5524/ 100040 (2012).
- Zhang G, Cowled C, Shi Z, Huang Z, Bishop-Lilly KA, et al: Comparative analysis of bat genomes provides insight into the evolution of flight and immunity. Science 339:456–460 (2013).
- Zhang G, Li B, Li C, Gilbert MT, Jarvis ED, Wang J: Comparative genomic data of the avian phylogenomics project. Gigascience 3:26 (2014a).
- Zhang G, Li C, Li Q, Li B, Larkin DM, et al: Comparative genomics reveals insights into avian genome evolution and adaptation. Science 346:1311–1320 (2014b).
- Zhang H, Wang ZP, Wang SZ, Li H: Progress of genome wide association study in domestic animals. J Anim Sci Biotech 3:26 (2012).
- Zhang H, Du ZQ, Dong JQ, Wang HX, Shi HY, et al: Detection of genome-wide copy number variation in two chicken lines divergently selected for abdominal fat content. BMC Genomics 15:517 (2014).
- Zhang XH, Chasin LA: Computational definition of sequence motifs governing constitutive exon splicing. Genes Dev 18:1241–1250 (2004).
- Zhang X, Moor AN, Merkler KA, Liu Q, McLean MP: Regulation of alternative splicing of liver scavenger receptor class B gene by estrogen and the involved regulatory splicing factors. Endocrinology 148:5295–5304 (2007).
- Zhang Y, Wang J, Huang S, Zhu X, Liu J, et al: Systematic identification and characterization of chicken (*Gallus gallus*) ncRNAs. Nucleic Acids Res 37:6562–6574 (2009).

- Zhang Y, Zhang X, O'Hare TH, Payne WS, Dong JJ, et al: A comparative physical map reveals the pattern of chromosomal evolution between the turkey (*Meleagris gallopavo*) and chicken (*Gallus gallus*) genomes. BMC Genomics 12:447 (2011).
- Zhang Z, Pal S, Bi Y, Tchou J, Davuluri RV: Isoform level expression profiles provide better cancer signatures than gene level expression profiles. Genome Med 5:33 (2013).
- Zhao D, McBride D, Nandi S, McQueen HA, Mc-Grew MJ, et al: Somatic sex identity is cell autonomous in the chicken. Nature 464:237– 242 (2010).
- Zhao H, Bourque G: Recovering genome rearrangements in the mammalian phylogeny. Genome Res 19:934–942 (2009).
- Zhao Z, Tavoosidana G, Sjölinder M, Göndör A, Mariano P, et al: Circular chromosome conformation capture (4C) uncovers extensive networks of epigenetically regulated intraand interchromosomal interactions. Nat Genet 38:1341–1347 (2006).
- Zhou Q, Zhang J, Bachtrog D, An N, Huang Q, et al: Complex evolutionary trajectories of sex chromosomes across bird taxa. Science 346: 1246338 (2014).
- Zhou W, Liu R, Zhang J, Zheng M, Li P, et al: A genome-wide detection of copy number variation using SNP genotyping arrays in Beijing-You chickens. Genetica 142:441–450 (2014).
- Zimmer R, Verrinder Gibbins AM: Construction and characterization of a large-fragment chicken bacterial artificial chromosome library. Genomics 42:217–226 (1997).
- Zoorob R, Billault A, Severac V, Fillon V, Vignal A, Auffray C: Two chicken genomic libraries in the PAC and BAC cloning systems: organization and characterization. Anim Genet 27 Suppl 2:69 (1996).
- Zwaenepoel I, Mustapha M, Leibovici M, Verpy E, Goodyear R, et al: Otoancorin, an inner ear protein restricted to the interface between the apical surface of sensory epithelia and their overlying acellular gels, is defective in autosomal recessive deafness DFNB22. Proc Natl Acad Sci USA 99:6240–6245 (2002).